

Geographic distribution.—Redding Formation, Hooten Gulch Mudstone Member (Haggart, 1986), Swede basin, Redding area, Shasta County, California.

Etymology.—For W. P. Popeno, who collected the holotype, in recognition of his study of aporrhaides of the Redding area.

ACKNOWLEDGMENTS

The authors acknowledge Peter Rodda (California Academy of Sciences), Peter Ward and Ron Eng (Burke Memorial Washington State Museum), Dick Squires (California State University, Northridge), and Jean Dougherty (Geological Survey of Canada) for assistance in the procurement and study of specimens residing in collections at those institutions. The paper benefited from critical review by and valuable suggestions of Warren Allmon, Bonnie Murchey, Bill Sliter, and two anonymous reviewers. The help of Ann Marie Davis in printing some of the photographs is appreciated.

REFERENCES

- ABBOTT, R. T. 1960. The genus *Strombus* in the Indo-Pacific. Indo-Pacific Mollusca 1: 33–62.
- ANDERSON, F. M. 1902. Cretaceous deposits of the Pacific Coast. California Academy of Sciences Proceedings, Series 3, 2, 143 p.
- . 1938. Lower Cretaceous deposits in California and Oregon. Geological Society of America Special Paper 16, 339 p.
- . 1958. Upper Cretaceous of the Pacific Coast. Geological Society of America Memoir 71, 378 p.
- BANNON, J. L., D. J. BOTTJER, S. P. LUND, AND L. R. SAUL. 1989. Campanian/Maastrichtian stage boundary in southern California: Resolution and implications for large-scale depositional patterns. Geology, 17: 80–83.
- BOLTON, T. E. 1965. Catalogue of type invertebrate fossils of the Geological Survey of Canada. Volume II. Geological Survey of Canada, 344 p.
- . 1968. Catalogue of type invertebrate fossils of the Geological Survey of Canada. Volume IV. Geological Survey of Canada, 221 p.
- CONRAD, T. A. 1860. Descriptions of new species of Cretaceous and Eocene fossils of Mississippi and Alabama. Journal of the Philadelphia Academy of Natural Sciences, Series 2, 4: 275–298.
- COLBURN, I. P., L. R. SAUL, AND A. A. ALMGREN. 1981. Chatsworth Formation: a new formation name for the Upper Cretaceous strata of the Simi Hills, California, p. 9–16. In M. H. Link, R. L. Squires, and I. P. Colburn (eds.), Simi Hills Cretaceous Turbidites, Southern California. Society of Economic Paleontologists and Mineralogists, Pacific Section, Volume and Guidebook.
- COSSMANN, M. 1904. Essais de Paléonconchologie Comparée. Paris, Volume 6, 151 p.
- DAVIES, A. M. [revised by F. E. Eames] 1971. Tertiary Faunas. Volume 1. The composition of the faunas. George Allen and Unwin Limited, London, 57 p.
- DAVIS, C. H. 1913. New species from the Santa Lucia Mountains, California, with a discussion of the Jurassic age of the slates at Slate's Springs. Journal of Geology, 21: 435–458.
- DOCKERY, D. T., III. 1993. The Streptoneuran gastropods, exclusive of the Stenoglossa, of the Coffee Sand (Campanian) of northeastern Mississippi. Mississippi Department of Environmental Quality, Office of Geology, Bulletin 129, 191 p.
- ELDER, W. P., AND L. R. SAUL. 1993. Paleogeographic implications of molluscan assemblages in the Upper Cretaceous (Campanian) Pigeon Point Formation, California, p. 171–185. In G. Dunne, and K. McDougall (eds.), Mesozoic Paleogeography of the Western United States—II. Society of Economic Paleontologists and Mineralogists, Pacific Section, Book 71.
- GABB, W. M. 1864. Description of the Cretaceous fossils. California Geological Survey, Palaeontology, 1: 57–243, plates 9–32, 1865.
- . 1868. An attempt at a revision of the two families, Strombidae and Aporrhaidae. American Journal of Conchology 4: 137–149.
- . 1869. Cretaceous and Tertiary fossils. California Geological Survey, Palaeontology vol. 2, 299 p.
- GOLDFUSS, G. A. 1826–1844. Petrefacta Germaniae, &c. Petrefacta Musei Universitatis Aggildungen und Beschreibungen der Petrefacten Deutschlands und der angrenzenden Länder . . . herausgegeben von A. Goldfuss. Düsseldorf. Volume 1, 1826–1833, 252 p.; Volume 2, 1833–1840, 165 p.; Volume 3, 1844, 128 p.
- GRABAU, A. W., AND H. W. SHIMER. 1909. North American index fossils. A. G. Seiler and Company, New York, Volume 1, 853 p.
- GRADSTEIN, F. M., F. P. AGTERBERG, J. G. OGG, J. HARDENBOL, P. VAN VEEN, J. THIERRY, AND Z. HUANG. 1994. A Mesozoic time scale. Journal of Geophysical Research, series B, 99: 24,051–24,074.
- GRAY, J. E. 1850. Systematic arrangement of the figures, p. 63–124. In M. E. Gray, Figures of molluscous animals selected from various authors; etched for the use of students. Long, Brown, etc., London, Volume 4, 219 p.
- HAGGART, J. W. 1986. Stratigraphy of the Redding Formation of north-central California and its bearing on Late Cretaceous paleogeography, p. 161–178. In P. L. Abbott (ed.), Cretaceous stratigraphy western North America. Society of Economic Paleontologists and Mineralogists, Pacific Section, Book 46.
- . AND P. D. WARD. 1984. Late Cretaceous (Santonian-Campanian) stratigraphy of the northern Sacramento Valley, California. Geological Society of America Bulletin, 95: 618–627.
- . AND —. 1989. New Nanaimo Group ammonites (Cretaceous, Santonian-Campanian) from British Columbia and Washington State. Journal of Paleontology, 63: 218–227.
- KIRBY, M. X. 1991. Macropaleontology and biostratigraphy across the Cretaceous/Tertiary boundary, San Francisquito Formation, Warm Springs Mountain, Los Angeles County, southern California. Unpublished M.A. thesis, California State University, Northridge, 134 p.
- . AND L. R. SAUL. 1995. The Tethyan bivalve *Roudairia* from the Upper Cretaceous of California. Palaeontology, 38: 23–38.
- LINNAEUS, CARL. 1758. Systema naturae per regna tria naturae. Editio decima, reformata. Stockholm, Volume 1, Regnum animale, 824 p.
- LOCH, J. D. 1989. A new genus of aporrhaid gastropod from southern California. Journal of Paleontology, 63: 574–577.
- MATSUMOTO, T. 1960. Upper Cretaceous ammonites of California, Part III. Kyushu University, Faculty of Science Memoirs, Series D, Geology, Special Volume 2, 204 p.
- MULLER, J. E., AND J. A. JELETZKY. 1970. Geology of the Upper Cretaceous Nanaimo Group, Vancouver Island and Gulf Islands, British Columbia. Geological Survey of Canada, Paper 69–25, 77 p.
- OBRADOVICH, J. D. 1993. A Cretaceous time scale, p. 379–396. In W. G. E. Caldwell, and E. G. Kauffman (eds.), Evolution of the Western Interior Basin. Geological Association of Canada Special Paper 39.
- OLSSON, A. A. 1944. Contributions to the paleontology of northern Peru: Part VII. The Cretaceous of the Paita region. Bulletins of American Paleontology, 111, 114 p.
- D'ORBIGNY, A. 1842–1847. Description des animaux invertébrés; 2 Gastéropodes. Paléontologie française, terrain Crétacé, Series 1, 2, 456 p.
- POPENO, W. P., AND L. R. SAUL. 1987. Evolution and classification of the Late Cretaceous-early Tertiary gastropod *Perissitys*. Los Angeles County Museum of Natural History, Contributions in Science 380, 37 p.
- . —. AND T. SUSUKI. 1987. Gyrodiform gastropods from the Pacific Coast Cretaceous and Paleocene. Journal of Paleontology 61: 70–100.
- RAFINESQUE, C. S. 1815. Analyses de la nature ou tableau de l'univers et des corps organisés. Palermo, 224 p.
- ROY, K. 1994. Effects of the Mesozoic marine revolution on the taxonomic, morphologic, and biogeographic evolution of a group: aporrhaid gastropods during the Mesozoic. Paleobiology, 20: 274–296.
- SAUL, L. R. 1983. *Turritella* zonation across the Cretaceous-Tertiary boundary, California. University of California Publications, Geological Sciences 125, 164 p.
- . 1988. New Late Cretaceous and Early Tertiary Perissityidae (Gastropoda) from the Pacific Slope of North America. Los Angeles County Natural History Museum, Contributions in Science 400, 25 p.
- . AND W. P. POPENO. 1992. Pacific Slope Cretaceous bivalves of the genus *Calva*. Los Angeles County Natural History Museum, Contributions in Science 433, 68 p.
- . AND —. 1993. Additions to Pacific Slope Turonian Gastropoda. The Veliger 36: 351–388.

- SHIMER, H. W., AND SHROCK, R. R. 1944. Index fossils of North America. John Wiley and Sons, Inc., New York, 837 p.
- SHUMARD, B. F. 1859. Descriptions of new fossils from the Tertiary formations of Oregon and Washington Territories and the Cretaceous of Vancouver's Island, collected by Dr. Jno. Ivans, U. S. geologist, under instructions from the Department of the Interior. Transactions of the Academy of Natural Sciences of Saint Louis, 1: 120-125.
- SOHL, N. F. 1960. Archaeogastropoda, Mesogastropoda, and stratigraphy of the Ripley, Owl Creek, and Prairie Bluff Formations. U. S. Geological Survey Professional Paper 331-A, 152 p.
- . 1977. Utility of gastropods in biostratigraphy, p. 519-539. In E. G. Kauffman, and J. E. Hazel (eds.), Concepts and methods of biostratigraphy. Dowden Hutchinson, and Ross, Inc., Stroudsburg, Pennsylvania.
- STEWART, R. B. 1927. Gabb's California fossil type gastropods. Proceedings of the Academy of Natural Sciences of Philadelphia, 78 [1926]: 287-447.
- SUNDBERG, F. A., AND B. RINEY. 1984. Preliminary report on the Upper Cretaceous macro-invertebrate faunas near Carlsbad, California, p. 103-107. In P. L. Abbott (ed.), Upper Cretaceous depositional systems, southern California-northern Baja California. Society of Economic Paleontologists and Mineralogists, Pacific Section, Guidebook.
- TAFF, H. A., G. D. HANNA, AND C. M. CROSS. 1940. Type locality of the Cretaceous Chico Formation. Geological Society of America Bulletin 51: 1311-1328.
- TALIAFERRO, N. L. 1944. Cretaceous and Paleocene of Santa Lucia Range, California. American Association of Petroleum Geologists Bulletin, 28: 449-521.
- VEROSUB, K. L., J. W. HAGGART, AND P. D. WARD. 1989. Magnetostratigraphy of Upper Cretaceous strata of the Sacramento Valley, California. Geological Society of America Bulletin, 101: 521-533.
- WADE, B. 1926. The fauna of the Ripley Formation on Coon Creek, Tennessee. U. S. Geological Survey Professional Paper 137, 272 p.
- WARD, P. D. 1978. Revisions to the stratigraphy and biochronology of the Upper Cretaceous Nanaimo Group, British Columbia and Washington State. Canadian Journal of Earth Sciences, 15: 405-423.
- WEBSTER, M. L. 1983. New species of *Xenophora* and *Anchura* (Mollusca: Gastropoda) from the Cretaceous of Baja California Norte, Mexico. Journal of Paleontology, 57: 1050-1097.
- WHITE, C. A. 1879. Contributions to invertebrate paleontology, no. 1: Cretaceous fossils of the western states and territories. U. S. Geological and Geographical Survey of the Territories (Hayden Survey), Annual Report 11: 273-320.
- WHITEAVES, J. F. 1879. On the fossils of the Cretaceous rocks of Vancouver and adjacent Islands in the Strait of Georgia. Canada Geological Survey, Mesozoic Fossils, 1 (2): 93-190.
- . 1903. On some additional fossils from the Vancouver Cretaceous, with a revised list of species therefrom. Canada Geological Survey, Mesozoic Fossils, 1 (5): 309-415.

ACCEPTED 23 OCTOBER 1995

CITED LOCALITIES

- 92 CIT (=LACMIP 10100): Concretions in shale 100' above stream and near fence on N side of Harding Canyon, about 0.25 mi N of road fork in Santiago Canyon at Harding/Modjeska Canyon junction, near section line NW 1/4, NW 1/4 sec. 28, T5S, R7W, Santiago Peak quadrangle, Santa Ana Mountains, Orange County, California. Collected by: B. N. Moore, 1928. Ladd Formation, basal Holz Shale Member; late Turonian.
- 974 CIT (=LACMIP 10105): SW slope of Aliso-Santiago Creek divide, 475'N, 1200'W of SE corner of sec. 29 and edge of topo sheet, El Toro quadrangle, Santa Ana Mountains, Orange County, California. Collected by: W. P. Popenoe, 1/14/33. Williams Formation, Pleasants Sandstone Member; late middle to early late Campanian, *Metaplacenticeras pacificum* Zone (Matsumoto, 1960, p. 99; Popenoe and Saul, 1987, p. 34; Saul and Popenoe, 1992, p. 60).
- 1018 CIT (=LACMIP 10833): Fossiliferous layers cropping out in beds of small gullies in field along Durham-Pentz Rd., approximately 0.75 mi W of Pentz, approximately 950'S, 350'E of NW corner sec. 25, T1N, R3E, W side Messila Valley, Cherokee quadrangle, Butte County, California. Collected by: W. P. Popenoe and D. Scharf, 8/19/1931. Chico Formation; late early Campanian.
- 1053 CIT (=LACMIP 10093): First prominent NE-SW spur north of Santiago Creek near junction with Modjeska Creek, 200'N, 2850'E of SW corner sec. 20, T5S, R7W, El Toro quadrangle, Santa Ana Mountains, Orange County, California. Collected by: W. P. Popenoe, 4/9/34. Ladd Formation, upper part of Holz Shale Member; late early Campanian (Matsumoto, 1960, p. 102).
- 1054 CIT (=LACMIP 10793): Shale beds above conglomerate, near head of Aliso Creek, 1725'N, 1575'W of SE corner sec. 33, T5S, R7W, Santiago Peak quadrangle, Santa Ana Mountains, Orange County, California. Collected by: W. P. Popenoe, 10/14/34. Ladd Formation, upper part of Holz Shale Member; late early Campanian.
- 1057 CIT (=LACMIP 10794): Shale beds overlying conglomerate lens, about 250' NW of CIT 1054, 1925'N, 1725'W of SE corner sec. 33, T5S, R7W, Santiago Peak quadrangle, Santa Ana Mountains, Orange County, California. Collected by: W. P. Popenoe, 10/26/34. Ladd Formation, upper part of Holz Shale Member; late early Campanian.
- 1060 CIT (=LACMIP 8196): Crest of high NE-SW trending ridge between Santiago and Williams Canyons, 1900'N, 1650'E of SW corner sec. 20, T5S, R7W, El Toro quadrangle, Santa Ana Mountains, Orange County, California. Collected by: W. P. Popenoe, 10/14/35. Ladd Formation, uppermost part of Holz Shale Member; late early Campanian.
- 1158 CIT (=LACMIP 10710): SE slope of Simi Hills, north bank Bell Canyon, 1.15 mi due west of Los Angeles-Ventura County line on boundary (extended) between T1 and 2N, 500'S, 9000'W of NE corner sec. 4, T1N, R17W. Calabasas quadrangle, Ventura County, California. Collected by: W. P. Popenoe, 6/18/35. Chatsworth Formation; middle Campanian.
- 1159 CIT (=LACMIP 10715): Prominent fossil bed on crest of spur between forks of Dayton Canyon, about 400'E of Los Angeles-Ventura County line, 400'N, 2350'W of SE corner sec. 28, T2N, R17W, Calabasas quadrangle, Los Angeles County, California. Collected by: W. P. Popenoe, H. L. Popenoe, and R. Durbin, 6/21/35. Chatsworth Formation; late middle Campanian (Matsumoto, 1960, p. 103; Popenoe, Saul, and Susuki, 1987, p. 99; Saul and Popenoe, 1992, p. 60).
- 1527 UCLA: South of Santiago Creek, along Santiago Truck Trail, SW 1/4, SW 1/4 sec. 28, T5S, R7W, Santiago Peak quadrangle, Santa Ana Mountains, Orange County, California. Collected by: T. Bear, 1940. Ladd Formation, Holz Shale Member; probably late early Campanian.
- 1545 CIT (=LACMIP 8147): Laguna Seca Section, 2500'S, 1300'E of sec. 13, T12S, R10E, on cliff in pebbly bed approximately 50' above contact with silty clay-stone, Los Baños quadrangle, Merced County, California. Collected by: B. Adams. Moreno Formation, Tierra Loma Shale Member; early late Maastrichtian.
- 2858 LACMIP: (Webster locality 25) Top of S slope, elevation 272', N fork of Ammonite Ravine, about 1125 m E of mouth of Ammonite Ravine, E side Arroyo Santa Catarina, about 6.4 km inland from Pacific Ocean, Baja California, Mexico. Collected by: M. Webster, 1966. Rosario Formation; late Campanian-early Maastrichtian.
- 3632 UCLA: West side Chico Creek about 1/3 mi up deep ravine, 2/3 mi S of Mickey's Place, 1750'S, 25'W of NE corner sec. 11, T23N, R2E, Paradise quadrangle, Butte County, California. Collected by: L. R. and R. B. Saul, 1952. Chico Formation, basal part of Tenmile Member, late Santonian or early Campanian.
- 3635 UCLA: On E bank of Chico Creek W from HB House and approximately 400' S of twin meadows, 1800'S, 400'E of NW corner sec. 13, T23N, R2E, Paradise quadrangle, Butte County, California. Collected by: L. R. and R. B. Saul, 8/17/1952. Chico Formation, Tenmile Member, early Campanian.
- 3637 UCLA: East bank Chico Creek, 1250'N of SE corner sec. 14, T23N, R2E, Paradise quadrangle, Butte County, California.

- Collected by: L. R. and R. B. Saul, 8/18/1952. Chico Formation, Tenmile Member, early Campanian.
- 3643 UCLA: W bank Chico Creek, 1500'S, 2500'W of NE corner sec. 26, T23N, R2E, Paradise quadrangle, Butte County, California. Collected by: L. R. and R. B. Saul, 8/22/1952. Chico Formation, Tenmile Member; early Campanian.
- 3648 UCLA = LACMIP 10861, 23648: Fossil Bluff, W side Chico Creek, 1750'S, 1800'E of NW corner sec. 35, T23N, R2E, Paradise quadrangle, Butte County, California. Collected by: L. R. and R. B. Saul, 8/21/1952. Chico Formation, Tenmile Member; early middle Campanian.
- 4082 UCLA: Tuscan Springs, on Little Salt Creek, about 10 mi NE of Red Bluff, near center NE 1/4 sec. 32, T28N, R2W, Tuscan Springs quadrangle, Tehama County, California. Collected by: W. P. Popenoe et al. Chico Formation; early middle Campanian.
- 4192 UCLA: Hills N of Santiago Canyon on crest of long NE trending ridge, 200'N, 2600'E of SW corner sec. 20, T5S, R7W, El Toro quadrangle, Santa Ana Mountains, Orange County, California. Collected by: M. A. Murphy, W. P. Popenoe and T. Susuki, 2/4/59. Ladd Formation, Holz Shale Member, about 105' below top; late early Campanian, *Sub-mortoniceras chicoense* Zone.
- 4224 UCLA: From nodules in mine tunnel on E bank of Butte Creek, about 10' above water's edge, 2.8 mi by road NW of Honey Run Road covered bridge, approximately 2000'S, 250'E of NW corner sec. 17, T22N, R3E, Paradise quadrangle, Butte County, California. Collected by: W. P. Popenoe, Aug. 29, 1952. Chico Formation, Ten Mile Member; middle Campanian.
- 4662 UCLA: N side Mill Creek, approximately 1700'S, 2250'W of NE corner sec. 19, T27N, R2E, Panther Spring quadrangle, Tehama County, California. Collected by: P. U. Rodda, 1954. Chico Formation; late(?) Santonian.
- 4664 UCLA: Just below contact of volcanics and Cretaceous, mouth of Rancheria Creek, 1700'S, 1550'E of NW corner sec. 19, T27N, R2E, Mill Creek Canyon, Panther Spring quadrangle, Tehama County, California. Collected by: P. U. Rodda, 1954. Chico Formation, Kingsley Cave Member (Haggart and Ward, 1984); late Santonian.
- 4878 UCLA: Sucia Island from bluffs on S side of Fossil Bay about 1/4 mile E of W (closed) end of the bay, sec. 26, T38N, R2W, San Juan County, Washington. Collected by: W. P. Popenoe, 8/23/1952. Cedar District Formation; middle Campanian.
- M5906 USGS: On E side of Dip Creek, 2300'S, 1000'W of NE corner sec. 30, T25S, R10E, Lime Mountain quadrangle, San Luis Obispo County, Santa Lucia Mountains, California. Collected by: D. L. Durham and R. J. McLaughlin, 1969; W. O. Addicott, Koichiro Masuda, D. L. Durham, and T. W. Dibblee, Jr., 1970. Asuncion Formation (Taliaferro, 1944); late late Maastrichtian.
- 5990 UCLA: Sandstone cropping out in bed of small northwestward-flowing gully tributary to French Creek, near S end of Swede Basin, 300'S, 1800'E of NW corner sec. 9, T33N, R2W, Millville quadrangle, Shasta County, California. Collected by: W. P. Popenoe, 1/1/1959. Redding Formation, Hooten Gulch Mudstone Member; Coniacian.
- 6044 UCLA: (P 1-72) Soft massive sandstone cropping out on left (east) bank of Butte Creek, just downstream from high bluff at water's edge, and across stream from A-frame house. About 1000'N and 2800'W of SE corner sec. 8, T22N, R3E, Paradise 15 minute (1953) quadrangle, Butte County, California. Collected by: W. P. Popenoe, 6/21/72. Chico Formation, Ten-mile Member; late early Campanian.
- 6950 UCLA: Roadcut N side Silverado Truck Trail, very fossiliferous bed (1'+ thick) immediately below Schultz Conglomerate Member of the Williams Formation, 950'S, 125'W of NE corner sec. 18, T5S, R7W, El Toro quadrangle, Santa Ana Mountains, Orange County, California. Collected by: L. R. Saul, 12/4/81. Ladd Formation, uppermost part of Holz Shale Member; early middle Campanian.
- 6965 LACMIP: Prominent fossil bed on crest of spur between forks of Dayton Canyon, about 400'E of Los Angeles-Ventura County line, 400'N, 2350'W of SE corner Sec. 33, T2N, R17W, Calabasas quadrangle, Simi Hills, Los Angeles County, California. Collected by: J. Alderson, 1974. Chatsworth Formation; middle Campanian, *Hoplitoplacenticeras bowersi* Zone.
- 6996 UCLA: N bank of Bell Canyon about 1 mi W of Los Angeles-Ventura County line on the boundary (extended) between T1N and T2N, R17W, Calabasas quadrangle, Simi Hills, Ventura County, California. Collected by: W. P. Popenoe, 3/27/46. Chatsworth Formation; middle Campanian.
- 7135 UCLA: Fossils collected within 3 m of section in highly fossiliferous zone, W side Horse Canyon near 1800' contour, 1,125 km N, 1.2 km E of SW corner of topo, T9N, R30W, Bates Canyon quadrangle, Sierra Madre Mountains, Santa Barbara County, California. Collected by: Greg Smith, January, 1984. Unnamed formation; late middle or late Campanian.
- 7235 UCLA: Carlsbad Research Park, N side of Faraday Street from cut slope destined to be used for filling Letterbox Canyon, approximately 0.7 mi N, 1.55 mi W of SE corner of San Luis Rey quadrangle, San Diego County, California. Collected by: L. R. Saul, A. R. Loeblich, and J. Loch, 6/21/1984. Point Loma Formation; late Campanian or earliest Maastrichtian.
- 8068 LACMIP: Punta San Jose, Baja California, Mexico. Collected by: unknown. Rosario Formation; late Campanian or early Maastrichtian.
- M8591 USGS Mesozoic: (J 979-199-11) Manzanita Mountain, Santa Barbara County, California. Approximately 11,400'N and 17,025'W of SE corner Manzanita Mountain quadrangle (1964), altitude approximately 1560' in South Fork La Brea Creek near mouth of Lion Canyon, approximately 1.5 miles NE of Manzanita Mountain. Collected by: J. Joyce and J. Vedder, 1979. Unnamed unit; middle or early late Campanian.
- M8611 USGS Mesozoic: Pebby sandstone just S of head 0.7 mi E of Pigeon Point, latitude 37° 10.86'N, longitude 122° 22.79'W, Pigeon Point 1:24000 quadrangle, San Mateo County, California. Collected by: W. P. Elder, 1990. Pigeon Point Formation; middle to early late Campanian.
- M8756 USGS Mesozoic: (91E-16) Sandstone turbidite 100 m NW of small creek, 1 km SE of Pigeon Point, Pigeon Point 1:24000 quadrangle, San Mateo County, California. Collected by: W. P. Elder, 1991. Pigeon Point Formation; middle or early late Campanian.
- M8759 USGS Mesozoic: (PPLH-A) W end of beach 200 m NE of Pigeon Point Lighthouse, Pigeon Point 1:24000 quadrangle, San Mateo County, California. Collected by: W. P. Elder, 1991. Pigeon Point Formation; middle or early late Campanian.
- A9254 UWBM: Sucia Island, San Juan County, Washington. Latitude 48°46', Longitude 122°52'. Collected by: Peter Ward?, 1972. Cedar District Formation; middle(?) Campanian.
- 10934 LACMIP: Fossils collected along 1450' outcrop of sandstone bed on N-facing slope of canyon N of Modjeska Canyon, 150'S, 1250'E of SW corner sec. 20, to 600'N, 2600'E of SW corner sec. 20, T5S, R7W, El Toro quadrangle, Santa Ana Mountains, Orange County, California. Collected by: L. J. Czel, 1957. Ladd Formation, Holz Shale Member, uppermost sandstone bed; early middle Campanian.
- 11950 LACMIP: Approximately 6.4 km inland from Pacific Ocean, on NW wall of Arroyo Santa Catarina near base (approximately 100 m), Baja California, Mexico. Collected by: R. Demetrio, January 1987. Rosario Formation; late Campanian and early Maastrichtian.
- 14313 LACMIP (=CSUN 1447P): About 450 meters north by northwest of lookout on Warm Springs Mountain, elevation 3460', 47 meters above basement complex—San Francisquito Formation contact in Kirby's measured section number 2 (Kirby, 1991, p. 119), Los Angeles County, California. Collected by: M. X. Kirby, 1990. San Francisquito Formation; late Maastrichtian.
- 14314 LACMIP (=CSUN 1447G): 460 meters north by northwest