

THE SNAKE LEMMA AND LONG EXACT HOMOLOGY SEQUENCE

Consider a commutative diagram of R -modules of the form

$$\begin{array}{ccccccc}
 & A & \xrightarrow{f} & B & \xrightarrow{g} & C & \longrightarrow 0 \\
 & \downarrow a & & \downarrow b & & \downarrow c & \\
 0 & \longrightarrow & A' & \xrightarrow{f'} & B' & \xrightarrow{g'} & C'
 \end{array}$$

If the rows are exact, there is an exact sequence

$$\ker(a) \rightarrow \ker(b) \rightarrow \ker(c) \xrightarrow{d} \operatorname{coker} a \rightarrow \operatorname{coker} b \rightarrow \operatorname{coker} c$$

constituting the picture of a snake in the diagram

$$\begin{array}{ccccccc}
 \ker a & \dashrightarrow & \ker b & \dashrightarrow & \ker c & & \\
 \downarrow \iota & & \downarrow & & \downarrow & & \\
 A & \xrightarrow{f} & B & \xrightarrow{g} & C & \longrightarrow & 0 \\
 \downarrow a & & \downarrow b & & \downarrow c & & \\
 0 & \longrightarrow & A' & \xrightarrow{f'} & B' & \xrightarrow{g'} & C' \\
 \downarrow \pi & & \downarrow & & \downarrow & & \\
 & & \operatorname{coker} a & \dashrightarrow & \operatorname{coker} b & \dashrightarrow & \operatorname{coker} c
 \end{array}$$

$\left. \begin{array}{c} \text{---} \\ \text{---} \end{array} \right\} d$

where the ι are the natural inclusion morphisms of the kernel and the π are the natural projection morphisms.

Proof.

Morphisms between the kernel

First we define the morphisms between the kernel. For example we define the map

$$\bar{f} : \ker a \rightarrow \ker b$$

by

$$x \mapsto f(x).$$

This is well defined, since

$$b(f(x)) = f'(a(x)) = f'(0) = 0,$$

meaning that indeed $f(x) \in \ker b$. Of course, the upper left square then commutes and \bar{f} is a homomorphism of R -modules.

Of course, $\bar{g} : \ker b \rightarrow \ker c$ is defined analogously.

Morphisms between the cokernel

We proceed to define the morphisms between the cokernel. Here we, for example, define the map

$$\bar{f}' : \operatorname{coker} a \rightarrow \operatorname{coker} b$$

by

$$[y] \mapsto [f'(y)]$$

for y in A' . Again we check that this gives us a well defined map:

Let y_1, y_2 be in A' with $[y_1] = [y_2]$. This means that $y_1 - y_2 \in \operatorname{im} a$, therefore we have z in A , such that $a(z) = y_1 - y_2$. Then

$$\begin{aligned} \bar{f}'([y_1]) - \bar{f}'([y_2]) &= \bar{f}'([y_1 - y_2]) = [f'(y_1 - y_2)] \\ &= [f'(a(z))] = [b(f(z))] = 0 \end{aligned}$$

thus $\bar{f}'[y_1] = \bar{f}'[y_2]$. Here we have already used the homomorphism property of \bar{f}' that needs a quick proof:

$$\begin{aligned} \bar{f}'([x] + [y]) &= \bar{f}'([x + y]) = [f'(x + y)] \\ &= [f'(x) + f'(y)] = \bar{f}'([x]) + \bar{f}'([y]) \\ \bar{f}'(r[x]) &= \bar{f}'([rx]) = [f'(rx)] = r[f'(x)] = r\bar{f}'([x]) \end{aligned}$$

for $r \in R$. Of course, the lower left square commutes, in fact, the way we defined \bar{f}' was the only chance to have it commute.

Again we use an analogous construction to obtain the morphism

$$\bar{g}' : \operatorname{coker} b \rightarrow \operatorname{coker} c.$$

The morphism $d : \ker c \rightarrow \operatorname{coker} a$

Now we define the morphism $d : \ker c \rightarrow \operatorname{coker} a$. Starting with an element $x \in \ker c$, we chase x through the diagram until we arrive in $\operatorname{coker} a$:

First we obviously want to map x to $\iota(x) \in C$ and then use the surjectivity of $g : B \rightarrow C$ to get an (not necessarily unique) element y in B , such that $g(y) = \iota(x)$. We then take $b(y) \in B'$ and from

$$g'(b(y)) = c(g(y)) = c(\iota(x)) = 0,$$

we use the exactness of the lower row to obtain a unique(!) element z in A' , with $f'(z) = b(y)$. With $\pi(z)$ we finally arrive in $\operatorname{coker} a$.

We now immediately want to define d as

$$d : \ker c \rightarrow \operatorname{coker} a, \quad x \mapsto \pi(z),$$

however, we first need to show that this way, this gives a well-defined map.

For this take y_1 and y_2 in B , with

$$g(y_1) = g(y_2) = \iota(x).$$

We want to show that this gives the same element $\pi(z)$ in $\operatorname{coker} a$. Since $y_2 - y_1 \in \ker g$, we have a $w \in \ker g$ such that $y_2 = y_1 + w$. Using the exactness of the upper row we can rewrite this as

$$y_2 = y_1 + f(u)$$

with $u \in A$.

We then have

$$z_1 = f'^{-1}(b(y_1)) \text{ and } z_2 = f'^{-1}(b(y_2))$$

as the elements of A' the result from chasing y_1 and y_2 to A' and compute

$$\begin{aligned} z_2 &= f'^{-1}(b(y_2)) = f'^{-1}(b(y_1 + f(u))) \\ &= f'^{-1}(b(y_1)) + f'^{-1}(b(f(u))) \\ &= z_1 + f'^{-1}(f'(a(u))) = z_1 + a(u) \end{aligned}$$

(mind the injectivity of f'), giving us

$$\pi(z_2) = \pi(z_1) + \pi(a(u)) = \pi(z_1)$$

as desired.

Again, d was the only choice to make the diagram commute, we now to need to ascertain that d is morphism of R -modules.

Taking $x_1 + x_2 \in \ker c$, we have $\iota(x_1) + \iota(x_2) \in C$ and since d is well defined, we may take $y_1 + y_2 \in B$, where $b(y_i) = \iota(x_i), i = 1, 2$, as the pre-image. Then the pre-image of $b(y_1) + b(y_2)$ clearly is $z_1 + z_2$, where $f'(z_i) = b(y_i), i = 1, 2$ and finally obtain $\pi(z_1) + \pi(z_2)$ for $d(x_1 + x_2)$, thus

$$d(x_1 + x_2) = d(x_1) + d(x_2).$$

For $rx \in \ker c, r \in R$, the argumentation is completely analogous:

□

We now want to use the snake lemma to show the existence of the long exact homology sequence:

Given a short exact sequence of chain complexes

$$0 \longrightarrow A_{\bullet} \longrightarrow B_{\bullet} \longrightarrow C_{\bullet} \longrightarrow 0$$

there is a long exact sequence

$$\dots \longrightarrow H_{n+1}(C) \longrightarrow H_n(A) \longrightarrow H_n(B) \longrightarrow H_n(C) \longrightarrow H_{n-1}(A) \longrightarrow \dots$$