

## YONEDA LEMMA

Given a category  $\mathcal{C}$ , we can form a category  $\mathbf{Set}^{\mathcal{C}}$ , consisting of all functors  $\mathcal{C} \rightarrow \mathbf{Set}$  with natural transformations as morphisms. The composition of morphisms in  $\mathbf{Set}^{\mathcal{C}}$  is then just the vertical composition of natural transformations.

One easily sees that this composition is associative and for all functors  $F : \mathcal{C} \rightarrow \mathbf{Set}$  the natural transformation  $F \rightarrow F$  with components  $\text{id}_C$  is an identity, thus we have indeed a category. We call  $\mathbf{Set}^{\mathcal{C}}$  the *functor category* of functors from  $\mathcal{C}$  to  $\mathbf{Set}$ .

For each object  $A$  in  $\mathcal{C}$  we obtain a special functor  $\mathcal{C} \rightarrow \mathbf{Set}$  called the *Hom-functor* of  $A$ . It sends each object  $X$  in  $\mathcal{C}$  to the Hom-Set  $\text{Mor}_{\mathcal{C}}$ .

$$h^A : \begin{array}{l} \mathcal{C} \longrightarrow \mathbf{Set} \\ X \longmapsto \text{Mor}_{\mathcal{C}}(A, X) \end{array}$$

$$(X \xrightarrow{f} Y) \longmapsto \left( \begin{array}{c} \text{Mor}_{\mathcal{C}}(A, X) \xrightarrow{h^A(f)} \text{Mor}_{\mathcal{C}}(A, Y) \\ A \rightarrow X \longmapsto A \xrightarrow{\quad} Y \\ \quad \searrow \quad \nearrow f \\ \quad \quad X \end{array} \right)$$

In order to secure that the Hom-sets are indeed sets, we demand that  $\mathcal{C}$  is *locally small*.

**Yoneda's lemma** now states that there is a bijection between the set of natural transformations  $h^A \rightarrow F$  and the set  $F(A)$  for any functor  $F : \mathcal{C} \rightarrow \mathbf{Set}$ .

*Proof.*

Given a natural transformation  $h^A \rightarrow F$  with components  $m_C$  we define the map

$$\begin{array}{ccc} f : \{ \text{natural transformations} \} & \longrightarrow & F(A) \\ h^A \rightarrow F & & \\ h^A \rightarrow F & \longmapsto & m_A(\text{id}_A) \end{array}$$

Obviously it is well defined, we now check for surjectivity and injectivity.

For  $c \in F(A)$  we just define a natural transformation by assigning  $\text{id}_A \mapsto c$  via the component  $m_A$ . This not only gives a natural transformation - it is actually completely determined by that assignment:

To see this consider any object  $X$  in  $\mathcal{C}$ . We need to give a map

$$h^A(X) = \text{Mor}(A, X) \rightarrow F(X).$$

If  $\text{Mor}(A, X)$  is empty, there is only one possible choice. Else let  $g \in \text{Mor}(A, X)$ . Then we consider the diagram

$$\begin{array}{ccc} \text{Mor}(A, A) & \xrightarrow{g \circ -} & \text{Mor}(A, X) \\ \downarrow m_A & & \downarrow m_X \\ F(A) & \xrightarrow{F(g)} & F(X) \end{array}$$

in which we don't know the vertical arrows, but we *do* know two things:

$$\text{id}_A \mapsto m_A(\text{id}_A) = c$$

and that the diagram has to constitute a naturality square.

Thus we get, by tracking  $\text{id}_A$ ,

$$\boxed{m_X(g \circ \text{id}_A) = F(g) \circ m_A(\text{id}_A)}.$$

Thus  $m_X(g)$  is completely determined by  $m_A(\text{id}_A)$ .  $\square$

**Lemma 0.1.** *Let  $F : \mathcal{C} \rightarrow \mathcal{D}$  be fully faithful functor. Then, whenever  $F(X) \xrightarrow{F(f)} F(X')$  is an isomorphism in  $\mathcal{D}$ ,  $X \xrightarrow{f} X'$  is an isomorphism in  $\mathcal{C}$ .*

**Corollary 0.2.** *Fully faithful functors are essentially injective.*

*Proof.*

Let  $F(X) \xrightarrow{F(f)} F(X')$  be an isomorphism. Then there exists

$$\varphi \in \text{Mor}_{\mathcal{D}}(F(X'), F(X))$$

such that  $F(f) \circ \varphi = \text{id}_{F(X')}$ ,  $\varphi \circ F(f) = \text{id}_{F(X)}$ .

Since  $F$  is fully faithful there exists a morphism  $g \in \text{Mor}(X, X')$ , such that  $F(g) = \varphi$ . (This is the injectivity of the map  $\text{Mor}(X, X') \rightarrow \text{Mor}(F(X), F(X'))$ .)

Then we have

$$F(g \circ f) = F(g) \circ F(f) = \text{id}_{F(X)} = F(\text{id}_X).$$

Injectivity then gives  $g \circ f = \text{id}_X$  and similarly  $f \circ g = \text{id}_{X'}$ .  $\square$

In summary one can say:

*Fully faithful functors reflect isomorphisms.*