

# Nucleotide Diversity Loss on a Plant Y Chromosome Following Recent Recombination Suppression

Josh Hough<sup>\*,†,1</sup>, Wei Wang<sup>†</sup>, Spencer C.H. Barrett<sup>†</sup> and Stephen I. Wright<sup>†</sup>

<sup>\*</sup>Department of Plant Sciences, University of California, Davis, <sup>†</sup>Department of Ecology and Evolutionary Biology, University of Toronto

**ABSTRACT** X and Y chromosomes differ in effective population size ( $N_e$ ), rates of recombination, and exposure to natural selection, all of which can affect levels of genetic diversity. On Y chromosomes with suppressed recombination, selection is expected to eliminate neutral variation and reduce the  $N_e$  of Y compared to X chromosomes or autosomes. However, non-selective factors including female biased sex ratios and high variance in male reproductive success can also reduce Y-linked  $N_e$ , making it difficult to infer the causes of low Y-diversity. Here, we investigate the factors affecting levels of polymorphism within the genome during sex chromosome evolution in *Rumex hastatulus* (Polygonaceae), a dioecious plant with young sex chromosomes. Strikingly, we find that neutral diversity for genes on the Y is on average ~2.1% of the value for their homologues on the X, corresponding to a chromosome-wide reduction of ~93% compared to the neutral expectation. We demonstrate that the magnitude of this diversity loss is inconsistent with a reduced male  $N_e$  caused by neutral processes including female-biased sex ratios and high variance in male reproductive success. Instead, using forward simulations, we show that the loss of diversity on the Y can be explained by interference among a large number ( $\geq 800$  Kb) of weakly selected mutations. Our results are in agreement with theory on "interference selection", and provide evidence that the effects of purifying selection over a large number of genetically-linked sites can substantially reduce neutral diversity. Given the recent origin of *R. hastatulus* sex chromosomes (~15MYA), our results imply that Y chromosome degeneration in the early stages may be largely driven by interference rather than positive selection for gene silencing followed by neutral genetic drift.

**KEYWORDS** Sex Chromosome Evolution; Nucleotide Diversity; Recombination; Deleterious Mutations

## Introduction

Morphologically distinct sex chromosomes have evolved multiple times independently in both plants and animals (Westergaard 1958; Ohno 1967; Bull 1983; Charlesworth 1991, 2015). Despite clear biological differences between these kingdoms, X and Y chromosomes in both lineages have undergone similar genetic changes. For example, in both groups the loss of recombination between X and Y chromosomes has been associated with an accumulation of deleterious mutations and a gradual loss of genes from the Y chromosome (Hough *et al.* 2014; Bergero *et al.* 2015; Bachtrog 2013), and in some species, the degeneration of the Y has led to the evolution dosage compensation of the X chromosome (Charlesworth 1996; Muyle *et al.*

2012; Mank 2013; Papadopoulos *et al.* 2015). The independent evolution of these phenomena in such taxonomically distant species suggest that general evolutionary mechanisms may be involved, but inferring the causes of molecular evolution and patterns of polymorphism in genomic regions that lack recombination is a longstanding challenge for both theoreticians and experimentalists (Charlesworth 1978; Feldman *et al.* 1980; Barton 1995; Charlesworth 1996; Otto and Feldman 1997; Charlesworth and Charlesworth 2000; McVean and Charlesworth 2000a).

One fundamental difference between the X and Y chromosomes is that there are 1/3 as many Y-linked gene copies as X-linked ones in a diploid population. Thus, genes on the Y chromosome are expected to experience an effective population size ( $N_e$ ) that is 1/4 that of autosomal genes, whereas the  $N_e$  for genes on the X chromosome should be 3/4 that of autosomal genes (assuming an equal number of reproducing females and males). The lowered  $N_e$  of the Y chromosome implies that the equilibrium level of neutral polymorphism

Copyright © 2016 by the Genetics Society of America

doi: 10.1534/genetics.XXX.XXXXXX

Manuscript compiled: Sunday 13<sup>th</sup> November, 2016%

<sup>1</sup>jhough@ucdavis.edu

- proportional to the product of  $N_e$  and the neutral mutation rate,  $\mu$  - should be lower for Y-linked genes than for their X-linked counterparts. In the absence of recombination, genes on the Y chromosome are also expected to be in strong linkage disequilibrium, making them vulnerable to diversity loss due to selection against strongly deleterious mutations (background selection) and selective sweeps of strongly beneficial mutations (genetic hitchhiking). Furthermore, the build-up of linkage disequilibrium between selected mutations on the Y means that selection will act non-independently across the chromosome such that selection at a focal site may "interfere" with selection at sites with which it is linked. Originally developed by Hill Robertson (1966), a large body of work has now shown that such "selective interference" can substantially reduce both the efficacy of selection and the level of neutral variability (Fisher 1930; Muller 1964; Hill and Robertson 1966; McVean and Charlesworth 2000b). These arguments all suggest that non-recombining Y chromosomes should harbor a lower amount of neutral genetic variability than predicted based on the number of Y chromosomes in a population, but the extent to which background selection, genetic hitchhiking, or selective interference have affected chromosome-wide levels of diversity in natural populations, and the relative importance of these processes, is not well-understood.

In addition to reduced diversity arising from selection, in species with female-biased sex ratios or extensive male-male competition, high variance in male reproductive success is also expected reduce the  $N_e$  experienced by genes on the Y chromosome (Caballero 1995; Charlesworth 2001; Laporte and Charlesworth 2002; Pool and Nielsen 2007; Ellegren 2009), suggesting that inferences about the effects positive of purifying selection need to be distinguished from these neutral processes. Because variance in male reproductive success reduces both Y-linked  $N_e$  and autosomal  $N_e$  (Kimura and Crow 1964; Nomura 2002), evidence for this can therefore be obtained by comparing levels of neutral diversity on X and Y chromosomes relative to values on autosomes. For example, high variance in male reproductive success causes a reduction in the Y/A diversity ratio, but an increase in the X/A ratio. Based on such comparisons, studies in humans, for example, have suggested that the inflated X/A ratio is attributable to a historical excess of breeding females over males (Hammer *et al.* 2008) (and see (Bustamante and Ramachandran 2009; Hammer *et al.* 2010; Cotter *et al.* 2016)).

Despite widespread interest in determining the evolutionary factors affecting neutral diversity on sex chromosomes (Ellegren 2011; Bachtrog 2013), we know very little about the influence of either sex ratio variation or linked selection in determining levels of diversity on more recently evolved sex chromosomes. The time scales over which these different effects are likely to be important is therefore not well understood. In humans, estimates of Y-linked diversity are considerably lower than predicted under neutral models, and simulations suggest that levels of diversity are consistent with strong purifying selection (Wilson Sayres *et al.* 2014). However, given that human sex chromosomes evolved from autosomes ~200 million years ago (MYA), is not immediately obvious whether purifying selection might have such strong effects on Y chromosomes that evolved *do novo* from autosomes over much more recent evolutionary time (e.g., within the last 20 MYA in the case of plants (Charlesworth 2015)). Y-linked diversity loss might on the one hand be expected to be lower in younger systems due

to a shorter history of recombination suppression. On the other hand, simulations of strong selection models (background selection and genetic hitchhiking) suggest that these processes may have the greatest effects during the earliest stages of sex chromosome evolution, before the Y has lost many of its genes (Bachtrog 2008). Moreover, work by (Kaiser and Charlesworth 2009) and (Good *et al.* 2014) have provided evidence that even weak purifying selection, if acting non-independently across large number of sites, can generate substantial deviations from neutrality, whereas classic background selection theory breaks down in such cases. Given that a large number of sites are likely to be in linkage disequilibrium on a recently evolved Y chromosome, such "interference selection" *sensu* (Good *et al.* 2014) is a good candidate model for exploring the evolutionary dynamics of young plant sex chromosomes.

To investigate the factors affecting nucleotide diversity in the early stages of sex chromosome evolution, we analyzed neutral polymorphism levels on X, Y, and autosomal chromosomes in the plant *R. hastatulus* (Polygonaceae). This species is a dioecious annual with heteromorphic X and Y chromosomes that evolved approximately 15 MYA (Quesada del Bosque *et al.* 2011; Grabowska-Joachimik *et al.* 2015; Navajas-Pérez *et al.* 2005), making Y chromosomes in this species over 100 million years younger than the highly degenerated Y chromosomes in mammals (Lahn and Page 1999; Ross *et al.* 2005). *Rumex hastatulus* has also received particular attention because of the occurrence of an intraspecific polymorphism in sex chromosome system, in which both XY and XY<sub>1</sub>XY<sub>2</sub> males occur in geographically distinct populations ("chromosomal races") (Smith 1963). The XY<sub>1</sub>XY<sub>2</sub> sex chromosome system in this species is thought to have originated through an X-autosome fusion, with the XY system maintaining the ancestral chromosome complement (Smith 1964). Despite the recent origin of sex chromosomes in both races, there is evidence that both Y's have undergone significant gene loss and functional deterioration (Hough *et al.* 2014). Here, to simplify our comparison of polymorphism levels on X, Y, and autosomes, we focus only on the ancestral Y chromosome, which occurs in both sex chromosome races.

Of particular importance for our study, *R. hastatulus* populations have been found to consistently exhibit female-biased reproductive sex ratios, with a mean sex ratio of  $N_f / (N_m + N_f) = 0.6$  (Pickup and Barrett 2013). Female-biased sex ratios are not uncommon in dioecious plants with heteromorphic X and Y chromosomes (Field *et al.* 2013; Hough *et al.* 2013), and their occurrence in *R. hastatulus* provides an excellent opportunity to study both the demographic and selective factors contributing to sex-linked variability.

## Materials and Methods

### Population Samples and Sex-Linked Genes

We analyzed sex-linked and autosomal genes identified from Illumina RNA sequence data from 12 males and 12 females (1 male and 1 female from each of 6 populations). Samples were collected in 2010 from throughout the native range of *R. hastatulus* (locations in Table S1), and plants were grown in the glasshouse at the University of Toronto from seeds collected from open-pollinated females. We extracted RNA from leaf tissue using Spectrum Plant Total RNA kits (Sigma-Aldrich). Isolation of mRNA and cDNA synthesis was conducted according to standard Illumina RNAseq procedures, and sequencing was conducted on two Illumina HiSeq lanes with 150-bp end reads at the Genome Quebec Innovation Cen-

ter. Reads from these samples were mapped to the *R. hastatus* reference transcriptome (Hough *et al.* 2014) using the BurrowsWheeler Aligner (Li and Durbin 2010), followed by Stampy (Lunter and Goodson 2011). We used Picard tools (<http://picard.sourceforge.net>) to process mapping alignments for the Genome Analysis Toolkit (McKenna *et al.* 2010) variant calling software, and subsequently removed genes with low coverage (<10x) and low Phred Quality Scores <20. The population samples analyzed here were previously reported in Hough *et al.* (2014), where they were used to validate the ascertainment of sex-linked genes identified through segregation analysis, and raw sequences are available from the GenBank Short Read Archive under accession no. SRP041588. Here, to consider sex linked genes that were identified in all of our sequenced population samples, we focused on the previously described set of 460 X/Y genes for which a Y homolog was found in both the Texas and North Carolina races (i.e., X/Y genes where the Y copy was inferred to be on the  $Y_1$  chromosome).

### Autosomal Genes

In evaluating evidence for nucleotide diversity differences between X and Y chromosomes, it is important to distinguish between reduced Y-linked diversity, and the possibility that X-linked diversity is elevated above the level predicted from a neutral model. To do this, we normalized our sex-linked diversity estimates by autosomal diversity, and compared empirical X/A and Y/A nucleotide diversity ratios to those predicted from neutral models and from simulations (described below). Because the criteria for ascertaining autosomal loci in Hough *et al.* (2014) were based on identifying four segregating SNPs per locus, and since this set of genes is likely to be higher in diversity than the average autosomal gene, here we instead used the larger set of all non-sex linked (putatively autosomal) genes as our autosomal reference. We filtered genes in this set to remove any genes that may have been sex-linked but were not identified as such by Hough *et al.*'s conservative ascertainment criteria. In particular, we removed: (i) any genes in which there was evidence for at least one SNP with a sex-linked segregation pattern, (ii) any genes where SNPs showed fixed heterozygosity in males and fixed homozygosity in females, (iii) genes with less than 10X coverage or greater than 100X coverage from independently obtained genomic coverage data (to filter out duplicates or genes with highly repetitive sequences), and (iv) any genes containing SNPs with large (>0.4) allele frequency differences between males and females. Finally, we removed genes with fewer than 100 synonymous sites to avoid biasing our results toward genes that may have been particularly short due to assembly problems. This filtering resulted in a final set of 12,356 autosomal genes.

### Phasing X and Y alleles

To estimate polymorphism for X and Y sequences separately, it is necessary to infer the phase of SNPs in sex-linked transcripts in males. In previous work, phasing alleles on *R. hastatus* sex chromosomes was achieved using segregation analysis from a genetic cross. Here, to phase SNPs from population samples where such segregation data was unavailable, we used HAPCUT (Bansal and Bafna 2008), a maximum-cut based algorithm that reconstructs haplotypes using sequenced fragments (Illumina read data) from the two homologous chromosomes to output a list of phased haplotype blocks containing the SNP variants on each chromosome. Because the resulting

haplotype blocks produced by HAPCUT contained SNPs that were phased relative to each other, but not designated to either the X or Y chromosome, we assigned individual variants to X or Y by independently identifying fixed X-Y differences within each haplotype block (i.e., sites where all females were homozygous, and all males were heterozygous). Identifying such fixed differences within phased haplotype blocks enabled us to then infer the correct phase (X or Y) of the polymorphisms from HAPCUTs output. In particular, this was done by matching the phase of fixed X-Y differences with their neighboring polymorphic sites: when a fixed X-Y difference occurred in the same phased haplotype block as a polymorphic site, then the polymorphic variants in that block were assigned to either X or Y based on the known phase of the fixed difference with which they were matched. SNPs that were identified outside of phased blocks, or in blocks without fixed X-Y differences, were recorded as missing data. Finally, we filtered out SNPs with coverage > 60, QUAL score > 60, and those within a distance of 10bp or less from indels. This filtering procedure resulted in fasta-formatted alignments of X and Y sequences for 372 sex-linked genes.

We further validated the results of HAPCUTs allele phasing by comparing the accuracy of this method with the phasing-by-segregation method that was conducted in Hough *et al.* (2014). To do this, we first phased the sequence data from parents and their progeny using HAPCUTs algorithm (using the same parameters as for the population data), and then identified cases where SNPs were inferred on the Y chromosome by HAPCUT, but where the true level polymorphism, obtained from the genetic cross, was zero. We identified 7 % of sex-linked genes that either had phasing errors of this kind genotyping errors. This corresponds to a SNP error rate estimate of  $1.7 \times 10^{-4}$ . Note that this rate is very low relative to population-based estimates of polymorphism on the X and autosomes (Table 1), and therefore should have minimal effects on our estimation of the X/A ratio. However, because this rate is high relative to the expected level of polymorphism on the Y chromosome, we further filtered genes in which we found evidence for false-positive SNP calls arising from: (i) phasing errors caused by gene duplicates (more than two haplotypes), (ii) polymorphisms around indels, and (iii) genotyping errors caused by low Y-expression. This final filtering was conducted by manually checking each individual putative polymorphism on the Y chromosome using IGV (Robinson *et al.* 2011).

### Estimating nucleotide diversity on sex chromosomes and autosomes

For each locus in our analysis, we calculated Wattersons (1975) estimator of the population parameter  $\theta = 4N_e\mu$ , where  $N_e$  is the effective population size, and  $\mu$  is the mutation rate (Watterson 1975), using a modified version of the Perl program Polymorphurama (Bachtrog and Andolfatto 2006). To compare sex-linked and autosomal loci, we calculated the average value of  $\theta$ , weighted by the number of synonymous sites in each gene (Figure 2). We obtained 95% confidence intervals for X/A and Y/A ratios by bootstrapping per gene using the BCa method (Efron and Tibshirani 1994) implemented in the Boot package in R (Canty and Ripley 2012), and calculating X/A and Y/A on each iteration for 20000 replicates each. Bootstrapping was conducted on the final filtered set of 173 sex-linked, and 12355 autosomal genes. Note that the lack of recombination on the Y chromosome implies that statistical assumptions about inde-

pendence across loci are violated, suggesting that the true uncertainty in the estimate  $Y/A$  may be wider than implied by bootstrapping results. To address this, we also used a maximum likelihood approach, implemented in a modified version of the MLHKA software (Wright and Charlesworth 2004), to independently estimate a credibility interval for the  $Y/A$  ratio (Figure S1). Because of the thousands of genes involved, a likelihood method incorporating divergence to control for heterogeneity in mutation rate was not feasible, as this would require maximizing the likelihood estimate of the mutation rate for each locus independently. Therefore, we assumed no heterogeneity in mutation rate, no recombination between Y-linked genes, and free recombination between autosomal loci. Our model thus had two parameters:  $\theta_{autosomal}$  and  $Y/A$ . We varied both parameters and evaluated the likelihood for  $Y/A$  from 0.001 to 1, and  $\theta_{autosomal}$  from 0.001 to 0.01.

### Neutral predictions and the effect of sex ratio bias on diversity

To test whether our estimated levels of diversity on X, Y and autosomal chromosomes could be explained by neutral processes, we compared our estimated levels diversity to predictions from a model of the expected  $N_e$  of males and females when the variance in offspring number is either smaller or larger than Poisson (Kimura and Crow 1964; Hedrick 2011). As we were primarily interested in the possible influence of variance in male reproductive success on Y-linked  $N_e$ , and the parameter space within which such a model might generate  $Y/A$  and  $X/A$  ratios consistent with our data, we ignored non-poisson variance in female offspring, and calculated the male  $N_e$  as:

$$N_{em} = \frac{N_m k_m - 1}{k_m - 1 + \frac{V_{k_m}}{\bar{k}}} \quad (1)$$

where  $k_m$  and  $V_{k_m}$  are the mean and variance in male offspring number, respectively, in a population of  $N_m$  reproducing males, and  $\bar{k}$  is the total mean number of progeny (Kimura and Crow 1964). Given expected values of  $N_{em}$  for a range of values for  $V_{k_m}$  and  $N_m$ , the corresponding expectation for the  $N_e$  of autosomes and sex chromosomes at equilibrium were calculated as:

$$N_{eA} = \frac{4N_m N_f}{N_m + N_f} \quad (2)$$


$$N_{eX} = \frac{9N_m N_f}{4N_m + 2N_f} \quad (3)$$

$$N_{eY} = \frac{N_m}{2} \quad (4)$$

following (Wright 1931). Note that with equal sex ratios  $N_f = N_m$ , and Poisson-distributed offspring numbers ( $V_k = \bar{k}$ ), then  $N_{eX}/N_{eA} = 0.75$ , and  $N_{eY}/N_{eA} = 0.25$ . Figure 1 shows these predictions as a function of the sex ratio for a range values of  $V_{k_m}$ , and we tested the fit of our empirical diversity ratios to these predictions (shown in Figure 2) assuming the estimated *R. hastatulus* sex ratio of 0.6 (Pickup and Barrett 2013).

### Simulations of purifying selection

To study the effects of purifying selection on expected levels of Y-chromosome diversity, we conducted forward-time simulations of haploid Y chromosomes using the software SFSCODE (Hernandez 2008). We first estimated the distribution of fitness effects of deleterious mutations from our polymorphism



**Figure 1** The relation between effective population size and sex ratio bias for genes on autosomes (A) and the X chromosome (B), and corresponding normalized X/A and Y/A ratios (C). The sex ratio is shown as the proportion of females,  $N_f/(N_f + N_m)$ , plotted against  $N_e/N$ , where  $N = N_m + N_f$ . Dotted curves show equilibrium predictions when both males and females produce Poisson-distributed offspring numbers and ( $V_k = \bar{k} = 2$ ). Solid curves correspond to increasing values for  $V_{k_m}$ , the variance in male offspring number, ranging from Poisson/2 (top solid curve in panels A and B) to  $3 \times \text{Poisson}$  (bottom most curve). For (C) we assumed equal neutral mutation rates among genes and calculated the expected neutral diversity as  $\theta = 4N_e\mu$ .

data for X-linked genes using the method of (Keightley and Eyre-Walker 2007), which fits a gamma distribution of selection coefficients to the observed frequency distribution of nonsynonymous and synonymous polymorphism. We used this estimated gamma distribution to parameterize the simulations, initializing each with our estimated  $\theta$  from autosomal genes, but appropriately adjusted to reflect the expected reduction in Y-linked  $N_e$  for a sex ratio of  $N_f/(N_f + N_m) = 0.6$ . To match our sample size and the number of synonymous sites sample from our data (see Supporting Information), the simulations sampled 6 haploid chromosomes, and simulated sequences contained 45,331bp of linked neutral sequence from which we calculated diversity. To examine the expected reduction in diversity ( $\pi/\pi_0$ ) as a function of the number of selected sites ( $L$ ), and to estimate this parameter from our data, we ran simulations over a range of values of  $L$  (up to a maximum of  $5 \times 10^6$ ) (Figure 3A). For each such simulation, we calculated the approximate likelihood of our observed data based on the proportion of simulations in which synonymous diversity,  $\pi_{simulated}$ , was not significantly different from our empirical estimate,  $\pi_{observed}$  (Figure 3B).

## Results and Discussion

### Extensive loss of Y-chromosome diversity

#### Female biased sex ratio and variance in male offspring number

#### Purifying selection


#### Conclusions

### Acknowledgments

### Literature Cited

Bachtrog, D., 2008 The temporal dynamics of processes underlying y chromosome degeneration. *Genetics* **179**: 1513–1525.





**Figure 2** new figure coming soon).

- Bachtrog, D., 2013 Y-chromosome evolution: emerging insights into processes of y-chromosome degeneration. *Nature Reviews Genetics* **14**: 113–124.
- Bachtrog, D. and P. Andolfatto, 2006 Selection, recombination and demographic history in *Drosophila miranda*. *Genetics* **174**: 2045–2059.
- Bansal, V. and V. Bafna, 2008 Hapcut: an efficient and accurate algorithm for the haplotype assembly problem. *Bioinformatics* **24**: i153–i159.
- Barton, N., 1995 A general model for the evolution of recombination. *Genetical research* **65**: 123–144.
- Bergero, R., S. Qiu, and D. Charlesworth, 2015 Gene loss from a plant sex chromosome system. *Current Biology* **25**: 1234–1240.
- Bull, J., 1983 *Evolution of sex determining mechanisms..* The Benjamin/Cummings Publishing Company, Inc.
- Bustamante, C. D. and S. Ramachandran, 2009 Evaluating signatures of sex-specific processes in the human genome. *Nature genetics* **41**: 8–10.
- Caballero, A., 1995 On the effective size of populations with separate sexes, with particular reference to sex-linked genes. *Genetics* **139**: 1007–1011.
- Canty, A. and B. Ripley, 2012 boot: Bootstrap r (s-plus) functions. R package version 1.
- Charlesworth, B., 1978 Model for evolution of y chromosomes and dosage compensation. *Proceedings of the National Academy of Sciences* **75**: 5618–5622.
- Charlesworth, B., 1991 The evolution of sex chromosomes. *Science* **251**: 1030–1033.
- Charlesworth, B., 1996 The evolution of chromosomal sex determination and dosage compensation. *Current Biology* **6**: 149–162.
- Charlesworth, B., 2001 The effect of life-history and mode of inheritance on neutral genetic variability. *Genetical research* **77**: 153–166.
- Charlesworth, B. and D. Charlesworth, 2000 The degeneration of y chromosomes. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* **355**: 1563–1572.
- Charlesworth, D., 2015 Plant contributions to our understanding of sex chromosome evolution. *New Phytologist* **208**: 52–65.
- Cotter, D. J., S. M. Brotman, and M. A. W. Sayres, 2016 Genetic diversity on the human x chromosome does not support a strict pseudoautosomal boundary. *Genetics* **203**: 485–492.
- Efron, B. and R. J. Tibshirani, 1994 *An introduction to the bootstrap*. CRC press.
- Ellegren, H., 2009 The different levels of genetic diversity in sex chromosomes and autosomes. *Trends in Genetics* **25**: 278–284.
- Ellegren, H., 2011 Sex-chromosome evolution: recent progress and the influence of male and female heterogamety. *Nature Reviews Genetics* **12**: 157–166.
- Feldman, M. W., F. B. Christiansen, and L. D. Brooks, 1980 Evolution of recombination in a constant environment. *Proceedings of the National Academy of Sciences* **77**: 4838–4841.
- Field, D. L., M. Pickup, and S. C. Barrett, 2013 Comparative analyses of sex-ratio variation in dioecious flowering plants. *Evolution* **67**: 661–672.
- Fisher, R. A., 1930 *The genetical theory of natural selection: a complete variorum edition*. Oxford University Press.
- Good, B. H., A. M. Walczak, R. A. Neher, and M. M. Desai, 2014 Genetic diversity in the interference selection limit. *PLoS Genet* **10**: e1004222.
- Grabowska-Joachimik, A., A. Kula, T. Książczyk, J. Chojnicka, E. Sliwiska, and A. J. Joachimik, 2015 Chromosome landmarks and autosome-sex chromosome translocations in *Rumex hastatulus*, a plant with xx/xy1y2 sex chromosome system. *Chromosome Research* **23**: 187–197.
- Hammer, M. F., F. L. Mendez, M. P. Cox, A. E. Woerner, and J. D. Wall, 2008 Sex-biased evolutionary forces shape genomic patterns of human diversity. *PLoS Genet* **4**: e1000202.
- Hammer, M. F., A. E. Woerner, F. L. Mendez, J. C. Watkins, M. P. Cox, and J. D. Wall, 2010 The ratio of human x chromosome to autosome diversity is positively correlated with genetic distance from genes. *Nature genetics* **42**: 830–831.
- Hedrick, P. W., 2011 *Genetics of populations*. Jones & Bartlett Learning.
- Hernandez, R. D., 2008 A flexible forward simulator for populations subject to selection and demography. *Bioinformatics* **24**: 2786–2787.
- Hill, W. G. and A. Robertson, 1966 The effect of linkage on limits to artificial selection. *Genetical research* **8**: 269–294.
- Hough, J., J. D. Hollister, W. Wang, S. C. Barrett, and S. I. Wright, 2014 Genetic degeneration of old and young y chromosomes in the flowering plant *Rumex hastatulus*. *Proceedings of the National Academy of Sciences* **111**: 7713–7718.
- Hough, J., S. Immler, S. C. Barrett, and S. P. Otto, 2013 Evolutionarily stable sex ratios and mutation load. *Evolution* **67**: 1915–1925.
- Kaiser, V. B. and B. Charlesworth, 2009 The effects of deleterious mutations on evolution in non-recombining genomes. *Trends in Genetics* **25**: 9–12.
- Keightley, P. D. and A. Eyre-Walker, 2007 Joint inference of the distribution of fitness effects of deleterious mutations and population demography based on nucleotide polymorphism frequencies. *Genetics* **177**: 2251–2261.
- Kimura, M. and J. F. Crow, 1964 The number of alleles that can be maintained in a finite population. *Genetics* **49**: 725–738.
- Lahn, B. T. and D. C. Page, 1999 Four evolutionary strata on the human x chromosome. *Science* **286**: 964–967.
- Laporte, V. and B. Charlesworth, 2002 Effective population size and population subdivision in demographically structured populations. *Genetics* **162**: 501–519.
- Li, H. and R. Durbin, 2010 Fast and accurate long-read alignment with burrows-wheeler transform. *Bioinformatics* **26**:

- 589–595.
- Lunter, G. and M. Goodson, 2011 Stampy: a statistical algorithm for sensitive and fast mapping of illumina sequence reads. *Genome research* **21**: 936–939.
- Mank, J. E., 2013 Sex chromosome dosage compensation: definitely not for everyone. *Trends in genetics* **29**: 677–683.
- McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis, A. Kernytsky, K. Garimella, D. Altshuler, S. Gabriel, M. Daly, *et al.*, 2010 The genome analysis toolkit: a mapreduce framework for analyzing next-generation dna sequencing data. *Genome research* **20**: 1297–1303.
- McVean, G. A. and B. Charlesworth, 2000a The effects of hill-robertson interference between weakly selected mutations on patterns of molecular evolution and variation. *Genetics* **155**: 929–944.
- McVean, G. A. and B. Charlesworth, 2000b The effects of hill-robertson interference between weakly selected mutations on patterns of molecular evolution and variation. *Genetics* **155**: 929–944.
- Muller, H. J., 1964 The relation of recombination to mutational advance. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* **1**: 2–9.
- Muyle, A., N. Zemp, C. Deschamps, S. Mousset, A. Widmer, and G. A. Marais, 2012 Rapid de novo evolution of x chromosome dosage compensation in *silene latifolia*, a plant with young sex chromosomes. *PLoS Biol* **10**: e1001308.
- Navajas-Pérez, R., R. de la Herrán, G. L. González, M. Jamilena, R. Lozano, C. R. Rejón, M. R. Rejón, and M. A. Garrido-Ramos, 2005 The evolution of reproductive systems and sex-determining mechanisms within *rumex* (polygonaceae) inferred from nuclear and chloroplastidial sequence data. *Molecular Biology and Evolution* **22**: 1929–1939.
- Nomura, T., 2002 Effective size of populations with unequal sex ratio and variation in mating success. *Journal of Animal Breeding and Genetics* **119**: 297–310.
- Ohno, S., 1967 *Sex chromosomes and sex-linked genes*, volume 1. Springer-Verlag.
- Otto, S. P. and M. W. Feldman, 1997 Deleterious mutations, variable epistatic interactions, and the evolution of recombination. *Theoretical population biology* **51**: 134–147.
- Papadopoulos, A. S., M. Chester, K. Ridout, and D. A. Filatov, 2015 Rapid y degeneration and dosage compensation in plant sex chromosomes. *Proceedings of the National Academy of Sciences* **112**: 13021–13026.
- Pickup, M. and S. C. Barrett, 2013 The influence of demography and local mating environment on sex ratios in a wind-pollinated dioecious plant. *Ecology and evolution* **3**: 629–639.
- Pool, J. E. and R. Nielsen, 2007 Population size changes reshape genomic patterns of diversity. *Evolution* **61**: 3001–3006.
- Quesada del Bosque, M., R. Navajas-Pérez, J. Panero, A. Fernández-González, and M. Garrido-Ramos, 2011 A satellite dna evolutionary analysis in the north american endemic dioecious plant *rumex hastatulus* (polygonaceae). *Genome* **54**: 253–260.
- Robinson, J. T., H. Thorvaldsdóttir, W. Winckler, M. Guttman, E. S. Lander, G. Getz, and J. P. Mesirov, 2011 Integrative genomics viewer. *Nature biotechnology* **29**: 24–26.
- Ross, M. T., D. V. Grafham, A. J. Coffey, S. Scherer, K. McLay, D. Muzny, M. Platzer, G. R. Howell, C. Burrows, C. P. Bird, *et al.*, 2005 The dna sequence of the human x chromosome. *Nature* **434**: 325–337.
- Smith, B. W., 1963 The mechanism of sex determination in *rumex hastatulus*. *Genetics* **48**: 1265.
- Smith, B. W., 1964 The evolving karyotype of *rumex hastatulus*. *Evolution* pp. 93–104.
- Watterson, G., 1975 On the number of segregating sites in genetical models without recombination. *Theoretical population biology* **7**: 256–276.
- Westergaard, M., 1958 The mechanism of sex determination in dioecious flowering plants. *Advances in genetics* **9**: 217–281.
- Wilson Sayres, M. A., K. E. Lohmueller, and R. Nielsen, 2014 Natural selection reduced diversity on human y chromosomes. *PLoS Genet* **10**: e1004064.
- Wright, S., 1931 Evolution in mendelian populations. *Genetics* **16**: 97–159.
- Wright, S. I. and B. Charlesworth, 2004 The hka test revisited. *Genetics* **168**: 1071–1076.