

Deep Inelastic Scattering Revisited

Felix Hekhorn
University of Jyväskylä

Lecture notes for the Jyväskylä Summer School 2026

Abstract

Discuss Yadism [\[1\]](#).

Contents

1	The basics	1
1.1	Introduction	1
1.1.1	Motivation and non-goals	1
1.1.2	Requirements	3
1.1.3	The very basics	4
1.2	Setting the stage	6
1.3	Distributions	9
1.4	Next-to-leading order	9
2	DIS in real life	11
2.1	DIS is diverse	11
2.1.1	HERA	11
2.1.2	Neutrino DIS	13
2.2	Interpolation grids	13
2.3	DGLAP	13
2.4	Scale variations	13
2.5	Heavy Quarks	13
2.6	γ_5	14
2.7	TMC	14
3	More advanced topics	15
3.1	Positivity	15
3.2	Higher orders	15
3.3	QED	15
3.4	Valo	15
	Bibliography	17

Chapter 1

The basics

1.1 Introduction

1.1.1 Motivation and non-goals

We aim in this course to make a meaningful comparison between the theoretical predictions and the experimental measurement of a fully inclusive Deeply Inelastic Scattering (DIS) cross section. Instead of an abstract, theoretical discussion about DIS, which can be found in a standard lecture on particle physics, we're looking here to make a connection with real-life physics solving current research questions. The main focus is be naturally on PDF, for which DIS plays a major role. In fact, DIS was, is, and will be a very important constrain in determining Parton Distribution Functions (PDFs) - only recently, e.g. starting from NNPDF4.0 [2], LHC data starts to dominate in PDF fits. Stressing the importance and relevance of PDFs for any kind of perturbative QCD (pQCD) application and to understand the inner structure of hadrons is left to the literature [3–5]. In turn, DIS itself can be used to answer unresolved questions about PDFs, e.g. whether the proton has an intrinsic charm component [6].

In order to obtain a more realistic DIS treatment, we review general pQCD features and exemplify them using DIS. In fact, fully inclusive DIS comes in very handy, since here the effects mostly appear in their simplest form and when considering hadronic collisions, e.g. at the LHC, the strategies are mostly the same just more involved. We keep a phenomenological point of view with a specific focus on the numerical implementation as, eventually, we want to compare to the experimental measurements. We refrain from giving detailed proofs in this course and point to the standard textbooks [7–11] for that.

Instead, we consider different features as given and focus on their practical implications. However, a major complication in this course is the interconnection between almost all features and while we discuss them here in turn, we highlight also how they are intertwined with each other.

In recent years (and much more so hopefully in the coming years), DIS has become a major research topic again thanks to the advent of the Electron Ion Collider (EIC) [12]. While the planned physics program of the EIC goes far beyond standard fully inclusive DIS, this simplest case can serve as an introduction into more complicated topics.

The structure of this course follows in large parts the implementation of Yadism (Yet Another DIS module) [1, 13], which is developed and used by the NNPDF collaboration in their PDF determinations. The online documentation, which is available at

<https://yadism.readthedocs.io/> ,

may serve as additional, more technical introduction into the topic. Note that Yadism is an open-source project to which everybody can contribute¹.

Since DIS is a very wide field, it is worth spelling out explicitly some non-goals:

- We only consider pQCD in the collinear framework here. However, because DIS is such a powerful experiment, which offers a clean access to the inner structure of the protons it can be studied using other frameworks as well, such as Color Glass Condensates [14, 15] → ask H. Mäntysaari or T. Lappi
- We only consider fully inclusive cross sections here (to be defined in Section 1.2 and with a tiny exception to be discussed in Section 2.5). In particular, we do not consider any more complicated final states, such as, e.g., single jet or di-jet production [16].
- We only consider pQCD in the collinear framework here. While, Monte Carlo Event generators, such as, e.g., Pythia [17], are rooted in collinear pQCD they eventually go much beyond and we do not consider their details here [18] → ask I. Helenius.
- We only consider DIS here. In particular, we do not consider photo-production or elastic or diffractive cross sections. We define the necessary kinematics for the distinction below in Section 1.1.3.

¹e.g. if some documentation can be improved




Figure 1.1: The Standard Model of Elementary Particles - taken from [19]

- We compute the theoretical predictions here. So while we try to keep the connection to the experimental side close, we do not discuss any specific details about how to measure cross sections.

1.1.2 Requirements

We need some basic knowledge from particle physics and, in particular, you need to be familiar with the Standard Model overview shown in Fig. 1.1. We discuss in this course each and everyone of these particles, with the exception of the three heaviest in each category: the heaviest quark, the top, the heaviest lepton, the tau, and the heaviest boson, the Higgs. They work, in principle, mostly like their lighter brothers and sisters, but in practice are basically always neglected. We give a more thorough justification to discard them when we get to the actual numbers. All other particles explicitly appear in this course and you need to know their charges - see also Fig. 1.1.

Conclusion 1.1: Summary forces, particles, and charges

- The carrier of the strong force is the gluon
- Quarks have a strong charge, leptons don't
- The carriers of the electro-weak force are the photon, the Z boson, and the W boson
- Quarks and charged leptons have electro-magnetic charges
- Quarks and leptons have weak charges

We also need some basic knowledge from Quantum Field Theory (QFT).

1.1.3 The very basics

Let's start from the top:

Conclusion 1.2: DIS in words

fully inclusive DIS = fully inclusive deeply inelastic scattering

What does that actually mean? We begin from the back:

- “scattering” means, of course, we are scattering two particles with each other. In this particular case, we are scattering a lepton off a hadron and to make things simple in the beginning we consider an electron is scattered off a proton.
- “inelastic” means we are breaking the target, the proton, apart; it shatters into many pieces.
- “deeply” is an adjective² to inelastic and it means we are looking “deep” into the inside of the proton. And in order to resolve small things, as usual, we need an highly energetic probe.
- “fully inclusive” means we are observing a single particle in the final state, which is the electron, but nothing else.

Thus, we can write the same information again with a formula

²thus it is “deeply inelastic” or “deep-inelastic”, but not a list of adjectives “deep inelastic”

Conclusion 1.3: DIS in a formula

fully inclusive DIS = $e^-(k) + p(P) \rightarrow e^-(k') + X$ with $-(k - k')^2 \gg \Lambda_{\text{QCD}}^2$

where we have assigned the four-momenta k, P, k' to the incoming electron, the incoming proton and the outgoing electron respectively. As usual X marks the inclusiveness, i.e. we don't observe anything else apart from the electron. By requiring a large momentum-transfer between the incoming and the outgoing electron, i.e. their difference four-momentum vector has a large magnitude, we probe the proton with highly energetic parton (which we can think of as photon).

Next, we need a way to actually compute theory predictions for the cross sections, which you can measure in an experiment. We use the factorization theorem [20],

Conclusion 1.4: Factorization Theorem

$$\sigma(x, Q^2) = \sum_j \int_x^1 \frac{dz}{z} C_j(z, Q^2) f_j(x/z, Q^2) + \mathcal{O}\left(\frac{\Lambda_{\text{QCD}}^2}{Q^2}\right) \quad (1.1)$$

$$= \sum_j (C_j(Q^2) \otimes f_j(Q^2))(x) + \mathcal{O}\left(\frac{\Lambda_{\text{QCD}}^2}{Q^2}\right) \quad (1.2)$$

$$= (\mathbf{C}(Q^2) \otimes \mathbf{f}(Q^2))(x) + \mathcal{O}\left(\frac{\Lambda_{\text{QCD}}^2}{Q^2}\right), \quad (1.3)$$

which states how to compute a cross section $\sigma(x, Q^2)$ from coefficient function $\mathbf{C}(z, Q^2)$, computable in pQCD, and a universal PDF $\mathbf{f}(\xi, Q^2)$. In the remaining part of the course we discuss, enhance and correct this formula in many different ways and basically challenge every aspect. We define each and every variable and function in Section 1.2, so we can focus for the moment on two other important aspects: universality and notation.

The crucial, physical aspect about Eq. (1.1) is universality: for any cross section σ measured at any x and Q^2 there is exactly one PDF $\mathbf{f}(\xi, Q^2)$. Instead, the coefficient functions $\mathbf{C}(z, Q^2)$ may depend on the definition of the cross section, but they are computable from first principles. This implies the predictive powers of our framework: if $\mathbf{C}$ is fixed and $\mathbf{f}$ is universal, we can measure σ in one set of experiments and then predict the outcome of another set of measurements σ' .

The other, more practical aspect of Eqs. (1.1) to (1.3) is the short-hand notation we introduce there. While Eq. (1.1) is the most canonical form of the factorization theorem, which can be found like this in most textbooks, we introduce in Eq. (1.2) the multiplicative convolution symbol $\otimes$. Note that we have suppressed the variable over which the convolution is performed and recall that convolution is associative, i.e. $(g \otimes h)(x) = (h \otimes g)(x)$. Finally, in Eq. (1.3) we introduce the vector notation, where we denote object, which have a non-trivial dependence on the parton flavor j , with bold font. Equation (1.3) looks nice and compact, but it worth recalling that it already hides a lot of details: specifically, $C_j(z, Q^2)$ is an object which depends on three different variables. We use vector notation and the index notation interchangeably depending on the context.

To Leading Order (LO) accuracy, the coefficient functions $\mathbf{C}$ greatly simplify and we find

$$\sigma(x, Q^2) = \sum_j e_{q_j}^2 f_j(x, Q^2), \quad (1.4)$$

i.e. the cross section is given by the sum of the quark PDFs weighted by their electrical charges squared. Note that here the variables, which are used for the experimentally measured cross section on the left-hand-side, are also used as arguments of the PDFs on the right-hand-side. This is a pure coincidence only valid in the simplest case at LO and two historical comments can be made here. First, when studying older literature this equivalence can lead to confusion as, e.g., PDFs have been also called “structure functions” at the time, or what exactly we mean when talking about a general variable “ x ”. Second, in the so-called “parton model”, which was popular before QCD was invented, the index j just referred to a “parton” with some properties. Instead, within QCD we now know exactly, what the object in question is: a quark, which has a strict definition in QFT with well-defined properties and quantum numbers. For simplicity, we still call an object, which can be either a quark or a gluon “parton”.

1.2 Setting the stage

In the following section we collect and define all ingredients needed for Eq. (1.1). As already said, DIS is given by

$$e^-(k) + p(P) \rightarrow e^-(k') + X \quad (1.5)$$

FH: copy DIS diagram from PDG where the four-momenta k, P, k' belong to the incoming electron, the incoming proton and the outgoing electron respectively. This allows us, first, to define four momentum of the exchanged photon

FH!

$$q = k - k' \quad (1.6)$$

and from there the set of four common variables:

$$s = (k + P)^2 = \frac{Q^2}{xy} \quad (1.7)$$

$$Q^2 = -q^2 \quad (1.8)$$

$$x = \frac{Q^2}{2q \cdot P} \quad (1.9)$$

$$y = \frac{q \cdot P}{k \cdot P} \quad (1.10)$$

Note that Eq. (1.7) gives an explicit relation among these, thus we have only three independent and one dependent variable. In practice, the invariant mass of the electron-proton system s is fixed by the experimental setup, i.e. by the energy of the colliding particles. Also the measurable range of y is in practice limited by the experimental geometry. DIS cross sections are typically provided as double-differential cross sections in x and y . Finally, it is convenient to define the mass squared of the system X recoiling against the scatter electron,

$$W^2 = (P + q)^2 = Q^2 \left(\frac{1}{x} - 1 \right) \quad (1.11)$$

Conclusion 1.5: DIS requirements

We require $Q^2 \gg \Lambda_{\text{QCD}}^2$ (deep) and $W^2 \gg \Lambda_{\text{QCD}}^2$ (inelastic) to call a lepton-hadron scattering DIS.

We can decompose the (double-differential) cross section into a leptonic interaction and a hadronic interaction, yielding each a tensor in Minkowski space, $L_{\mu\nu}$ and $W_{\mu\nu}$ respectively. Thus, we can write

$$\frac{d^2\sigma}{dx dy} = \frac{2\pi y \alpha_{\text{em}}^2}{Q^4} L_{\mu\nu} W^{\mu\nu} \quad (1.12)$$

where we have introduced the electro-magnetic coupling α_{em} , which appears once on the leptonic side and once on the hadronic side. We can express the leptonic tensor using plain Quantum Electrodynamics (QED) and provide a decomposition of the hadronic tensor in terms of scalar functions. Together, we obtain

$$\frac{d^2\sigma}{dxdy} = \frac{4\pi\alpha_{\text{em}}^2}{xyQ^2} \left(\left(1 - y + \frac{y^2x}{2}\right) F_2(x, Q^2) - \frac{y^2x}{2} F_L(x, Q^2) \right) \quad (1.13)$$

where we have introduced the two common scalar functions, now called structure functions F_2 and F_L , which are often measured in experiments. It is convenient to also define a third, linearly dependent structure function via

$$2xF_1(x, Q^2) = F_2(x, Q^2) - F_L(x, Q^2). \quad (1.14)$$

Keep in mind that the discussion so far is a major simplification of DIS and in Section 2.1 we reintroduce the main complexity.

Let us continue to collect the remaining ingredient for Eq. (1.1):

- The sum over partons j includes quarks and gluons and we refine in Section 2.5 how large the sum actually is.
- The parton momentum fraction $\xi = x/z$ is the fraction of the four momentum the parton carries with respect to its parent proton.
- The probability of such a configuration to occur is given by the PDF, i.e. $f_j(\xi, Q^2)d\xi$ is the probability to find a parton j with momentum ξP inside a proton with momentum P at an experiment performed at the scale Q^2 .
- The integration over z sums all configuration to generate the final state configuration.

Finally, the coefficient functions $\mathbf{C}(x, Q^2)$ are computable order-by-order in pQCD and thus can be decomposed as

$$\mathbf{C}(x, Q^2) = \sum_{k=0} \alpha_s^k(Q^2) \mathbf{C}^{(k)}(x). \quad (1.15)$$

Note that the Q^2 dependency of $\mathbf{C}(x, Q^2)$ is fully captured by the strong coupling $\alpha_s(Q^2)$.

1.3 Distributions

As we have seen the coefficient functions may contain distributions. For example the quark coefficient function at LO is just a Dirac delta distribution:

$$C_q^{(0)}(z) = \delta(1 - z) . \quad (1.16)$$

To better compare with our next step, let's repeat the definition of the Dirac delta distribution:

$$\int_0^1 dz \delta(1 - z) f(z) = f(1) . \quad (1.17)$$

However, this is only the simplest case, which appears at LO, and at higher order we get more complicated distributions. For example the quark coefficient function at NLO is given by

$$C_q^{(1)}(z) = C_F \left[4 \left(\frac{\ln(1 - z)}{1 - z} \right)_+ - 3 \left(\frac{1}{1 - z} \right)_+ - (9 + 4\zeta_2) \delta(1 - z) - 2(1 + z) \ln \left(\frac{1 - z}{z} \right) - 4 \frac{\ln(z)}{1 - z} + 6 + 4z \right] \quad (1.18)$$

where the first line only contains distributions. Here, we have introduced the usual $+$ -distributions, which are defined via

$$\int_0^1 dz a(z)_+ f(z) = \int_0^1 dz a(z) (f(z) - f(1)) . \quad (1.19)$$

We review in Section 1.4 again briefly on how these distributions appear, but for the moment we turn to more practical matters as the goal of our course is to get an actual comparison to experimental data. The problem is: distributions are (complicate) math objects, defined only via integral relations, Eqs. (1.17) and (1.19), which are unsuitable for numeric evaluation.

1.4 Next-to-leading order

At LO accuracy, DIS is trivial **FH: add LO diagram** and we find

FH!

$$C_u^{(0)}(z) = e_u^2 \delta(1-z) = C_{\bar{u}}^{(0)}(z) = C_c^{(0)}(z) = C_{\bar{c}}^{(0)}(z) = \dots \quad (1.20)$$

$$C_d^{(0)}(z) = e_d^2 \delta(1-z) = C_{\bar{d}}^{(0)}(z) = C_s^{(0)}(z) = C_{\bar{s}}^{(0)}(z) = \dots \quad (1.21)$$

which, eventually, yields Eq. (1.4).

Starting from next-to-leading order (NLO), the diagrams become more complicated and we encounter new features.

- NLO_q
- NLO_g

We need the Riemann Zeta function $\zeta(n)$ below, which is defined at integer values n by

$$\zeta_n = \zeta(n) = \sum_{j=1}^{\infty} \frac{1}{j^n} \quad (1.22)$$

Chapter 2

DIS in real life

2.1 DIS is diverse

In this section we give an overview many different DIS datasets commonly used in PDF fits. By doing so, we quickly realize that the picture in Chapter 1 is far too simplistic.

2.1.1 HERA

We start with the Hadron-Elektron-Ringanlage (HERA), which was in operation between 1992 and 2007. It was hosted by DESY in Hamburg, Germany. **FH: add HERA image** While the promoted physics discovery, leptoquarks, eventually did not happen, HERA has become one of the most influential DIS machines so far. In fact, it is until this day the only lepton-proton collider in the world. It recorded a huge amount of data, which is still analyzed nowadays, 20 years after the shutdown. HERA had four experiments (H1, ZEUS, HERA-B and HERMES), but we focus here on the combined analysis of H1 and ZEUS [21]. In Table 2.1 we given an overview of the most important fully-inclusive DIS measurements from HERA.

FH!

FH!

FH: complete HERA table

We start by looking at the projectile column and immediately we realize that HERA was most of the time not colliding electrons, but positrons. This is to remind us that DIS is lepton-hadron scattering and, indeed, we have to consider all leptons. We expand our selection further below in Section 2.1.2, but remain for the moment with the electron and the positron, which are exact copies of each other, but with opposite charge.

Table 2.1: HERA measurement [21] used in NNPDF4.0 [2]

projectile	$\sqrt{s}$ [GeV]	x	Q^2	$N_{\text{dat}}^{\text{fit}}/N_{\text{dat}}$	obs.
e^-	318	?	?	?/201-43	σ_r^{NC}
e^+	318	?	?	?	σ_r^{NC}
e^+	300	?	?	?	σ_r^{NC}
e^+	251	?	?	?	σ_r^{NC}
e^+	225	?	?	?	σ_r^{NC}
e^-	318	?	?	?	σ_r^{CC}
e^+	318	?	?	?	σ_r^{CC}

Next, we turn to the energy column $\sqrt{s}$, which spreads across four different energies. In practice the lepton beam was operated with a fixed beam energy ($E_e = 27.5$ GeV) and varying energies for the protons ($E_p = 920$ GeV, 820 GeV, 575 GeV and 460 GeV). If we then look to the x and Q^2 columns and recall Eq. (1.7), we see that the table only indicates the maximal available range of those variables. Specifically, HERA was able to measure activity in the detector for $0.005 < y < 0.95$.

However, not only the accessible kinematical phase space limits our kinematic coverage, but also our requirement to study DIS, conclusion 1.5. In practice, e.g. NNPDF4.0 [2] applies a cut of $Q^2 \geq 8 \text{ GeV}^2$ and $W^2 \geq 12.5 \text{ GeV}^2$. Mathematically, the Q^2 cut ensures we are in a perturbative regime, Eq. (1.15), while the W^2 cut has multiple motivations. First, it ensures we are in an inelastic regime, i.e. we are not finding resonances or excitations of the proton. Second and related to the former, we know it suppresses the correction term $\mathcal{O}\left(\frac{\Lambda_{\text{QCD}}^2}{Q^2}\right)$ in Eq. (1.1). Studying and trying to estimate the correction term is a topic in current research [22–24]. Looking to the column with the number of data points, we note that even after the kinematic cuts we are left with a total of **FH:** $N_{\text{data}}^{\text{HERA}}$ measured cross sections.

Finally, we turn to the observables column for which we see two different entries exist: the reduced neutral current (NC) cross section σ_r^{NC} and the reduced charge current (CC) cross section σ_r^{CC} . The “current” here refers to the current between the leptonic part and the hadronic part, in particular the exchanged boson. The “reduced” cross section refers to a slight rescaling of the double differential cross section, Eq. (1.13),

$$\sigma_r = \frac{xyQ^2}{4\pi\alpha_{\text{em}}^2} \frac{d^2\sigma}{dx dy} \quad (2.1)$$

such that it refers to a specific PDF combination at LO accuracy.

Next, we examine the CC case by recalling two important facts from conclusion 1.1: first, charged leptons, and thus also electrons and positrons, have weak charges and, second, W bosons have an electric charge. Thus, by CC we consider the lepton interacts with the hadron via the exchange of a W boson, which behaves quite different than a photon. First and foremost, due to the electric charge conservation the W boson can only interact with specific leptons and quarks.

2.1.2 Neutrino DIS

2.2 Interpolation grids

2.3 DGLAP

- resummation
- Mellin

2.4 Scale variations

- scheme A/B/C
- MHOU
- covariance matrix

2.5 Heavy Quarks

- F_2^c
- FONLL

2.6 γ_5

- helicity
- HVBM + Larin

2.7 TMC

Chapter 3

More advanced topics

3.1 Positivity

- DIS vs. MSbar

3.2 Higher orders

- IHOU
- N3LO
- channels
- loops and legs

3.3 QED

- Qiu et al.

3.4 Valo

- $C_\gamma \rightarrow k$

Bibliography

- [1] A. Candido, F. Hekhorn, G. Magni, T. R. Rabemananjara, and R. Stegeman, Eur. Phys. J. C **84**, 698 (2024), 2401.15187.
- [2] NNPDF, R. D. Ball *et al.*, Eur. Phys. J. C **82**, 428 (2022), 2109.02653.
- [3] J. Gao, L. Harland-Lang, and J. Rojo, Phys. Rept. **742**, 1 (2018), 1709.04922.
- [4] G. Heinrich, Phys. Rept. **922**, 1 (2021), 2009.00516.
- [5] S. Amoroso *et al.*, Acta Phys. Polon. B **53**, 12 (2022), 2203.13923.
- [6] NNPDF, R. D. Ball *et al.*, Phys. Rev. D **109**, L091501 (2024), 2311.00743.
- [7] E. Leader and E. Predazzi, *An Introduction to gauge theories and modern particle physics. Vol. 1: Electroweak interactions, the new particles and the parton model* (Cambridge University Press, 2011).
- [8] E. Leader and E. Predazzi, *An Introduction to gauge theories and modern particle physics. Vol. 2: CP violation, QCD and hard processes* (Cambridge University Press, 1996).
- [9] M. E. Peskin and D. V. Schroeder, *An Introduction to quantum field theory* (Addison-Wesley, Reading, USA, 1995).
- [10] F. Halzen and A. D. Martin, *QUARKS AND LEPTONS: AN INTRODUCTORY COURSE IN MODERN PARTICLE PHYSICS* (Wiley, 1984).
- [11] A. S. Kronfeld and C. Quigg, Am. J. Phys. **78**, 1081 (2010), 1002.5032.
- [12] A. Accardi *et al.*, Eur. Phys. J. A **52**, 268 (2016), 1212.1701.
- [13] A. Barontini *et al.*, Nnpdf/yadism: v0.13.10: Update n3lo coeff fnc, 2026.
- [14] M. Bertilsson, T. Lappi, H. Mäntysaari, and X.-B. Tong, (2026), 2601.07302.

- [15] C. Casuga and H. Mäntysaari, (2026), 2604.22332.
- [16] H1, V. Andreev *et al.*, Eur. Phys. J. C **75**, 65 (2015), 1406.4709.
- [17] C. Bierlich *et al.*, SciPost Phys. Codeb. **2022**, 8 (2022), 2203.11601.
- [18] I. Helenius, J. O. Laulainen, and C. T. Preuss, (2026), 2605.00502.
- [19] Wikipedia, File:Standard Model of Elementary Particles.svg, online, 2026.
- [20] J. C. Collins, D. E. Soper, and G. F. Sterman, Adv. Ser. Direct. High Energy Phys. **5**, 1 (1989), hep-ph/0409313.
- [21] H1, ZEUS, H. Abramowicz *et al.*, Eur. Phys. J. C **75**, 580 (2015), 1506.06042.
- [22] R. D. Ball, A. Chiefa, and R. Stegeman, Eur. Phys. J. C **86**, 281 (2026), 2511.14387.
- [23] L. A. Harland-Lang, T. Cridge, M. Reader, and R. S. Thorne, Eur. Phys. J. C **86**, 115 (2026), 2510.03753.
- [24] M. Cerutti *et al.*, Phys. Rev. D **111**, 094013 (2025), 2501.06849.