

Deep Inelastic Scattering Revisited

Felix Hekhorn
University of Jyväskylä

Lecture notes for the Jyväskylä Summer School 2026

Abstract

Discuss Yadism [\[1\]](#).

Contents

1	The basics	1
1.1	Introduction	1
1.1.1	Motivation and non-goals	1
1.1.2	Requirements	3
1.1.3	The very basics	4
1.2	Setting the stage	7
1.3	Distributions	9
1.4	Next-to-leading order	10
2	DIS in real life	13
2.1	DIS is diverse	13
2.1.1	HERA	14
2.1.2	Nuclear DIS	17
2.1.3	Neutrino DIS	17
2.2	Interpolation grids	17
2.3	Resummation	19
2.3.1	The strong coupling	20
2.3.2	DGLAP	22
2.4	Scale variations	24
2.4.1	Renormalization scale	24
2.4.2	Factorization scale	27
2.4.3	Combined variation	31
2.4.4	Discussion	32
2.4.5	Implementation	33
2.4.6	Interplay with previous content	35
2.5	Heavy quarks	36
2.5.1	Flavor space	37
2.5.2	ZM-VFNS	38

2.5.3	FFNS	39
2.5.4	Matching schemes	42
2.5.5	FONLL as an example of a GM-VFNS	43
2.5.6	Heavy quark structure functions	43
3	More advanced topics	45
3.1	TMC	45
3.2	γ_5	45
3.3	Positivity	45
3.4	Higher orders	45
3.5	QED	46
3.6	Valo	46
	Bibliography	47

List of Corrections

Add QFT requirements	4
copy DIS diagram from PDG	7
Introduce RSL	10
add LO diagram	10
explain we encounter a certain set of special functions	10
complete HERA table	14
$N_{\text{data}}^{\text{HERA}}$	15
Point to LO DIS exercise?	16
explain interpolation?	19
excercise: proof lambda in alphas?	21
exercise: proof NLO alphas?	22
show exp alphas is LL	22

Chapter 1

The basics

1.1 Introduction

1.1.1 Motivation and non-goals

We aim in this course to make a meaningful comparison between the theoretical predictions and the experimental measurement of a fully inclusive Deeply Inelastic Scattering (DIS) cross section. Instead of an abstract, theoretical discussion about DIS, which can be found in a standard lecture on particle physics, we're looking here to make a connection with real-life physics solving current research questions. The main focus is on Parton Distribution Functions (PDFs), for which DIS plays a major role. In fact, DIS was, is, and will be a very important constrain in determining PDFs - only recently, e.g. starting from NNPDF4.0 [2], LHC data starts to dominate in PDF fits. Stressing the importance and relevance of PDFs for any kind of perturbative QCD (pQCD) application and to understand the inner structure of hadrons is left to the literature [3–5]. In turn, DIS itself can be used to answer unresolved questions about PDFs, e.g. whether the proton has an intrinsic charm component [6, 7].

In order to obtain a more realistic DIS treatment, we review general pQCD features and exemplify them using DIS. In fact, fully inclusive DIS comes in very handy, since here the effects mostly appear in their simplest form and when considering hadronic collisions, e.g. at the LHC, the strategies are mostly the same just more involved. We keep a phenomenological point of view with a specific focus on the numerical implementation as, eventually, we want to compare to the experimental measurements. We refrain from giving detailed proofs in this course and point to the standard textbooks [8–12] for that.

Instead, we consider different features as given and focus on their practical implications. However, a major complication in this course is the interconnection between almost all features and while we discuss them here in turn, we highlight also how they are intertwined with each other.

In recent years (and much more so hopefully in the coming years), DIS has become a major research topic again thanks to the advent of the Electron Ion Collider (EIC) [13]. While the planned physics program of the EIC goes far beyond standard fully inclusive DIS, this simplest case can serve as an introduction into more complicated topics.

The structure of this course follows in large parts the implementation of Yadism (Yet Another DIS module) [1, 14], which is developed and used by the NNPDF collaboration in their PDF determinations. The online documentation, which is available at

<https://yadism.readthedocs.io/> ,

may serve as additional, more technical introduction into the topic. Note that Yadism is an open-source project to which everybody can contribute¹.

Since DIS is a very wide field, it is worth spelling out explicitly some non-goals:

- We only consider pQCD in the collinear framework here. However, because DIS is such a powerful experiment, which offers a clean access to the inner structure of the protons it can be studied using other frameworks as well, such as Color Glass Condensates [15, 16] → ask H. Mäntysaari or T. Lappi
- We only consider fully inclusive cross sections here (to be defined in Section 1.2 and with a tiny exception to be discussed in Section 2.5). In particular, we do not consider any more complicated final states, such as, e.g., single jet or di-jet production [17].
- We only consider pQCD in the collinear framework here. While, Monte Carlo Event generators, such as, e.g., Pythia [18], are rooted in collinear pQCD they eventually go much beyond and we do not consider their details here [19] → ask I. Helenius.
- We only consider DIS here. In particular, we do not consider photo-production or elastic or diffractive cross sections. We define the necessary kinematics for the distinction below in Section 1.1.3.

¹e.g. if some documentation can be improved




Figure 1.1: The Standard Model of Elementary Particles - taken from [20]

- We compute the theoretical predictions here. So while we try to keep the connection to the experimental side close, we do not discuss any specific details about how to measure cross sections.

1.1.2 Requirements

We need some basic knowledge from particle physics and, in particular, you need to be familiar with the Standard Model overview shown in Fig. 1.1. We discuss in this course each and everyone of these particles, with the exception of the three heaviest in each category: the heaviest quark, the top, the heaviest lepton, the tau, and the heaviest boson, the Higgs. They work, in principle, mostly like their lighter brothers and sisters, but in practice are basically always neglected. We give a more thorough justification to discard them when we get to the actual numbers. All other particles explicitly appear in this course and you need to know their charges - see also Fig. 1.1.

Conclusion 1.1: Summary forces, particles, and charges

- The carrier of the strong force is the gluon
- Quarks have a strong charge, leptons don't
- The carriers of the electro-weak force are the photon, the Z boson, and the W boson
- Quarks and charged leptons have electro-magnetic charges
- Quarks and leptons have weak charges

FH!

We also need some basic knowledge from Quantum Field Theory (QFT). FH: Add QFT requirements

1.1.3 The very basics

Let's start from the top:

Conclusion 1.2: DIS in words

fully inclusive DIS = fully inclusive deeply inelastic scattering

What does that actually mean? We begin from the back:

- “scattering” means, of course, we are scattering two particles with each other. In this particular case, we are scattering a lepton off a hadron and to make things simple in the beginning we consider an electron is scattered off a proton.
- “inelastic” means we are breaking the target, the proton, apart; it shatters into many pieces.
- “deeply” is an adverb² to inelastic and it means we are looking “deep” into the inside of the proton. And in order to resolve small things, as usual, we need an highly energetic probe.
- “fully inclusive” means we are observing a single particle in the final state, which is the electron, but nothing else.

Thus, we can write the same information again with a formula

²thus it is “deeply inelastic” or “deep-inelastic”, but not a list of adjectives “deep inelastic”

Conclusion 1.3: DIS in a formula

fully inclusive DIS = $e^-(k) + p(P) \rightarrow e^-(k') + X$ with $-(k - k')^2 \gg \Lambda_{\text{QCD}}^2$

where we have assigned the four-momenta k, P, k' to the incoming electron, the incoming proton and the outgoing electron respectively. As usual X marks the inclusiveness, i.e. we don't observe anything else apart from the electron. By requiring a large momentum-transfer between the incoming and the outgoing electron, i.e. their difference four-momentum vector has a large magnitude, we probe the proton with a highly energetic parton (which we can think of as photon for now).

Next, we need a way to actually compute theory predictions for the cross sections, which you can measure in an experiment. We use the factorization theorem [21],

Conclusion 1.4: Factorization Theorem

$$\sigma(x, Q^2) = \sum_j \int_x^1 \frac{dz}{z} C_j(z, Q^2) f_j(x/z, Q^2) + \mathcal{O}\left(\frac{\Lambda_{\text{QCD}}^2}{Q^2}\right) \quad (1.1)$$

$$= \sum_j (C_j(Q^2) \otimes f_j(Q^2))(x) + \mathcal{O}\left(\frac{\Lambda_{\text{QCD}}^2}{Q^2}\right) \quad (1.2)$$

$$= (\mathbf{C}^T(Q^2) \otimes \mathbf{f}(Q^2))(x) + \mathcal{O}\left(\frac{\Lambda_{\text{QCD}}^2}{Q^2}\right), \quad (1.3)$$

which states how to compute a cross section $\sigma(x, Q^2)$ from coefficient function $\mathbf{C}(z, Q^2)$, computable in pQCD, and a universal PDF $\mathbf{f}(\xi, Q^2)$. In the remaining part of the course we discuss, enhance and correct this formula in many different ways and basically challenge every aspect. We define each and every variable and function in Section 1.2, so we can focus for the moment on two other important aspects: universality and notation.

The crucial, physical aspect about Eq. (1.1) is universality: for any cross section σ measured at any x and Q^2 there is exactly one PDF $\mathbf{f}(\xi, Q^2)$. Instead, the coefficient functions $\mathbf{C}(z, Q^2)$ may depend on the definition of the cross section, but they are computable from first principles. This implies the predictive powers of our framework: if $\mathbf{C}$ is fixed and $\mathbf{f}$ is universal, we can measure σ in one set of experiments and then predict the outcome of another set of measurements σ' .

The other, more practical aspect of Eqs. (1.1) to (1.3) is the short-hand notation we introduce there. While Eq. (1.1) is the most canonical form of the factorization theorem, which can be found like this in most textbooks, we introduce in Eq. (1.2) the multiplicative convolution symbol $\otimes$. Note that we have suppressed the variable over which the convolution is performed and recall that convolution is associative, i.e. $(f \otimes g) \otimes h = f \otimes (g \otimes h)$. Finally, in Eq. (1.3) we introduce the vector notation, where we denote objects, which have a non-trivial dependence on the parton flavor j , with bold font. We use vector notation and the index notation interchangeably depending on the context. Equation (1.3) looks nice and compact, but it worth recalling that it already hides a lot of details: specifically, $C_j(z, Q^2)$ is an object which depends on three different variables.

To Leading Order (LO) accuracy, the coefficient functions $\mathbf{C}$ greatly simplify and we find

$$\sigma(x, Q^2) = \sum_j e_{q_j}^2 f_j(x, Q^2), \quad (1.4)$$

i.e. the cross section is given by the sum of the quark PDFs weighted by their electrical charges squared. Note that here the variables, which are used for the experimentally measured cross section on the left-hand-side, are also used as arguments of the PDFs on the right-hand-side. This is a pure coincidence only valid in the simplest case at LO and two historical comments can be made here. First, when studying older literature this equivalence can lead to confusion as, e.g., PDFs have been also called “structure functions” at the time, or what exactly we mean when talking about a general variable “ x ”. Second, in the so-called “parton model”, which was popular before QCD was invented, the index j just referred to a “parton” with some properties. Instead, within QCD we now know exactly, what the object in question is: a quark, which has a strict definition in QFT with well-defined properties and quantum numbers. For simplicity, we still call an object, which can be either a quark or a gluon “parton”.

1.2 Setting the stage

In the following section we collect and define all ingredients needed for Eq. (1.1). As already said, DIS is given by

$$e^-(k) + p(P) \rightarrow e^-(k') + X \quad (1.5)$$

FH: copy DIS diagram from PDG where the four-momenta k, P, k' belong to the incoming electron, the incoming proton and the outgoing electron respectively. This allows us, first, to define four momentum of the exchanged photon **FH!**

$$q = k - k' \quad (1.6)$$

and from there the set of four common variables:

$$s_l = (k + P)^2 = \frac{Q^2}{xy} \quad (1.7)$$

$$Q^2 = -q^2 \quad (1.8)$$

$$x = \frac{Q^2}{2q \cdot P} \quad (1.9)$$

$$y = \frac{q \cdot P}{k \cdot P} \quad (1.10)$$

We refer to Q^2 as virtuality (of the exchanged boson), to x as Bjorken- x (named after James Bjorken), and to y as inelasticity. Note that Eq. (1.7) gives an explicit relation among these, thus we have only three independent and one dependent variable. In practice, the invariant mass of the electron-proton system s_l is fixed by the experimental setup, i.e. by the energy of the colliding particles. Also the measurable range of y is in practice limited by the experimental geometry. DIS cross sections are typically provided as double-differential cross sections in x and y . Finally, it is convenient to define the mass squared of the system X recoiling against the scatter electron,

$$W^2 = (P + q)^2 = Q^2 \left(\frac{1}{x} - 1 \right) \quad (1.11)$$

Conclusion 1.5: DIS requirements

We require $Q^2 \gg \Lambda_{\text{QCD}}^2$ (deep) and $W^2 \gg \Lambda_{\text{QCD}}^2$ (inelastic) to call a lepton-hadron scattering DIS.

We can decompose the (double-differential) cross section into a leptonic interaction and a hadronic interaction, yielding each a tensor in Minkowski space, $L_{\mu\nu}$ and $W_{\mu\nu}$ respectively. Thus, we can write

$$\frac{d^2\sigma}{dx dy} = \frac{2\pi y \alpha_{\text{em}}^2}{Q^4} L_{\mu\nu} W^{\mu\nu} \quad (1.12)$$

where we have introduced the electro-magnetic coupling α_{em} , which appears once on the leptonic side and once on the hadronic side. We can express the leptonic tensor using plain Quantum Electrodynamics (QED) and provide a decomposition of the hadronic tensor in terms of scalar functions. Together, we obtain

$$\frac{d^2\sigma}{dx dy} = \frac{4\pi \alpha_{\text{em}}^2}{xy Q^2} \left(\left(1 - y + \frac{y^2 x}{2} \right) F_2(x, Q^2) - \frac{y^2 x}{2} F_L(x, Q^2) \right) \quad (1.13)$$

where we have introduced the two common scalar functions, now called structure functions F_2 and F_L , which are often measured in experiments. It is convenient to also define a third, linearly dependent structure function via

$$2xF_1(x, Q^2) = F_2(x, Q^2) - F_L(x, Q^2). \quad (1.14)$$

Keep in mind that the discussion so far is a major simplification of DIS and in Section 2.1 we reintroduce the main complexity.

Let us continue to collect the remaining ingredient for Eq. (1.1):

- The sum over partons j includes quarks and gluons and we refine in Section 2.5 how large the sum actually is.
- The parton momentum fraction $\xi = x/z$ is the fraction of the four momentum the parton carries with respect to its parent proton.
- The probability of such a configuration to occur is given by the PDF, i.e. $f_j(\xi, Q^2)d\xi$ is the probability to find a parton j with momentum ξP inside a proton with momentum P at an experiment performed at the scale Q^2 .
- The integration over z sums all configuration to generate the final state configuration.

Finally, the coefficient functions $\mathbf{C}(z, Q^2)$ are computable order-by-order in pQCD and thus can be decomposed as

$$\mathbf{C}(z, Q^2) = \sum_{k=0} \alpha_s^k(Q^2) \mathbf{C}^{(k)}(z). \quad (1.15)$$

Note that the Q^2 dependency of $\mathbf{C}(z, Q^2)$ is fully captured by the strong coupling $\alpha_s(Q^2)$.

1.3 Distributions

As we have seen the coefficient functions may contain distributions. For example the quark coefficient function at LO is just a Dirac delta distribution:

$$\bar{C}_q^{(0)}(z) = \delta(1 - z). \quad (1.16)$$

To better compare with our next step, let's repeat the definition of the Dirac delta distribution:

$$\int_0^1 dz \delta(1 - z) f(z) = f(1). \quad (1.17)$$

However, this is only the simplest case, which appears at LO, and at higher order we get more complicated distributions. For example the quark coefficient function at NLO is given by

$$\begin{aligned} \bar{C}_q^{(1)}(z) = C_F \left[4 \left(\frac{\ln(1 - z)}{1 - z} \right)_+ - 3 \left(\frac{1}{1 - z} \right)_+ - (9 + 4\zeta_2) \delta(1 - z) \right. \\ \left. - 2(1 + z) \ln \left(\frac{1 - z}{z} \right) - 4 \frac{\ln(z)}{1 - z} + 6 + 4z \right] \end{aligned} \quad (1.18)$$

where the first line only contains distributions. Here, we have introduced the usual $+$ -distributions, which are defined via

$$\int_0^1 dz a(z)_+ f(z) = \int_0^1 dz a(z) (f(z) - f(1)). \quad (1.19)$$

We review in Section 1.4 again briefly on how these distributions appear, but for the moment we turn to more practical matters as the goal of our course is to get an actual

Figure 1.2: LO DIS Feynman diagram

comparison to experimental data. The problem is: distributions are (complicate) math objects, defined only via integral relations, Eqs. (1.17) and (1.19), which are unsuitable for numeric evaluation.

FH!

FH: Introduce RSL

1.4 Next-to-leading order

FH!

At LO accuracy, DIS is trivial FH: add LO diagram and we find

$$C_q^{(0)}(z) = C_{\bar{q}}^{(0)}(z) = e_q^2 \bar{C}_q^{(0)}(z) \quad \forall q, \quad C_g^{(0)}(z) = 0 \quad (1.20)$$

where q represents any quark and g represent the gluon. Inserting Eq. (1.20) into Eq. (1.1) yields Eq. (1.4).

Starting from next-to-leading order (NLO), the diagrams become more complicated and we encounter new features.

- NLO_q
- NLO_g

Eventually, we get

$$C_q^{(1)}(z) = C_{\bar{q}}^{(1)}(z) = e_q^2 \bar{C}_q^{(1)}(z) \quad \forall q \quad (1.21)$$

and

$$C_g^{(1)}(z) = \quad (1.22)$$

FH!

FH: explain we encounter a certain set of special functions We need the Riemann Zeta function $\zeta(n)$ below, which is defined at integer values n by

$$\zeta_n = \zeta(n) = \sum_{j=1}^{\infty} \frac{1}{j^n} \quad (1.23)$$

Chapter 2

DIS in real life

2.1 DIS is diverse

In this section we give an overview many different DIS datasets commonly used in PDF fits. By doing so, we quickly realize that the picture in Chapter 1 is far too simplistic.



Figure 2.1: “The view into the 6.3-kilometre-long HERA tunnel shows the superconducting magnets for guiding the protons on top and the electron ring beneath them.” - taken from [22]

Table 2.1: HERA measurement [23] used in NNPDF4.0 [2]

projectile	$\sqrt{s}$ [GeV]	x	Q^2	$N_{\text{dat}}^{\text{fit}}/N_{\text{dat}}$	obs.
e^-	318	?	?	?/201-43	σ_r^{NC}
e^+	318	?	?	?	σ_r^{NC}
e^+	300	?	?	?	σ_r^{NC}
e^+	251	?	?	?	σ_r^{NC}
e^+	225	?	?	?	σ_r^{NC}
e^-	318	?	?	?	σ_r^{CC}
e^+	318	?	?	?	σ_r^{CC}

2.1.1 HERA

We start with the Hadron-Elektron-Ringanlage (HERA), which was in operation between 1992 and 2007. HERA was hosted by DESY in Hamburg, Germany and Fig. 2.1 shows an image from the HERA tunnel. While the promoted physics discovery, leptiquarks, eventually did not happen, HERA has become one of the most influential DIS machines so far. In fact, it is until this day the only lepton-proton collider in the world. It recorded a huge amount of data, which is still analyzed nowadays, 20 years after the shutdown. HERA had four experiments (H1, ZEUS, HERA-B and HERMES), but we focus here on the combined analysis of H1 and ZEUS [23]. In Table 2.1 we given an overview of the most important fully-inclusive DIS measurements from HERA.

FH!

FH: complete HERA table

We start by looking at the projectile column and immediately we realize that HERA was most of the time not colliding electrons, but positrons. This is to remind us that DIS is lepton-hadron scattering and, indeed, we have to consider all leptons. We expand our selection further below in Section 2.1.3, but remain for the moment with the electron and the positron, which are exact copies of each other, but with opposite charge.

Next, we turn to the energy column $\sqrt{s}$, which spreads across four different energies. In practice the lepton beam was operated with a fixed beam energy ($E_e = 27.5$ GeV) and varying energies for the protons ($E_p = 920$ GeV, 820 GeV, 575 GeV and 460 GeV). If we then look to the x and Q^2 columns and recall Eq. (1.7), we see that the table only indicates the maximal available range of those variables. Specifically, HERA was able to measure activity in the detector for $0.005 < y < 0.95$.

However, not only the accessible kinematical phase space limits our kinematic coverage, but also our requirement to study DIS, conclusion 1.5. In practice, e.g. NNPDF4.0 [2] applies a cut of $Q^2 \geq 8 \text{ GeV}^2$ and $W^2 \geq 12.5 \text{ GeV}^2$. Mathematically, the Q^2 cut ensures we are in a perturbative regime, Eq. (1.15), while the W^2 cut has multiple motivations. First, it ensures we are in an inelastic regime, i.e. we are not finding resonances or excitations of the proton. Second and related to the former, we know it suppresses the correction term $\mathcal{O}\left(\frac{\Lambda_{\text{QCD}}^2}{Q^2}\right)$ in Eq. (1.1). Studying and trying to estimate the correction term is a topic in current research [24–26]. Looking to the column with the number of data points, we note that even after the kinematic cuts we are left with a total of FH: $N_{\text{data}}^{\text{HERA}}$ measured cross sections. FH!

Finally, we turn to the observables column for which we see two different entries exist: the reduced neutral current (NC) cross section σ_r^{NC} and the reduced charge current (CC) cross section σ_r^{CC} . The “current” here refers to the current between the leptonic part and the hadronic part, in particular the exchanged boson. The “reduced” cross section refers to a slight rescaling of the double differential cross section, Eq. (1.13),

$$\sigma_r = \frac{xyQ^2}{4\pi\alpha_{\text{em}}^2} \frac{d^2\sigma}{dx dy} \quad (2.1)$$

such that it refers to a specific PDF combination at LO accuracy.

Next, we examine the CC case by recalling two important facts from conclusion 1.1: first, charged leptons, and thus also electrons and positrons, have weak charges and, second, W bosons have an electric charge. Thus, by CC we consider the lepton interacts with the hadron via the exchange of a W boson, which behaves quite different than a photon. First and foremost, due to the electric charge conservation the W boson can only interact with specific leptons and quarks. On the leptonic side this implies that, first, e.g., the positron always emits a W^+ boson, and, second, the recoiling lepton is a neutrino, which in practice is impossible to track. Specifically, it is impossible to measure its momentum four-vector k' , Eq. (1.5), and thus measure the W boson momentum q , Eq. (1.6), or any other derived variable (x, Q^2, y) . Instead, the full reconstruction of the hadronic final state also allows to extract the relevant kinematic variables, which is used instead [23]. Although the production of CC events is suppressed by the propagator of the W boson, $\sim (Q^2 + m_W^2)^{-1}$, it is a LO effect and thus no further suppression. Moreover, since the W are maximally parity violating this allows a third, independent structure function (typically referred to as F_3) to appear in the decomposition of the

cross section, Eq. (1.13). We review the problems introduced by γ_5 , which causes the parity violation, further in Section 3.2.

We now consider the NC case, which is a generalization of the photon exchange discussed in Chapter 1. As we recall from conclusion 1.1 the Z boson is practically a massive copy of the photon γ and they share all other quantum numbers. This implies whenever a photon is exchanged, also a Z boson can be exchanged, with the main difference that its contributions are suppressed by the respective propagator, $\sim (Q^2 + m_Z^2)^{-1}$. More specifically, we get three contributions to our final cross section: a pure photon contribution σ^γ (where the superscript γ is often suppressed), a pure Z contribution σ^Z , and an interference contribution¹ $\sigma^{\gamma Z}$, which have to be summed together:

$$\sigma^{\text{NC}} = \sigma^\gamma + \sigma^{\gamma Z} + \sigma^Z. \quad (2.2)$$

While at low virtualities $Q^2 \ll m_Z^2$ the Z contributions can be neglected, this does not apply to the largest HERA measurements. Similar to W boson, also Z bosons have an axial-vectorial coupling to fermions, $\sim \gamma^\mu \gamma^5$, which introduces a contribution from F_3 to Eq. (1.13). The interference contribution, $F_{1,2}^{\gamma Z}$, yields an electric charge (from the photon) and a vectorial coupling (from the Z boson) both on the leptonic and the hadronic side, whereas the pure Z contribution, $F_{1,2}^Z$, yields squares of both vectorial and axial-vectorial couplings. On the other side, as F_3 is per-se a parity-violating observable, there the interference contribution $F_3^{\gamma Z}$ yields an electric charge (from the photon) and an axial-vectorial coupling (from the Z) and the pure Z contribution, F_3^Z yields a product of the vectorial and axial-vectorial couplings.

The fact that NC and CC couple completely different to fermions, i.e. quarks and leptons, and the definition of three independent structure functions in both cases allows one to fully determine PDFs from DIS [27]. **FH: Point to LO DIS exercise?**

FH!

Let us return to the key point of this section here. Based on the discussion above, we can now better justify why we don't consider all particles in Fig. 1.1. According on our definition of DIS, conclusion 1.3, in theory also the Higgs boson would contribute to NC cross section, but since its coupling is proportional to the mass of the particle and we always² consider massless leptons it does in practice not contribute. As the tau has a very short lifetime they are in practice impossible to accelerate in a collider and

¹the interference term typically contains the required factor of 2 from the binomial formula inside

²almost always [28] - but even so Higgs would only contribute a very large Q^2

thus not suitable for DIS collider experiments. Instead, while muons are also difficult to accelerate concrete experiments for muon-proton scattering have been proposed [29].

2.1.2 Nuclear DIS

After having considered all available bosons and a few more leptons, we now need to consider more hadrons. While we focus here on the proton PDF point-of-view, DIS is equally important for nuclear PDFs [30–33], which describe the internal structure of nuclei. Moreover, the two are typically directly intertwined: proton PDF extractions often contain measurements on nuclear target [2], while nuclear PDF extraction often require a proton baseline.

2.1.3 Neutrino DIS

- CHORUS, Sami
- FASER
- astro

2.2 Interpolation grids

Since PDFs are non-perturbative objects, encoding low-scale physics, they are not computable from first principles inside pQCD. Instead, PDFs are obtained from a fitting procedure, which can be summarized as follows:

Conclusion 2.1: PDF fitting

1. Guess a candidate PDF
2. Compute the theory predictions with this candidate PDF
3. Compare the theory predictions to the experimental measurement

Repeat the above steps until the procedure converges, i.e. the comparison is “good enough”.

The specific details on how to guess a candidate PDF, how to exactly compute the theory predictions, which experimental data to use, and what “good enough” exactly means is a choice. That is the reason why competing PDF fitting groups exist and why they do not necessarily obtain the same PDF, although we all assume the latter to be universal.

The main point for us here is that we need to compute theory predictions very many times for slightly different PDFs. This is a serious problem for complicated hadronic observables, such as, e.g., jet cross sections [34], where the computation can take many hours. The solution is to compute fast interpolation grids, using, e.g., PineAPPL [35, 36], APPLgrid [37], or fastNLO [38], which store the partonic matrix elements in an efficient way, such that any subsequent convolution with an arbitrary PDF is fast.

In DIS this is not a very big problem, since typically coefficient functions, Eq. (1.1), are known analytically and thus the convolution with PDFs is straightforward. However, PineAPPL allows also DIS cross sections to be stored thus we can use this simplified case to introduce the general concept here. Moreover, having all observables in a PDF fit sharing the same technology also simplifies their consistent treatment and inside the NNPDF collaboration this is realized in the pipeline framework [39].

Interpolation grids are only a discretized version of the coefficient functions which we can recombine in a suitable way a posteriori. Specifically, they are higher dimensional tensors on which we perform fast linear algebra operations. The first three major dimensions are:

bin This is the list of cross section that we need to compute simultaneously. In DIS this typically corresponds to all N_{dat} data point measured in a given configuration, e.g. in Table 2.1. Thus, in practice the actual list is given by the experimental measurement.

order This is the list of perturbative orders, i.e. we store each $\mathbf{C}^{(k)}(z)$, Eq. (1.15), separately. Having this list available allows to use a given grid also at lower orders and thus study, e.g., perturbative stability. In practice, the actual list is given by the requested accuracy of the user (provided we can actual fulfill that request). Recall that LO, Eq. (1.16), and NLO, e.g. Eq. (1.18), are quite different and this generates correlations with the momentum fraction dimension (see below).

channel This is the list of PDF flavor combinations, i.e. the sum over j in Eq. (1.1). Having this list available allows to study the impact of different PDF combination

and, e.g., determine which PDF the observable can potentially constrain most. In DIS it is given by the coupling structure of the exchanged boson, i.e. γ/Z vs. $W^\pm$. In practice, the actual list is the easiest given directly in terms of individual parton flavors as the available combinations are too complex (see Section 2.4 for additional complications).

These three dimensions are present in any interpolation grid and in DIS there are two additional dimensions:

scale This is the list of factorization and renormalization scales, discussed in Section 2.4.

In DIS this dimension turns out to be trivial, i.e. it has support by a single value, as in fully inclusive DIS there is only a single scale, Q^2 , available and we set typically $\mu_F^2 = \mu_R^2 = Q^2$. In practice, the actual list is given by the Q^2 specified in the bin configuration.

momentum fraction This is the list of momentum fractions z , Eq. (1.1), over which the coefficient functions are convolved with the PDF. In DIS we are facing the problem of distributions, see Section 1.3, i.e. we can not directly tabulate the coefficient functions themselves. In practice, the actual list is given by a convolution with a suitable set of interpolation functions, which form a basis for PDFs.

For fully inclusive DIS we end up with five dimensional tensors, where each inner dimension may depend on the outer.

FH!

FH: explain interpolation?

2.3 Resummation

In preparation to Section 2.4 we need to review two main ingredients into pQCD: the strong coupling $\alpha_s(Q^2)$, Eq. (1.15), and the PDFs $f(\xi, Q^2)$, Eq. (1.1). Specifically, we need to remind ourselves on their scale dependency (also referred to as evolution) if we want to vary them in Section 2.4. While the two objects obey very similar equations, the strong coupling $\alpha_s(Q^2)$ is conceptually simpler and we start with that.

2.3.1 The strong coupling

The strong coupling α_s is directly linked to the coupling of the gluons to quarks,

$$\mathcal{L}_{\text{QCD}} \supset g T_a \bar{\psi} \gamma^\mu A_\mu^a \psi, \quad \alpha_s = \frac{g^2}{4\pi} \quad (2.3)$$

and is the central object in pQCD around which we organize our perturbative series. If one considers higher order corrections to the gluon-quark couplings, one encounters quickly divergencies of ultra-violet origin, i.e. for large loop momenta. This forces the introduction of a regularization procedure, typically dimensional regularization, which eventually leads to a scale dependent definition of the renormalized strong coupling $\alpha_s(\mu_R^2)$. The introduced scale, called renormalization scale μ_R henceforth, is an arbitrary scale, which is an artifact of our regularization procedure and which effectively represents a separation between perturbative and non-perturbative regimes. However, the scale dependence of $\alpha_s(\mu_R^2)$ is rigorously predicted in pQCD, organizing the terms in such a way that within the requested accuracy any dependence of μ_R is cancelled. Recall that the scale dependence can be computed once the Lagrangian $\mathcal{L}$ is defined and does not require any actual observable.

Eventually, we obtain the renormalization group equation (RGE) of the strong coupling,

$$\mu_R^2 \frac{d\alpha_s(\mu_R^2)}{d\mu_R^2} = \beta(\alpha_s) = - \sum_{j=0} \beta_j(\alpha_s(\mu_R^2))^{2+j} \quad (2.4)$$

which is also the definition³ of the beta function $\beta(\alpha_s)$. The upper limit of the sum in Eq. (2.4) defines the perturbative accuracy, to which all other objects in pQCD refer to.

At LO accuracy we find

$$\mu_R^2 \frac{d\alpha_s(\mu_R^2)}{d\mu_R^2} = -\beta_0(\alpha_s(\mu_R^2))^2, \quad \beta_0 = \frac{11C_A - 4T_R n_f}{12\pi} \quad (2.5)$$

where we used the second Casimir constant of the adjoint representation $C_A = N_c = 3$, the normalization of the fundamental generators $T_R = 1/2$, and the number of active

³Note that in the literature many conventions for the left- and/or right-hand-side exist.

flavors n_f , discussed in Section 2.5. In this specific case we can solve Eq. (2.5) and obtain

$$\alpha_s^{\text{LO}}(\mu_R^2) = \frac{\alpha_s(\mu_{R,0}^2)}{1 + \alpha_s(\mu_{R,0}^2)\beta_0 \ln(\mu_R^2/\mu_{R,0}^2)} \quad (2.6)$$

for a given boundary condition $\alpha_s(\mu_{R,0}^2)$ at the scale $\mu_{R,0}^2$. Recall, that Eq. (2.6) has not two free parameters, $\mu_{R,0}$ and $\alpha_s(\mu_{R,0})$ as one would naively think, but only one Λ_{QCD} . Also remember that Λ_{QCD} corresponds to the pole in Eq. (2.6) and pQCD only makes sense above that scale, $\mu_R > \Lambda_{\text{QCD}}$. **FH: exercise: proof lambda in alphas?**

FH!

Now, coming back to the title of this section, note that Eq. (2.6) is a resummation:

$$\alpha_s^{\text{LO}}(\mu_R^2) = \alpha_s(\mu_{R,0}^2) \sum_{k=0}^{\infty} \left(-\alpha_s(\mu_{R,0}^2)\beta_0 \ln(\mu_R^2/\mu_{R,0}^2) \right)^k \quad (2.7)$$

i.e. it collects an infinite tower of logarithms, here $\ln(\mu_R^2/\mu_{R,0}^2)$, into a compact expression, Eq. (2.6). Since $\mu_{R,0}$ is an arbitrary scale, often chosen to coincide with the mass of the Z boson $\mu_{R,0} = m_Z$, it is basically always required to perform the resummation and use Eq. (2.6). However, in some situations, to be discussed in Section 2.4, we need Eq. (2.6) truncated to a finite order, i.e. unraveling Eq. (2.7), and we get

$$\alpha_s^{\text{LO}}(\mu_R^2) = \alpha_s(\mu_{R,0}^2) - (\alpha_s(\mu_{R,0}^2))^2 \beta_0 \ln(\mu_R^2/\mu_{R,0}^2) + \dots \quad (2.8)$$

While for the special case $\mu_{R,0} = \mu_R$ we need to be consistent of course and we thus find that $\alpha_s^{\text{LO}}(\mu_R^2)$ is $\alpha_s(\mu_{R,0}^2)$ at the LO truncation, we note that they differ by higher order terms. Equation (2.8) is the first, simplest case of a perturbative expansion of a resummed expression, Eq. (2.6), and can also be obtained by using the Taylor expansion of $\alpha_s(\mu_R^2)$ around $\mu_{R,0}^2$ together with Eq. (2.5). In other words, Eq. (2.8) is a perturbative expansion of Eq. (2.6) in powers of $\alpha_s(\mu_{R,0}^2)$. Note that Eq. (2.7) has a finite radius of convergence (as all geometric series) and that is also reflected in Eq. (2.6) by the pole Λ_{QCD} .

Starting from NLO accuracy the RGE, e.g. in this specific case given by

$$\mu_R^2 \frac{d\alpha_s(\mu_R^2)}{d\mu_R^2} = -\beta_0 (\alpha_s(\mu_R^2))^2 - \beta_1 (\alpha_s(\mu_R^2))^3, \quad (2.9)$$

can no longer be solved exactly with elementary functions. However, it is still possible to find an analytic solution which solves Eq. (2.9) at the requested accuracy, i.e. in this

FH! case NLO. FH: exercise: proof NLO alphas? This solution still performs a resummation,
 FH! as we have to, but it only collects the so-called leading logarithms FH: show exp alphas
 is LL A completely different way to solve Eq. (2.9) is using numerical tools instead. We
 refer to the literature on how to implement a specific solution in practice and remain
 here just with the fact that it is a complicated problem on its own, which has no unique
 solution and for which approximations are mandatory.

Conclusion 2.2: Solution of RGE

- The LO RGE of the strong coupling, Eq. (2.5), can be solved exactly, Eq. (2.6)
- The RGE of the strong coupling at higher orders, e.g. Eq. (2.9), can not be solved exactly and one needs to choose a solution method

Any proper solution of the RGE must solve the RGE itself at the requested accuracy else it is not a solution, but they can and do differ by terms beyond - we refer to the uncertainty which is associated with this freedom as Missing Higher Order Uncertainty (MHOU). Truncating the perturbative series is a fundamental part of pQCD and thus we are always bound to the accuracy which we request at the beginning. While in many simple application MHOU are often neglected they are an important ingredient in modern PDF fits [40] and in Section 2.4 we demonstrate one possible way to estimate them.

2.3.2 DGLAP

For PDFs $\mathbf{f}(\xi, \mu_F^2)$ the situation is essentially the same as for the strong coupling $\alpha_s(\mu_R^2)$, except the latter is just a number and the former are a set of functions. The scale dependence is again predicted by pQCD, but now we need an arbitrary physical observable to do so. The specific observable does not matter since PDFs are universal, but due to the linear dependence on PDF and the “simplicity” of fully inclusive DIS it often serves as reference case. The arbitrary scale that is introduced in the regularization procedure, typically called collinear factorization, is referred to as factorization scale μ_F henceforth.

Eventually, we obtain the RGE of PDFs, often referred to as DGLAP equation named after the researcher who established it [41–43],

$$\mu_F^2 \frac{d\mathbf{f}(\xi, \mu_F^2)}{d\mu_F^2} = (\mathbf{P}(\mu_F^2) \otimes \mathbf{f}(\mu_F^2))(\xi) = \sum_{j=0} (\alpha_s(\mu_F^2))^{1+j} (\mathbf{P}^{(j)} \otimes \mathbf{f}(\mu_F^2))(\xi) \quad (2.10)$$

where $\mathbf{P}(y, \mu_F^2)$ is the matrix of Altarelli-Parisi splitting functions. Note that the upper limit of the sum in Eq. (2.10) must be set to the requested perturbative accuracy, i.e. consistent with Eq. (2.5).

When attempting to solve Eq. (2.10) we must worry about the two complications in PDFs with respect of the strong coupling. First, since PDFs are a vector and thus the Altarelli-Parisi splitting functions are combined in a matrix $\mathbf{P}$, we need to worry about non-commutativity. Indeed, while at LO accuracy we only have a single matrix $\mathbf{P}^{(0)}$, which trivially commutes with itself, starting from NLO we encounter a second matrix $\mathbf{P}^{(1)}$, which does not commute with the former. Second, at any fixed scale PDFs are a function of the momentum fraction ξ and, moreover, the right-hand-side of Eq. (2.10) features an integral (with the definition of the convolution $\otimes$ in Eqs. (1.1) and (1.2)).

To overcome the second problem, one possibility is to apply an integral transformation, the so-called Mellin transformation, defined by

$$\tilde{\mathbf{f}}(N, \mu_F^2) = \int_0^1 d\xi \xi^{N-1} \mathbf{f}(\xi, \mu_F^2) \quad (2.11)$$

$$\gamma(N, \mu_F^2) = \int_0^1 dy y^{N-1} \mathbf{P}(y, \mu_F^2) \quad (2.12)$$

where the Mellin-transformed splitting functions γ are often referred to as anomalous dimensions. Inserting Eqs. (2.11) and (2.12) into Eq. (2.10) yields a “simple” first order differential equation, which can be solved by a “simple” exponentiation. However, in practice PDFs are mostly needed in momentum fraction space, $\mathbf{f}(\xi, \mu_F^2)$, and not in Mellin space, $\tilde{\mathbf{f}}(N, \mu_F^2)$, thus if we use the Mellin transform, we must also use its inverse transformation.

We can write the general solution to Eq. (2.10) in terms of an Evolution Kernel Operator (EKO) [44] $\mathbf{E}$,

$$\mathbf{f}(\xi, \mu_F^2) = (\mathbf{E}(\mu_F^2 \leftarrow \mu_{F,0}^2) \otimes \mathbf{f}(\mu_{F,0}^2))(\xi), \quad (2.13)$$

but we skip spelling out an explicit solution here and refer to the literature instead [44]. The important part for us is that the operator (which is almost an exponential) again captures a resummation, in this case of $\ln(\mu_F^2/\mu_{F,0}^2)$. Again, we can perform a perturbative

expansion of the LO resummed solution to see the logarithm explicitly,

$$\mathbf{f}^{\text{LO}}(\xi, \mu_F^2) = \mathbf{f}(\xi, \mu_{F,0}^2) + \alpha_s(\mu_{F,0}^2) \ln(\mu_F^2/\mu_{F,0}^2) \left(\mathbf{P}^{(0)} \otimes \mathbf{f}(\mu_{F,0}^2) \right)(\xi) + \dots \quad (2.14)$$

and starting from NLO accuracy, we face the same issues as for the strong coupling and conclusion 2.2 applies in the same way.

2.4 Scale variations

We now attempt to address a problem we first see in Section 2.3: Missing Higher Order Uncertainty (MHOU). We repeat again, the truncation of the perturbative series is a fundamental part of our framework and we are bound to it. The naive solution of just computing the next higher correction is not a solution, as we would face the problem just again there. Thus the task is to get an estimate of the missing terms based on current knowledge, which by definition is an ill-posed problem. Several possible strategies have been suggested in the literature [40, 45–51], which are based on resummation of specific variables and/or statistical arguments.

From the practical point of view as a PDF fitter, however, there is one big problem with many of the proposed algorithms: they only work for some specific set of observables and require a detailed analysis of the kinematics. Thus, we review here the most common approach to estimate MHOU, which can be rigorously derived in pQCD and applied to any observable: scale variations. On the other side, especially in recent years, scale variations have been challenged as suitable tool, due to the inherent arbitrariness and the lack of statistical interpretation (to which we return in the end). For now we focus on their practical implementation in DIS.

2.4.1 Renormalization scale

We start again from the simpler case of the renormalization scale μ_R associated with the strong coupling $\alpha_s(\mu_R^2)$ via the RGE, Eq. (2.5). As we say above, the renormalization scale captures the ultra-violet divergencies, which arise, e.g., from self-energy diagrams and which are thus associated with the partonic diagram itself. Moreover, as we see from Eq. (1.15), we expand the partonic coefficient functions in powers of the strong coupling.

Conclusion 2.3: Renormalization scale dependence

The renormalization scale dependence is captured by the partonic coefficient function $\mathbf{C}(z, Q^2, \mu_R^2)$.

Since we have to choose something for the arbitrary scale μ_R and we only have a single scale available, the virtuality Q^2 , we start by setting them equal to each other $\mu_R^2 = Q^2$, which is also the implicit choice in Eq. (1.1). We then need to find a way to estimate the missing terms in $\mathbf{C}(z, Q^2, \mu_R^2 = Q^2)$, i.e. something that is identical at the requested perturbative accuracy, but differs from it beyond. This can be achieved by using Eq. (2.8) together with a suitable redefinition of the perturbative coefficients of the partonic coefficient function $\mathbf{C}^{(j)}(z)$. Specifically, we require

$$\mathbf{C}(z, Q^2, \mu_R^2) = \mathbf{C}(z, Q^2, \mu_R^2 = Q^2)(1 + \mathcal{O}(\alpha_s(Q^2))) \quad (2.15)$$

with

$$\mathbf{C}(z, Q^2, \mu_R^2) = \sum_{j=0} (\alpha_s(\mu_R^2))^j \mathbf{C}^{(j)}(z, \ln(\mu_R^2/Q^2)). \quad (2.16)$$

Before we continue with deriving explicit expressions for $\mathbf{C}^{(j)}(z, \ln(\mu_R^2/Q^2))$ some comments are in order:

- $\mathbf{C}(z, Q^2)$ in Eq. (1.1) is an abbreviation of $\mathbf{C}(z, Q^2, \mu_R^2 = Q^2)$ and $\mathbf{C}^{(j)}(z)$ of $\mathbf{C}^{(j)}(z, 0)$
- in Eq. (2.16) μ_R has its intended meaning: it is the scale of the strong coupling
- from Section 2.3.1 we know $\alpha_s(\mu_R^2)$ is a pure resummation of logarithms, thus $\mathbf{C}^{(j)}$ may only depend on $\ln(\mu_R^2/Q^2)$
- $\mathbf{C}^{(j)}(z, 0)$ can be computed from diagrams by assuming explicitly $\mu_R = Q$ there

We now derive the explicit next-to-next-to-leading order (NNLO) expression for fully inclusive DIS. In particular, we recall that LO is given by $\mathbf{C}^{(0)}(z)$, i.e. proportional to $(\alpha_s(Q^2))^0$ in Eq. (1.15), NLO is given by $\mathbf{C}^{(1)}(z)$ proportional to $(\alpha_s(Q^2))^1$, and NNLO by $\mathbf{C}^{(2)}(z)$ proportional to $(\alpha_s(Q^2))^2$. Moreover, we assume that the coefficient functions at the central scale $\mu_R = Q$ are known and we can write Eq. (1.15) up to NNLO

$$\mathbf{C}(z, Q^2) = \mathbf{C}^{(0)}(z) + \alpha_s(Q^2)\mathbf{C}^{(1)}(z) + (\alpha_s(Q^2))^2\mathbf{C}^{(2)}(z) \quad (2.17)$$

Expanding Eq. (2.16) up to the same accuracy we get

$$\begin{aligned} \mathbf{C}(z, Q^2, \mu_R^2) &= \mathbf{C}^{(0)}(z, \ln(\mu_R^2/Q^2)) + \alpha_s(\mu_R^2) \mathbf{C}^{(1)}(z, \ln(\mu_R^2/Q^2)) \\ &\quad + (\alpha_s(\mu_R^2))^2 \mathbf{C}^{(2)}(z, \ln(\mu_R^2/Q^2)) \end{aligned} \quad (2.18)$$

We are now left with the task to make Eq. (2.17) and Eq. (2.18) perturbatively equivalent, and although we are considering NNLO the equivalence must hold at any lower order. Thus, we first can consider LO: since there is no dependency on the strong coupling in the first place, the scale-varied coefficient function may not depend on the renormalization scale and we find

$$\mathbf{C}^{(0)}(z, \ln(\mu_R^2/Q^2)) = \mathbf{C}^{(0)}(z). \quad (2.19)$$

Starting from NLO an explicit dependency on the strong coupling appears, however, from Eq. (2.8) we know that the two choices, $\alpha_s(Q^2)$ and $\alpha_s(\mu_R^2)$ differ by higher order terms. Specifically, we have

$$\alpha_s(Q^2) = \alpha_s(\mu_R^2) + \mathcal{O}\left((\alpha_s(\mu_R^2))^2\right) \quad \text{and} \quad \alpha_s(\mu_R^2) = \alpha_s(Q^2) + \mathcal{O}\left((\alpha_s(Q^2))^2\right) \quad (2.20)$$

Thus, if Eq. (2.17) and Eq. (2.18) have to be the same up to $\mathcal{O}\left((\alpha_s(Q^2))^2\right)$, we find

$$\mathbf{C}^{(1)}(z, \ln(\mu_R^2/Q^2)) = \mathbf{C}^{(1)}(z). \quad (2.21)$$

Note that with this identification Eq. (2.17) and Eq. (2.18) (truncated to NLO) are not the same in practice, they are only perturbative equivalent. In any actual calculation with $\mu_R \neq Q$ also $\alpha_s(\mu_R^2)$ and $\alpha_s(Q^2)$ are different: they differ by an amount which is beyond our fixed-order framework and if that difference is “small” depends on the actual values of the strong coupling of the respective scales. However, with that we achieve the main purpose of this game: we generate some higher order terms, which we can then use to estimate MHOUs.

Starting from NNLO we see that the matching becomes non-trivial. In the case at hand we can insert Eq. (2.8) into Eq. (2.17) (with the choice $(\mu_R, \mu_{R,0}) \rightarrow (Q, \mu_R)$) and re-expand in powers of $\alpha_s(\mu_R^2)$ as requested. We find

$$\mathbf{C}^{(2)}(z, \ln(\mu_R^2/Q^2)) = \mathbf{C}^{(2)}(z) + \beta_0 \ln(\mu_R^2/Q^2) \mathbf{C}^{(1)}(z), \quad (2.22)$$

so the scale-varied coefficient receives a correction from a lower-order coefficient function.

Let's take a step back and review the general features.

Conclusion 2.4: Renormalization scale variation

Scale-varied coefficient functions

- can always be computed a posteriori from their non-scale-varied counterparts
- are a linear combination of their lower order non-scale-varied counterparts
- are derived by re-expanding the strong coupling in the perturbative sum up to the required perturbative accuracy

We also know that the dependency on the scale variation $\ln(\mu_R^2/Q^2)$ must be a polynomial dependence since the running of strong coupling is a pure resummation of this logarithm. By re-expanding we mean a Taylor expansion of the strong coupling at the default scale, here $\alpha_s(Q)$, in terms of the value at the renormalization scale, $\alpha_s(\mu_R^2)$. For computing the strong coupling at any scale $\alpha_s(\mu_R^2)$ we must use required perturbative accuracy, but for deriving the scale-varied coefficient functions we only need one order less. In the special case of fully inclusive DIS, which has no dependence on the strong coupling at LO, we even need only two orders lower.

For the sake of correcting Eq. (1.3), we can improve it by writing

$$\sigma(x, Q^2, \mu_R^2) = \left(\mathbf{C}^T(Q^2, \mu_R^2) \otimes \mathbf{f}(Q^2) \right)(x). \quad (2.23)$$

2.4.2 Factorization scale

As in Section 2.3.2 the general algorithm repeats for the factorization scale μ_F , which is the intrinsic scale of PDFs, except everything is more complicated. However, it is not only more complicated because PDFs are a set of function, but also because the PDF explicitly appears in Eq. (1.1). This implies, that the factorization scale dependence is not only be captured by the coefficient function, in contrast to the renormalization scale. In particular, the factorization scale captures collinear divergencies, which arise in diagrams, and re-attributes them to the PDF, thus there is a direct connection between the two. Recall our main goal here: we want to estimate MHOU by generating higher

order terms and we simply have more freedom here to do so. In order to simplify the discussion we explicitly neglect renormalization scale variations for the moment.

The most common way to deal with factorization scale variation is to directly modify the coefficient functions, as in the case for Eq. (2.16), and we refer to this choice as “factorization scale variation scheme C” [40]. Specifically, we make a similar ansatz and write

$$\sigma(x, Q^2, \mu_F^2) = \left(\mathbf{C}^T(Q^2, \mu_F^2) \otimes \mathbf{f}(\mu_F^2) \right)(x) \quad (2.24)$$

where now μ_F has its intended place and the usual perturbative expansion of the coefficient function still holds, i.e.

$$\mathbf{C}(z, Q^2, \mu_F^2) = \sum_{j=0} (\alpha_s(Q^2))^j \mathbf{C}^{(j)}(z, \ln(\mu_F^2/Q^2)). \quad (2.25)$$

However, in order to derive an explicit expression for $\mathbf{C}(z, Q^2, \mu_F^2)$ it is instructive to start from “factorization scale variation scheme B” [40]. In this scheme, we make a different choice and attribute the variation instead to the PDF:

$$\sigma(x, Q^2, \mu_F^2) = \left(\mathbf{C}^T(Q^2) \otimes \mathbf{f}^{SV}(\mu_F^2, \ln(\mu_F^2/Q^2)) \right)(x) \quad (2.26)$$

Note that Eq. (2.26) only uses the central coefficient functions $\mathbf{C}^T(Q^2)$, but requires the use of a special PDF $\mathbf{f}^{SV}$. In a PDF fit, where full control over the PDF is required any way and partonic matrix elements (coefficient functions) come from a variety of places this scheme can be actually preferable.

In order to define $\mathbf{f}^{SV}$ we start by observing, that we can write Eq. (2.14) also as a convolution with a factorization scale operator $\mathbf{K}$

$$\mathbf{f}(\xi, \mu_F^2) = (\mathbf{K}(\mu_F^2 \leftarrow \mu_{F,0}^2) \otimes \mathbf{f}(\mu_{F,0}^2))(\xi) \quad (2.27)$$

where $\mathbf{K}$ admits a perturbative expansion

$$\mathbf{K}(y, \mu_F^2 \leftarrow \mu_{F,0}^2) = \sum_{j=0} (\alpha_s(\mu_{F,0}^2))^j \mathbf{K}^{(j)}(y, \ln(\mu_F^2/\mu_{F,0}^2)) \quad (2.28)$$

and is given up to NLO accuracy by

$$\mathbf{K}(y, \mu_F^2 \leftarrow \mu_{F,0}^2) = \mathbb{1} \delta(1-y) + \alpha_s(\mu_{F,0}^2) \ln(\mu_F^2/\mu_{F,0}^2) \mathbf{P}^{(0)}(y) + \dots \quad (2.29)$$

Here, $\mathbb{1}$ refers to an identity operation in flavour space and $\delta(1-y)$ is the identity in the function convolution space as $\mathbf{K}$ is acting on both.

Note that $\mathbf{K}$ is a fixed order equivalent of the EKO $\mathbf{E}$, but the two are fundamentally different: the latter is a true solution to DGLAP, which must resum all necessary terms, but the former is only an approximation for a small amount of evolution. Specifically, $\mathbf{K}$ only is meaningful when we consider scale variations and in particular we use the interplay between $\mathbf{K}$ and $\mathbf{E}$ to generate the missing higher order terms we are looking for, because we get

$$\mathbf{f}(\xi, \mu_1^2) = (\mathbf{K}(\mu_1^2 \leftarrow \mu_2^2) \otimes \mathbf{E}(\mu_2^2 \leftarrow \mu_1^2) \otimes \mathbf{f}(\mu_1^2))(\xi) + \dots \quad (2.30)$$

where $\dots$ denotes the missing terms. Inserting the equation back into Eq. (1.3) with $(\mu_1, \mu_2) \rightarrow (Q, \mu_F)$ we obtain the definition of our sought-after special PDF

$$\mathbf{f}^{SV}(\xi, \mu_F^2, \ln(\mu_F^2/Q^2)) = (\mathbf{K}(Q^2 \leftarrow \mu_F^2) \otimes \mathbf{f}(\mu_F^2))(\xi) = \mathbf{f}(\xi, Q^2) + \dots \quad (2.31)$$

where we apply Eq. (2.13) for the PDF.

We can now return to the definition of scheme C: we insert Eq. (2.31) into Eq. (2.26) and obtain

$$\sigma(x, Q^2, \mu_F^2) = \left[\mathbf{C}^T(Q^2) \otimes (\mathbf{K}(Q^2 \leftarrow \mu_F^2) \otimes \mathbf{f}^{SV}(\mu_F^2)) \right] (x) \quad (2.32)$$

$$= \left[(\mathbf{C}^T(Q^2) \otimes \mathbf{K}(Q^2 \leftarrow \mu_F^2)) \otimes \mathbf{f}^{SV}(\mu_F^2) \right] (x) \quad (2.33)$$

where we only used the associative property of the convolution. However, we can now re-expand terms in a different order. First, it is convenient to use the perturbative expansion of the strong coupling to rewrite $\mathbf{K}$ in powers of $\alpha_s(Q^2)$ (as opposed to $\alpha_s(\mu_F^2)$, Eq. (2.29)):

$$\mathbf{K}'(y, Q^2 \leftarrow \mu_F^2) = \sum_{j=0} (\alpha_s(Q^2))^j \mathbf{K}'^{(j)}(y, \ln(\mu_F^2/Q^2)) \quad (2.34)$$

with

$$\mathbf{K}'(y, Q^2 \leftarrow \mu_F^2) = (\mathbf{K}(Q^2 \leftarrow \mu_F^2) \otimes [\mathbb{1} + \mathcal{O}(\alpha_s(Q^2))]) (y) \quad (2.35)$$

i.e. $\mathbf{K}'$ and $\mathbf{K}$ differ by higher order terms. Up to NLO accuracy we find from Eq. (2.29)

$$\mathbf{K}'(y, Q^2 \leftarrow \mu_F^2) = \mathbb{1}\delta(1-y) - \alpha_s(Q^2) \ln(\mu_F^2/Q^2) \mathbf{P}^{(0)}(y) + \dots \quad (2.36)$$

Second, we use the definition of $\mathbf{C}(z, Q^2, \mu_F^2)$

$$\mathbf{C}^T(z, Q^2, \mu_F^2) = \left(\mathbf{C}^T(Q^2) \otimes \mathbf{K}'(Q^2 \leftarrow \mu_F^2) \right) (z) \quad (2.37)$$

which reorganizes the product of sums in Eq. (2.33) into a single sum, Eq. (2.25). Let's see this in practice.

As before, we start from LO accuracy and from Eq. (2.36) we find immediately

$$\mathbf{C}^{(0)}(z, \ln(\mu_F^2/Q^2)) = \mathbf{C}^{(0)}(z), \quad (2.38)$$

i.e., also here scale variations do not alter the LO coefficient function. However, unlike in the case for renormalization scale variations, factorization scale variations have always an impact starting from LO accuracy, because PDFs are present per-se in the factorization formula, Eq. (1.1). Specifically, in any actual calculation with $\mu_F \neq Q$ also $\mathbf{f}(\mu_F^2)$ and $\mathbf{f}(Q^2)$ are different: they differ by an amount which is beyond our fixed-order framework and if that difference is “small” depends on the actual values of the PDF of the respective scales. However, with that we achieve the main purpose of this game: we generate some higher order terms, which we can then use to estimate MHOUs. This is an intentional copy from Section 2.4.1.

Starting from NLO the matching becomes non-trivial and, e.g., at this order we find

$$\mathbf{C}^{(1),T}(z, \ln(\mu_F^2/Q^2)) = \mathbf{C}^{(1),T}(z) - \ln(\mu_F^2/Q^2) \left(\mathbf{C}^{(0),T} \otimes \mathbf{P}^{(0)} \right) (z). \quad (2.39)$$

Always remember that for PDF operations, such as acting with the matrix of splitting functions $\mathbf{P}(y, \mu_F^2)$, non-commutativity is crucial, i.e., the flavor index dictated by the left-hand-side of Eq. (2.39) is matched by the second term on the right-hand-side by a vector-matrix multiplication (from the left). Note that starting from NNLO also explicit coefficients of the beta function start to appear, since the logarithm $\ln(\mu_F^2/Q^2)$

is coming directly from the running of the strong coupling, Eq. (2.7). In other words: PDF evolution resums logarithms through the running of the strong coupling.

In the following we assume scheme C as our default choice, thus conclusion 2.4 holds also for factorization scale variations and we can improve Eq. (1.3) with Eq. (2.24). The case of scheme B⁴ is slightly more complicated, as explained above, but straightforward to obtain.

2.4.3 Combined variation

We recall, the renormalization and factorization scales are two independent scales, related to two different kind of physics. The former captures ultra-violet and the latter collinear divergencies. While we discuss them one at a time above, in practice we have to consider them simultaneously. In order to do so, we must first consider factorization scale variation, which yields Eq. (2.24),

$$\sigma(x, Q^2, \mu_F^2, \mu_R^2 = Q^2) = \left(\mathbf{C}^T(Q^2, \mu_F^2, \mu_R^2 = Q^2) \otimes \mathbf{f}(\mu_F^2) \right)(x) \quad (2.40)$$

but now we make the renormalization scale dependence explicit and $\mathbf{C}(Q^2, \mu_F^2, \mu_R^2 = Q^2)$ corresponds to Eq. (2.25). Then in the second step, we apply the renormalization scale variation onto $\mathbf{C}(Q^2, \mu_F^2, \mu_R^2 = Q^2)$, which still depend on $\alpha_s(Q^2)$, and get

$$\sigma(x, Q^2, \mu_F^2, \mu_R^2) = \left(\mathbf{C}^T(Q^2, \mu_F^2, \mu_R^2) \otimes \mathbf{f}(\mu_F^2) \right)(x). \quad (2.41)$$

We can make the dependence on the various scales in Eq. (2.41) more explicit by rewriting it as

$$\sigma(x, Q^2, \mu_F^2, \mu_R^2) = \left(\mathbf{C}^T(\alpha_s(\mu_R^2), \ln(\mu_F^2/Q^2), \ln(\mu_R^2/Q^2)) \otimes \mathbf{f}(\mu_F^2) \right)(x) \quad (2.42)$$

where, again, finally μ_F and μ_R fulfill their dedicated role. Note that in Eq. (2.42) two additional scales, $\mu_{F,0}$ and $\mu_{R,0}$ are left implicit and those eventually guarantee that it is a dimensionless equation.

⁴scheme A is doing yet something different - see Ref. [40]

Let's see this in practice and remember that in order to see renormalization scale logarithms we need to play at NNLO. We find

$$\begin{aligned}
& \mathbf{C}(z, \alpha_s(\mu_R^2), \ln(\mu_F^2/Q^2), \ln(\mu_R^2/Q^2)) \\
&= \mathbf{C}^{(0)}(z) \\
&+ \alpha_s(\mu_R^2) \left[\mathbf{C}^{(1)}(z) + \mathbf{C}^{(1),F}(z) \ln(\mu_F^2/Q^2) \right] \\
&+ (\alpha_s(\mu_R^2))^2 \left[\mathbf{C}^{(2)}(z) + \mathbf{C}^{(2),F}(z) \ln(\mu_F^2/Q^2) + \mathbf{C}^{(2),R}(z) \ln(\mu_R^2/Q^2) \right. \\
&\quad \left. + \mathbf{C}^{(2),FF}(z) \ln^2(\mu_F^2/Q^2) + \mathbf{C}^{(2),FR}(z) \ln(\mu_F^2/Q^2) \ln(\mu_R^2/Q^2) \right] + \dots \quad (2.43)
\end{aligned}$$

We see that the number of terms per perturbative order quickly grows as the respective algorithms collect more of the full behavior. The appearance of $\mathbf{C}^{(2),FR}$ also demonstrate how the two variations become intertwined.

2.4.4 Discussion

The renormalization and factorization scale are remnants of the necessary regularization of UV and collinear divergencies. Although, scale variations are a genuine pQCD property they have some features, which questions their usefulness for estimating MHOUs.

When introducing the scales, we are required to set them to “a typical scale of the process”, which makes them inherently arbitrary from the start before doing any variation. In fully inclusive DIS there is a single scale available, the virtuality Q^2 , and thus the common choice is $\mu_F^2 = Q^2 = \mu_R^2$, but any multiplicative there of could work just as well⁵. If we want to use scale variations to estimate MHOUs, we need to make a further arbitrary choice about which range we want to use for the respective scales and if we want to correlate them. As we see in conclusion 2.4 the scale-varied coefficient functions are always a linear combination of the existing ones, i.e. they can not introduce new channels (new PDF combinations). However, it is often straightforward to show that a new channel should appear although its size is unknown and scale variations are unable to predict this. An example for such a scenario in fully inclusive DIS can be seen at NLO: starting from NLO the gluon can participate in the scattering, but the LO only contains quark PDFs. In other cases, we know that higher orders must contain certain kinematic logs, e.g. from particles radiating at small and/or large transverse momentum or in the

⁵For example CT18X [52] makes a different choice.

soft limit and in fully inclusive DIS the latter is the case⁶. Also those corrections can (in many cases) not be predicted by scale variations.

Scale variations also provide no statistical interpretation, which is in stark contrast to experimental uncertainties, where we usually assume Gaussian uncertainties and thus the provided error has a clear interpretation. However, strictly speaking this only applies to statistical errors but not systematic errors and the question which of the dominates depends very much on the specific measurement.

Despite all these (potential) problems, scale variations have a solid foundation in QFT and can be applied (straightforwardly) to any pQCD process, which makes them an easy tool to use MHO. In order to make a faithful statement, we can then simply say: when we compute this process with this specific choice of scale, we obtain this number; if we vary that scale in this range, we obtain this range of numbers.

2.4.5 Implementation

Let us return now to more practical consequences. As we see above, renormalization scale variations are just a scalar operation, but factorization scale variations are more complicated and it is worth spelling out their explicit construction to see what is actually happening.

The factorization scale variation is directly related to the PDFs themselves and, in particular, eventually, they derive from Eq. (2.10). However, the splitting functions $\mathbf{P}$ are encoding how partons couple among each other, which is through the gluon⁷ of course. This is in contrast to DIS, where instead the coefficient functions are encoding how partons couple to the electro-weak vector boson, which is completely different, see conclusion 1.1.

As usual, at LO accuracy the problem does not appear yet, because there are no scale-varied coefficient functions, see Eq. (2.38), but starting from NLO accuracy we need to worry. In particular, we know that the gluon couples the same to all quarks, thus, if we assume four flavors (see Section 2.5) for the moment and combine them in

⁶More mathematically speaking: soft gluon resummation is a double logarithmic resummation $\tilde{c}_q^{(j)} \sim \alpha_s^j \ln^{2j}(N)$ for $N \rightarrow \infty$ in Mellin space, Eq. (2.11).

⁷for quark-photon interaction see Ref. [53]

the following way [53]⁸,

$$f_{\Sigma_\Delta}(x, Q^2) = f_u(x, Q^2) + f_{\bar{u}}(x, Q^2) + f_c(x, Q^2) + f_{\bar{c}}(x, Q^2) \\ - f_d(x, Q^2) - f_{\bar{d}}(x, Q^2) - f_s(x, Q^2) - f_{\bar{s}}(x, Q^2) \quad (2.44)$$

we know that this combination does not interact with the gluon. In order to see more clearly what happens we need to go back to the vector notation from Eq. (1.3). We repeat: we denote objects, which have a non-trivial dependence on the parton flavors, with bold font. This applies to vectors in flavor space, e.g. coefficient functions $C_j \rightarrow \mathbf{C}$ or PDFs $f_j \rightarrow \mathbf{f}$, but also to matrices, e.g. splitting functions $P_{jk} \rightarrow \mathbf{P}$ or the factorization scale operator $K_{jl} \rightarrow \mathbf{K}$. By introducing the vector notation we hide any reference to a given basis in the flavor space, which now becomes relevant (see also Section 2.5).

For computing LO and NLO DIS coefficient functions it is convenient to consider each quark individually and we can find easily Eqs. (1.4) and (1.20). We refer to this choice of basis as the flavor basis. However, for the discussion in Section 2.3.2 and in particular for dealing with splitting functions, $\mathbf{P}$, a different basis is more convenient: the evolution basis. The two choices are both orthogonal bases, i.e. they are simply related by a rotation⁹ in flavor space. Specifically, $\mathbf{P}$ has a much simpler representation if one of the basis vectors is the singlet

$$f_\Sigma(x, Q^2) = f_u(x, Q^2) + f_{\bar{u}}(x, Q^2) + f_c(x, Q^2) + f_{\bar{c}}(x, Q^2) \\ - f_d(x, Q^2) - f_{\bar{d}}(x, Q^2) - f_s(x, Q^2) - f_{\bar{s}}(x, Q^2) \quad (2.45)$$

and all other basis vectors can be chosen such that they do not couple to the gluon, e.g. Eq. (2.44).

Let's see this in practice for LO photon DIS: we start from Eq. (1.20), which is written in flavor basis, i.e. for each given quark flavor and the gluon. Inserting Eq. (1.20) into Eq. (1.1) yields Eq. (1.4) and we find explicitly

$$\sigma(x, Q^2) = e_d^2(f_d(x, Q^2) + f_{\bar{d}}(x, Q^2) + f_s(x, Q^2) + f_{\bar{s}}(x, Q^2)) \\ + e_u^2(f_u(x, Q^2) + f_{\bar{u}}(x, Q^2) + f_c(x, Q^2) + f_{\bar{c}}(x, Q^2)) \quad (2.46)$$

⁸for the more general case see Ref. [53]

⁹actually, it is a rotation and a scaling, i.e. the evolution basis is not orthonormal, which may cause some trouble

Using Eqs. (2.44) and (2.45) we can rewrite this as

$$\sigma(x, Q^2) = (e_d^2 + e_u^2)f_\Sigma(x, Q^2) + (e_u^2 - e_d^2)f_{\Sigma_\Delta}(x, Q^2) \quad (2.47)$$

or equivalently

$$C_\Sigma^{(0)}(z) = (e_d^2 + e_u^2)\delta(1-z), \quad C_{\Sigma_\Delta}^{(0)}(z) = (e_u^2 - e_d^2)\delta(1-z), \quad C_j(z) = 0 \forall j \neq \{\Sigma, \Sigma_\Delta\} \quad (2.48)$$

We stress that Eqs. (1.20) and (2.48) are two equivalent representation of the coefficient function $\mathbf{C}^{(0)}$ using two different bases.

Moving to NLO, we have the same structure in the quark sector, Eq. (1.21), and we encounter for the first time the gluon channel, Eq. (1.22). Thus, when we want to execute Eq. (2.39), we need to consider three directions in flavor space: the gluon, the singlet f_Σ and Eq. (2.44). The preferred basis for $\mathbf{P}$ is manifest by the following statements,

$$P_{\Sigma\Sigma_\Delta}(y) = P_{g\Sigma_\Delta}(y) = P_{\Sigma_\Delta\Sigma}(y) = P_{\Sigma_\Delta g}(y) = 0 \quad (2.49)$$

which would not occur in flavor basis. Eventually, Eq. (2.39) becomes explicitly

$$C_{\Sigma_\Delta}^{(1)}(z, \ln(\mu_F^2/Q^2)) = C_{\Sigma_\Delta}^{(1)}(z) - \ln(\mu_F^2/Q^2) \left(C_{\Sigma_\Delta}^{(0)} \otimes P_{\Sigma_\Delta\Sigma_\Delta}^{(0)} \right)(z) \quad (2.50)$$

$$C_\Sigma^{(1)}(z, \ln(\mu_F^2/Q^2)) = C_\Sigma^{(1)}(z) - \ln(\mu_F^2/Q^2) \left(C_\Sigma^{(0)} \otimes P_{\Sigma\Sigma}^{(0)} \right)(z) \quad (2.51)$$

$$C_g^{(1)}(z, \ln(\mu_F^2/Q^2)) = C_g^{(1)}(z) - \ln(\mu_F^2/Q^2) \left(C_\Sigma^{(0)} \otimes P_{\Sigma g}^{(0)} \right)(z) \quad (2.52)$$

where the required splitting functions can be found in the literature [8–10].

Actually, also LO splitting functions $\mathbf{P}^{(0)}$ obey symmetry relations, in particular all quarks couple identically to the gluon and it is not possible (yet) to change the quark flavor. This implies $P_{\Sigma_\Delta\Sigma_\Delta}^{(0)}(y) = P_{\Sigma\Sigma}^{(0)}(y) = P_{qq}^{(0)}(y)$ and thus Eq. (2.39) can also be executed straightforwardly in flavor space. However, this symmetry breaks down starting from NLO splitting functions $\mathbf{P}^{(1)}$, which would be required for NNLO DIS cross sections.

2.4.6 Interplay with previous content

- Renormalization scale variation is a reshuffling of lower order coefficient functions and factorization scale variation adds convolutions between splitting functions and

lower order coefficient functions. This implies that the mathematical complexity remains the same or in other words: we never encounter new mathematical objects in scale variations; see Section 1.4.

- Although scale-varied coefficient functions can always be computed a posteriori, see conclusion 2.4, in practice it usually makes sense to tabulate them into interpolation grids as well, see Section 2.2. These additional coefficient functions can be easily accommodated when we generalize our concept of “order”: it can not only represent the perturbative order per se, but also additional powers of renormalization or factorization logarithms: $\alpha_s^k(\mu_R^2) \ln^m(\mu_R^2/Q^2) \ln^n(\mu_R^2/Q^2)$.
- The necessary projection onto a singlet component is particularly tricky for CC DIS, see Section 2.1. Recall that the $W^\pm$ boson only couple to specific PDFs and thus if we want to project the coefficient functions into evolution basis we need the full basis (and not just Eqs. (2.44) and (2.45)).
- Note that the coupling structure is always retained, i.e.

$$C_g^{(1)}(z) \sim e_u^2 + e_d^2 \sim C_g^{(1)}(z, \ln(\mu_F^2/Q^2)) \quad (2.53)$$

This implies that a given “channel” in an interpolation grid (Section 2.2) has a unique coupling, though recall that some channels only appear at high enough perturbative order.

2.5 Heavy quarks

When we compute LO DIS from Fig. 1.2 we say it is easy to identify the respective coefficient function, Eqs. (1.16) and (1.20). However, what exactly do we mean with the “ $\forall q$ ”? Which quarks are we exactly talking about? This is directly linked to the flavor space to which we hint when using bold font. However, unlike the problem we discuss in Section 2.4.5 about different bases (flavor basis vs. evolution basis), here we are more directly concerned what the flavor space is in the first place.

2.5.1 Flavor space

We define the flavor space $\mathcal{F}$ as the vector space spanned by all quark flavors and the gluon. We consider the naive flavor basis as our canonical basis and write

$$\mathcal{F} = \mathcal{F}_{fl} = \text{span}(g, u, \bar{u}, d, \bar{d}, s, \bar{s}, c, \bar{c}, b, \bar{b}, t, \bar{t}). \quad (2.54)$$

We have a 13 dimensional vector space, $\dim \mathcal{F} = 13$ ¹⁰, consisting of the gluon, six quarks, and six anti-quarks. Our bold vectors are an element in that space, e.g. $\mathbf{C} \in \mathcal{F}$, or more plainly,

$$\mathbf{C}^T \sim (C_g, C_u, C_{\bar{u}}, C_d, C_{\bar{d}}, C_s, C_{\bar{s}}, C_c, C_{\bar{c}}, C_b, C_{\bar{b}}, C_t, C_{\bar{t}}) \quad (2.55)$$

and, as said in Section 2.4.5, any other basis works just as well. The same applies also to the PDF $\mathbf{f}$, which is the other vector in Eq. (1.3), and any operator, e.g. splitting functions $\mathbf{P}$, just lives in the product space, $\mathbf{P} \in \mathcal{F} \otimes \mathcal{F}$.

The question we need to address here is: which quarks exactly can participate in Eqs. (1.4) and (1.20)? We refer to this number as the number of active quarks n_f . This dependency is completely hidden in Eq. (1.1), but, as we demonstrate in the following, is a major concern for both coefficient functions and PDF. Note that the gluon is always considered active and contributes to all diagrams always.

The number of flavors n_f is arbitrary per se and there is no absolutely correct answer, since there is no theory argument, which would fix this choice. However, in practice for a given observable $\sigma(x, Q^2)$ we can consider, which options are phenomenologically reasonable and if some option would yield unstable results.

For the sake of the argument, let's consider a fully inclusive DIS cross section $\sigma(x, Q^2 = 25 \text{ GeV}^2)$, so we probe the proton at $\mu_F = Q = 5 \text{ GeV}$. We can now consider the different quark masses as a guideline: on the one side the masses of up, down, and strange are significantly below the non-perturbative regime of QCD $m_u, m_d, m_s \ll \Lambda_{\text{QCD}} \sim 1 \text{ GeV}$ and we can safely assume we can find them in the proton. On the other side the mass of the top is significantly above the probed scale, $m_t \gg Q$, thus it is very unlikely to find it in the proton and we can neglect it. We can shut down such contributions by setting $f_t(\xi, Q^2) = 0 = f_{\bar{t}}(\xi, Q^2)$ and this way the top quark does

¹⁰one can add the photon and make it 14 dimensional [53]

not contribute to Eq. (1.1). In addition, we can also define $C_t(z, Q^2) = 0 = C_{\bar{t}}(z, Q^2)$ to explicitly say that we don't let the top contribute.

Now, we consider the two remaining quarks, charm and bottom, for which maybe there is no clear answer. First, for the charm quark we have $m_c \sim 1.5 \text{ GeV}$ and we can assume reasonably that it can contribute to DIS. On the other side, for the bottom quark we have $m_b \sim 4.5 \text{ GeV}$ and maybe we can both allow or deny it. The specific choice on which quarks we allow to participate and in which way they do so is referred to as flavor number scheme.

2.5.2 ZM-VFNS

One very popular choice is the Zero-Mass Variable Flavor Number Scheme (ZM-VFNS), which is a telling name and which we have silently adopted in the previous parts. Formally, we can define it in the following way: first, whenever the probed scale Q crosses a quark mass that quark becomes active; second, in coefficient functions all active quarks are considered massless and all non-active quarks are considered infinite massive. The infinite mass ensures $f_H(\xi, Q^2) = 0 = f_{\bar{H}}(\xi, Q^2)$ and $C_H(z, Q^2) = 0 = C_{\bar{H}}(z, Q^2)$ for all non-active quarks H and the zero mass ensures that all formulas in Section 1.4 hold. If we denote the new dependency on the number of active flavors n_f by a superscript (n_f) we can define the ZM-VFNS in the following way

$$\sigma^{\text{ZM-VFNS}}(x, Q^2) = \begin{cases} \sigma^{(3)}(x, Q^2) & \text{if } Q^2 < m_c^2 \\ \sigma^{(4)}(x, Q^2) & \text{if } m_c^2 \leq Q^2 < m_b^2 \\ \sigma^{(5)}(x, Q^2) & \text{if } m_b^2 \leq Q^2 < m_t^2 \\ \sigma^{(6)}(x, Q^2) & \text{if } Q^2 \geq m_t^2 \end{cases} \quad (2.56)$$

where

$$\sigma^{(n_f)}(x, Q^2) = \left(\mathbf{C}^{T, (n_f)}(Q^2) \otimes \mathbf{f}^{(n_f)}(Q^2) \right) (x) \quad (2.57)$$

i.e. each contribution has its own factorization formula and we have, e.g.,

$$\begin{aligned} C_H^{(3)}(z, Q^2) &= 0 = C_{\bar{H}}^{(3)}(z, Q^2) \\ f_H^{(3)}(\xi, Q^2) &= 0 = f_{\bar{H}}^{(3)}(\xi, Q^2) \end{aligned} \quad H \in \{c, \bar{c}, b, \bar{b}, t, \bar{t}\} \quad (2.58)$$

and similar for the others. Specifically, we can improve Eq. (1.20) by writing

$$C_q^{(0),(3)}(z) = e_q^2 \delta(1 - z) \quad q \in \{c, \bar{c}, b, \bar{b}, t, \bar{t}\} \quad (2.59)$$

$$C_H^{(0),(3)}(z) = 0 \quad H \in \{c, \bar{c}, b, \bar{b}, t, \bar{t}\} \quad (2.60)$$

$$C_g^{(0),(3)}(z) = 0 \quad (2.61)$$

and we finally give full details on the available flavor space $\mathcal{F}$.

Note that it does not make sense to consider less than three quarks, since $m_u, m_d, m_s \ll \Lambda_{\text{QCD}} \sim 1 \text{ GeV}$ and pQCD only works for $Q > \Lambda_{\text{QCD}}$. All quarks are considered with their actual mass for the decision if they are in or out, Eq. (2.56), but then after that their mass is discarded when computing $\mathbf{C}^{(n_f)}(z, Q^2)$. Thus, the VFNS indicates that the number of active quarks n_f depends on the probed scale Q , i.e. it is variable, and the ZM explains how to compute diagrams. However, once we have an actual scale in hand, n_f is also fixed.

Conclusion 2.5: ZM-VFNS

In the ZM-VFNS

- the number of active quarks n_f depends on the probed scale Q^2
- contain the diagrams to compute coefficient functions only massless quarks

2.5.3 FFNS

For the number of active quarks we can make another simple choice: fix n_f once and forever - the Fixed Flavor Number Scheme (FFNS). Here, the probed scale Q does not matter, but we use the same coefficient functions for any scale. Actually, saying FFNS is not sufficient, we still have some choices and the most obvious is n_f .

However, we can also improve with respect to other parts: while it is significantly easier to compute diagrams with only massless quarks, we can compute them also with massive quarks. For simplicity we assume in the following we have exactly one¹¹ massive quark with mass m^2 . Moreover, while we keep the notation general you can always consider as a concrete example $n_f = 3$ and thus $m^2 = m_c^2$. We recall two important

¹¹the case of two massive quarks is e.g. discussed in Ref. [54]

features of the propagator of a massive quark,

$$D_q(p) \sim \frac{p_\mu \gamma^\mu + m \mathbb{1}}{p^2 - m^2}; \quad (2.62)$$

first, the mass appears explicitly in the numerator, which makes computing Dirac traces significantly more lengthy, and, second, the mass appears in the denominator, which shields the propagator from a collinear pole. When we consider a finite mass we eventually have three different scales in our diagrams: the total energy of the colliding partons $\hat{s}$, the photon virtuality Q^2 , and the quark mass m^2 . This means that unlike the massless case discussed so far, the coefficient functions now depend on two scales, for example the partonic momentum fraction $z = Q^2/(\hat{s} + Q^2)$ and $\xi_q = Q^2/m^2$ or any other bijective combination¹². This in turn implies that the available mathematical function space is much bigger, which makes the computation of these coefficient functions $\mathbf{C}(z, \xi_q, Q^2)$ significantly more complex.

However, we focus here on the new dependency on ξ_q , which captures the new features. First, we consider the case $Q^2 \gg m^2$ or $\xi_q \rightarrow \infty$ where the denominator of Eq. (2.62) plays the important role. Indeed, since it is shielding the propagator from a collinear divergency, we encounter some signals thereof when we put $m^2 \rightarrow 0$. Specifically, we can rewrite the coefficient functions in this limit,

$$\lim_{Q^2/m^2 \rightarrow \infty} \mathbf{C}^{(n_f),T}(Q^2, m^2) = \mathbf{C}^{(n_f+1),T}(Q^2) \mathbf{A}^{(n_f+1)}(Q^2, m^2) \quad (2.63)$$

i.e. we can connect the coefficient functions using two neighboring schemes and a new, universal operator $\mathbf{A}$, which admits a perturbative series

$$\mathbf{A}^{(n_f+1)}(Q^2, m^2) = \sum_{j=0} (\alpha_s(Q^2))^j \mathbf{A}^{(j), (n_f+1)}(\ln(Q^2/m^2)). \quad (2.64)$$

We discuss more details of this new operator in a moment, but the important thing to stress here is the universality. This universality derives from the universality of PDFs, which makes the operator closely connected to collinear splitting functions as we expect. Moreover, Eq. (2.63) puts strong constraints on the computation of any coefficient function¹³ once the universal operator $\mathbf{A}$ is known at the requested accuracy.

¹²For example, for NC DIS one can show [55] that $\chi = \frac{1 - \sqrt{1 - 4m^2/s}}{1 + \sqrt{1 - 4m^2/s}}$ and $\chi_q = \frac{\sqrt{1 + 4m^2/Q^2} - 1}{\sqrt{1 + 4m^2/Q^2} + 1}$ are good combinations.

¹³or any partonic matrix elements as similar equations hold, e.g., in hadron-hadron collisions

Note that the coefficient functions in Eq. (2.63) depend on different sets of variables, since we consider the “+1” quark massless in the $(n_f + 1)$ scheme. Thus, as expected, the full mass dependence on the right-hand-side of Eq. (2.63) is carried by the universal operator $\mathbf{A}$ and it is a pure, actual polynomial, dependency on $\ln(Q^2/m^2)$. So, the remnant of the collinear divergency for a massive quark is a logarithm $\ln(Q^2/m^2)$ as we may have guessed from the beginning.

Next, we consider the case $Q^2 \sim m^2$ where the nominator of Eq. (2.62) plays the important role. While we conclude above that the FFNS coefficient functions contain logarithms $\ln(Q^2/m^2)$ they also naturally contain terms which are relatively suppressed to them. Specifically, if we expand the FFNS coefficient function around $Q^2 \sim m^2$ we obtain

$$\mathbf{C}^{(n_f)}(Q^2, m^2) = \mathbf{C}_{thr,0}^{(n_f)} + \mathbf{C}_{thr,1}^{(n_f)} \left(\frac{Q^2}{m^2} \right) + \mathbf{C}_{thr,2}^{(n_f)} \left(\frac{Q^2}{m^2} \right)^2 + \dots \quad (2.65)$$

with some coefficients $\mathbf{C}_{thr,k}^{(n_f)}$. Note that the equation is to be understood as a Taylor expansion at the requested point as in practice the actual dependency on ξ_q is very intricate¹⁴. However, it is clear that in the region $Q^2 \sim m^2$ these terms may not be small and thus neglecting such contributions may yield wrong results. That is the precisely the advantage of FFNS over ZM-VFNS: while in the latter we explicitly deny any massive quark in the diagrams, the former allows them to be included in the sum over diagrams.

Conclusion 2.6: FFNS

In the FFNS

- massive quarks are allowed in the computation of diagrams for coefficient functions
- we counter universal logarithms $\ln(Q^2/m^2)$ for $Q^2 \rightarrow m^2$ due to collinear divergencies

But if the FFNS calculation is actually better than ZM-VFNS, why don't we always use the former? Before answering the question, we need to return to the new operator $\mathbf{A}$.

¹⁴for example via χ_q

2.5.4 Matching schemes

The number of active quarks is not only relevant for coefficient functions, but, in fact, for the whole pQCD program and in particular for the strong coupling α_s and PDFs $\mathbf{f}$. As we see from Eq. (2.57) also PDFs depend on the number of flavors, e.g. $\mathbf{f}^{(3)}(\xi, \mu_F^2)$, and we can improve Eq. (1.15) with

$$\mathbf{C}^{(n_f)}(z, Q^2, m^2) = \sum_{k=0} \left(\alpha_s^{(n_f)}(Q^2) \right)^k \mathbf{C}^{(k), (n_f)}(z, \xi_q) \quad (2.66)$$

where we now also explicitly denote the flavor dependence of the strong coupling. Actually, from Eq. (2.5) we already known that the beta coefficients depend explicitly on n_f and thus α_s must depend on it as well. The same applies to PDFs since also splitting functions carry an explicit n_f dependence and all appearances above, indeed, correspond to $\mathbf{P}^{(n_f)}$.

The question is: how are those different schemes, e.g. $\alpha_s^{(3)}$ and $\alpha_s^{(4)}$, connected with each other? For coefficient functions the difference originates from the different diagrams which are included into the calculation of the coefficient functions, $\mathbf{C}^{(3)}$ and $\mathbf{C}^{(4)}$, but what about the strong coupling α_s and the PDFs $\mathbf{f}$? In order to answer this question it is important to return to the basic principles: we know that only cross sections σ are physical objects for which we can derive physical properties, e.g. that they must be positive. In contrast, the strong coupling α_s , PDFs $\mathbf{f}$, and coefficient functions $\mathbf{C}$ are not physical objects, which can not be measured directly in an experiment and which only are defined within our framework of pQCD. The concept of having different number of active flavors n_f is a theoretical idea, which we employ to make a consistent approximation in our calculation, but is irrelevant to nature. In pQCD we can still derive certain rules for our relevant objects in order to obtain a consistent theory, which we can then use to make predictions.

The important physical properties, which we need here is consistency and perturbative equivalence. Specifically, we require that the cross section in two different scheme are perturbatively equivalent, i.e.

$$\sigma^{(n_f)}(x, Q^2) = \sigma^{(n_f+1)}(x, Q^2) \left[1 + \mathcal{O}\left(\alpha_s^{(n_f+1)}(Q^2)\right) \right], \quad (2.67)$$

where the two cross section are consistently computed in their respective flavor number scheme. In particular, they use their own coefficient functions, their own strong coupling, and their own PDFs. However, since the coefficient functions, the beta coefficients, and

the splitting functions are computed from the respective relevant diagrams this makes the condition quite restrictive. Moreover, if we recall Section 2.3 we know that both, α_s and $\mathbf{f}$, perform a resummation, so Eq. (2.67) is to be understood in a logarithmic sense and thus, in practice, it is convenient to consider Eq. (2.67) in the limit $Q^2 \rightarrow \infty$.

One can show that the connection between two different schemes can be done by a universal matching procedure for the strong coupling and the PDFs. Specifically, for the strong coupling we find

$$\alpha_s^{(n_f+1)}(\mu_R^2) = \zeta^{(n_f+1)}(\mu_R^2, m^2) \alpha_s^{(n_f)}(\mu_R^2) \quad (2.68)$$

with a universal matching operator ζ , which admits a perturbative series and has a polynomial dependence on $\ln(\mu_R^2/m^2)$. Analogously, for the PDFs we find

$$\mathbf{f}^{(n_f+1)}(\xi, \mu_F^2) = \left(\mathbf{A}^{(n_f+1)}(\mu_F^2, m^2) \otimes \mathbf{f}^{(n_f)}(\mu_F^2) \right) (\xi) \quad (2.69)$$

with a universal matching operator $\mathbf{A}$, which is also used in Eq. (2.63). The scheme change, Eq. (2.68) and Eq. (2.69), can be done at any scale, but, following Eq. (2.56), is conventionally done at the mass of the heavy quark, $\mu_R^2 = m^2 = \mu_F^2$.

Let's see this in practice: consider a cross section $\sigma(x, Q^2)$ with $Q^2 > m^2$. As we say above the number of active quarks is a free choice, we can compute the cross section either in the n_f scheme, Eq. (2.57), or in the $(n_f + 1)$ scheme,

$$\sigma^{(n_f+1)}(x, Q^2) = \left(\mathbf{C}^{T, (n_f+1)}(Q^2) \otimes \mathbf{f}^{(n_f+1)}(Q^2) \right) (x). \quad (2.70)$$

Inserting Eq. (2.69) into Eq. (2.70) we obtain

$$\sigma^{(n_f+1)}(x, Q^2) = \left(\mathbf{C}^{T, (n_f+1)}(Q^2) \otimes \mathbf{A}^{(n_f+1)}(Q^2, m^2) \otimes \mathbf{f}^{(n_f)}(Q^2) \right) (x) \quad (2.71)$$

and if Eq. (2.63) holds, the two expressions, Eq. (2.57) and Eq. (2.70), are actually perturbatively equivalent. Recall that Eq. (2.63) only holds in the $Q^2 \gg m^2$ limit and so in practice for finite Q^2 the two expression are only perturbatively equivalent.

2.5.5 FONLL as an example of a GM-VFNS

2.5.6 Heavy quark structure functions

Chapter 3

More advanced topics

3.1 TMC

3.2 γ_5

- helicity
- HVBM + Larin

3.3 Positivity

- DIS vs. MSbar

3.4 Higher orders

- IHOU
- N3LO
- channels
- loops and legs

3.5 QED

- Qiu et al.

3.6 Valo

- $C_\gamma \rightarrow k$

Bibliography

- [1] A. Candido, F. Hekhorn, G. Magni, T. R. Rabemananjara, and R. Stegeman, Eur. Phys. J. C **84**, 698 (2024), 2401.15187.
- [2] NNPDF, R. D. Ball *et al.*, Eur. Phys. J. C **82**, 428 (2022), 2109.02653.
- [3] J. Gao, L. Harland-Lang, and J. Rojo, Phys. Rept. **742**, 1 (2018), 1709.04922.
- [4] G. Heinrich, Phys. Rept. **922**, 1 (2021), 2009.00516.
- [5] S. Amoroso *et al.*, Acta Phys. Polon. B **53**, 12 (2022), 2203.13923.
- [6] NNPDF, R. D. Ball *et al.*, Nature **608**, 483 (2022), 2208.08372.
- [7] NNPDF, R. D. Ball *et al.*, Phys. Rev. D **109**, L091501 (2024), 2311.00743.
- [8] E. Leader and E. Predazzi, *An Introduction to gauge theories and modern particle physics. Vol. 1: Electroweak interactions, the new particles and the parton model* (Cambridge University Press, 2011).
- [9] E. Leader and E. Predazzi, *An Introduction to gauge theories and modern particle physics. Vol. 2: CP violation, QCD and hard processes* (Cambridge University Press, 1996).
- [10] M. E. Peskin and D. V. Schroeder, *An Introduction to quantum field theory* (Addison-Wesley, Reading, USA, 1995).
- [11] F. Halzen and A. D. Martin, *QUARKS AND LEPTONS: AN INTRODUCTORY COURSE IN MODERN PARTICLE PHYSICS* (Wiley, 1984).
- [12] A. S. Kronfeld and C. Quigg, Am. J. Phys. **78**, 1081 (2010), 1002.5032.
- [13] A. Accardi *et al.*, Eur. Phys. J. A **52**, 268 (2016), 1212.1701.
- [14] A. Barontini *et al.*, Nnpdf/yadism: v0.13.10: Update n3lo coeff fnc, 2026.

- [15] M. Bertilsson, T. Lappi, H. Mäntysaari, and X.-B. Tong, (2026), 2601.07302.
- [16] C. Casuga and H. Mäntysaari, (2026), 2604.22332.
- [17] H1, V. Andreev *et al.*, Eur. Phys. J. C **75**, 65 (2015), 1406.4709.
- [18] C. Bierlich *et al.*, SciPost Phys. Codeb. **2022**, 8 (2022), 2203.11601.
- [19] I. Helenius, J. O. Laulainen, and C. T. Preuss, (2026), 2605.00502.
- [20] Wikipedia, File:Standard Model of Elementary Particles.svg, online, 2026.
- [21] J. C. Collins, D. E. Soper, and G. F. Sterman, Adv. Ser. Direct. High Energy Phys. **5**, 1 (1989), hep-ph/0409313.
- [22] Wikipedia, HERA-Tunnel.jpg, online, 2026.
- [23] H1, ZEUS, H. Abramowicz *et al.*, Eur. Phys. J. C **75**, 580 (2015), 1506.06042.
- [24] R. D. Ball, A. Chiefa, and R. Stegeman, Eur. Phys. J. C **86**, 281 (2026), 2511.14387.
- [25] L. A. Harland-Lang, T. Cridge, M. Reader, and R. S. Thorne, Eur. Phys. J. C **86**, 115 (2026), 2510.03753.
- [26] M. Cerutti *et al.*, Phys. Rev. D **111**, 094013 (2025), 2501.06849.
- [27] T. Lappi, H. Mäntysaari, H. Paukkunen, and M. Tevio, Eur. Phys. J. C **85**, 429 (2025), 2412.09589.
- [28] E. Spezzano, T. Jezo, M. Klasen, P. Risse, and I. Schienbein, JHEP **05**, 012 (2026), 2512.11569.
- [29] D. Akturk *et al.*, PTEP **2026**, 033C04 (2026), 2506.15445.
- [30] K. J. Eskola, P. Paakkinen, H. Paukkunen, and C. A. Salgado, Eur. Phys. J. C **82**, 413 (2022), 2112.12462.
- [31] R. Abdul Khalek *et al.*, Eur. Phys. J. C **82**, 507 (2022), 2201.12363.
- [32] M. Klasen, PoS DIS2025, 036 (2025), 2510.05880.
- [33] I. Helenius, M. Walt, and W. Vogelsang, Phys. Rev. D **105**, 094031 (2022), 2112.11904.
- [34] NNLOJET, A. Huss *et al.*, SciPost Phys. Codeb. **69**, 1 (2026), 2503.22804.

-
- [35] S. Carrazza, E. R. Nocera, C. Schwan, and M. Zaro, JHEP **12**, 108 (2020), 2008.12789.
 - [36] C. Schwan *et al.*, Nnpdf/pineappl: v1.0.0, 2025.
 - [37] T. Carli *et al.*, Eur. Phys. J. C **66**, 503 (2010), 0911.2985.
 - [38] fastNLO, M. Wobisch, D. Britzger, T. Kluge, K. Rabbertz, and F. Stober, (2011), 1109.1310.
 - [39] A. Barontini, A. Candido, J. M. Cruz-Martinez, F. Hekhorn, and C. Schwan, Comput. Phys. Commun. **297**, 109061 (2024), 2302.12124.
 - [40] NNPDF, R. D. Ball *et al.*, Eur. Phys. J. C **84**, 517 (2024), 2401.10319.
 - [41] G. Altarelli and G. Parisi, Nucl. Phys. B **126**, 298 (1977).
 - [42] V. N. Gribov and L. N. Lipatov, Sov. J. Nucl. Phys. **15**, 438 (1972).
 - [43] Y. L. Dokshitzer, Sov. Phys. JETP **46**, 641 (1977).
 - [44] A. Candido, F. Hekhorn, and G. Magni, Eur. Phys. J. C **82**, 976 (2022), 2202.02338.
 - [45] M. A. Lim and R. Poncelet, Phys. Rev. D **112**, L111901 (2025), 2412.14910.
 - [46] F. J. Tackmann, JHEP **08**, 098 (2025), 2411.18606.
 - [47] Z. Kassabov, M. Ubiali, and C. Voisey, JHEP **03**, 148 (2023), 2207.07616.
 - [48] M. Bonvini, Eur. Phys. J. C **80**, 989 (2020), 2006.16293.
 - [49] M. Cacciari and N. Houdeau, JHEP **09**, 039 (2011), 1105.5152.
 - [50] C. Duhr, A. Huss, A. Mazeliauskas, and R. Szafron, JHEP **09**, 122 (2021), 2106.04585.
 - [51] A. Ghosh *et al.*, SciPost Phys. Core **6**, 045 (2023), 2210.15167.
 - [52] T.-J. Hou *et al.*, Phys. Rev. D **103**, 014013 (2021), 1912.10053.
 - [53] NNPDF, R. D. Ball *et al.*, Eur. Phys. J. C **84**, 540 (2024), 2401.08749.
 - [54] A. Barontini, A. Candido, F. Hekhorn, G. Magni, and R. Stegeman, JHEP **10**, 004 (2024), 2408.07383.
 - [55] F. Hekhorn, *Next-to-Leading Order QCD Corrections to Heavy-Flavour Production in Neutral Current DIS*, PhD thesis, Tubingen U., Math. Inst., 2019, 1910.01536.