

# CS70 - Combinatorial Arguments

Name: Felix Su    SID: 25794773

Spring 2016    GSI: Gerald Zhang

## Strategy

1. Define equivalent quantity  $Q$
2. Count  $Q$  in a way to determine  $LHS = Q$
3. Count  $Q$  in a way to determine  $RHS = Q$
4. Conclude  $LHS = RHS$

## Examples

$$2^n = \sum_{k=0}^n \binom{n}{k} \quad (1)$$

**LHS:** Using counting product rule, this is the number subsets of  $n$  decided by either choosing or not choosing each item. There are a total of  $2^n$  possible subsets of size  $k \in (0, n)$  from a total of  $n$  items.

**RHS:** The summation of all possible ways to choose  $k \in (0, n)$  items from  $n$ , which is the same as left.

$$\sum_{m=k}^n \binom{m}{k} = \binom{n+1}{k+1} \quad (2)$$

**LHS:**

**RHS:**

$$\sum_{k=0}^n \binom{m+k}{k} = \binom{n+m+1}{n} \quad (3)$$

**LHS:** Specify the smallest element from  $\{1, \dots, n+m+1\}$  not in the selected subset. If it is 1, you must choose the remaining  $n$  items from  $\{2, \dots, n+m+1\}$ ,  $\binom{m+n}{n}$ , which is the  $k=n$  term. If it is 2, 1 is included in the subset, so you only need to choose from  $n-1$  remaining items from  $\{2, \dots, n+m+1\}$ ,  $\binom{m+n-1}{n-1}$ . This continues down to having the smallest number not in the set be  $n+1$ , which would cause,  $\binom{m+0}{0}$ . This equates to all possible ways to select select  $n$  items from  $n+m+1$  choices

**RHS:** All possible ways to select select  $n$  items from  $n+m+1$  choices.

$$\sum_{k=0}^n \binom{n}{k}^2 = \binom{2n}{n} \quad (4)$$

**LHS:** Split  $\{\pm 1, \dots, \pm n\}$  into 2 sets, 1 of positive and 1 of negative elements. Choose  $k$  positive items and  $n-k$  negative items (which is still equivalent to  $\binom{n}{k}$ ). This is equivalent to choosing  $n$  items out of  $\{\pm 1, \dots, \pm n\}$ .

**RHS:** All possible ways to directly choose  $n$  items out of  $\{\pm 1, \dots, \pm n\}$ .

$$\sum_{k=0}^r \binom{m}{k} \binom{n}{r-k} = \binom{n+m}{r} \quad (5)$$

**LHS:** Split the collection of books into a set of textbooks and one of comic books. Select  $k$  books from the set of textbooks and  $r - k$  books from the set of  $m$  comic books. This is the same as selecting  $r$  total books from a collection of  $n$  textbooks and  $m$  comic books.

**RHS:** Choose  $r$  books from a collection of  $n$  textbooks and  $m$  comic books.

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k} \quad (6)$$

**LHS:** All possible ways to select  $k$  items from a collection of  $n$  choices.

**RHS:** Set aside one element. Assuming this item must be in the selected subset of  $k$  items, select the remaining  $k - 1$  items from the remaining  $n - 1$  choices  $\binom{n-1}{k-1}$ . Then assume the item set aside is not in the selected subset of  $k$  items. Select the rest of the  $k$  items from the remaining  $n - 1$  choices  $\binom{n-1}{k}$ . These two disjoint sets add up to the original selection of  $k$  items from  $n$  choices

$$\binom{2n}{2} = 2\binom{n}{2} + n^2 \quad (7)$$

**LHS:** All possible ways to choose 2 items from a set of  $2n$  choices.

**RHS:** There are 3 ways to choose 2 items from a set of  $2n$  choices if you split the  $2n$  choices into 2 sets of  $n$  choices. You can choose both from the first set of  $n$  choices  $\binom{n}{2}$ , choose both from the second set of  $n$  choices  $\binom{n}{2}$ , or you can choose 1 from each set  $\binom{n}{1}^2 = n^2$ . Add these disjoint sets together to get LHS.

$$\sum_{k=0}^n k \binom{n}{k} = n2^{n-1} \quad (8)$$

**LHS:**

**RHS:**