Lorentz Symmetry of spin $\frac{1}{2}$ fields

Do a Lorentz transformation

$$x^\mu \to x'^\mu = \Lambda^\mu{}_\nu x^\nu, \text{ where } \Lambda^\mu{}_\rho \Lambda^\nu{}_\sigma g_{\mu\nu} = g_{\rho\sigma}$$

then $\psi'_a(x') = S_{ab}(\Lambda)\psi_b(x)$, note that $\Lambda^\mu{}_\nu$ are constants.

(recall that for scalar field $\phi'(x') = \phi(x)$,
for vector field $A'^\mu(x') = \Lambda^\mu{}_\nu A^\nu(x)$)

Since the theory should not change after a Lorentz transformation, then $\psi'(x')$ should also satisfies Dirac equation.

$$(i\gamma^\mu \partial'_\mu - m)\psi'(x') = 0.$$

times S from the left to $(i\gamma^\mu \partial_\mu - m)\psi(x) = 0$

$$S(i\gamma^\mu \partial_\mu - m)\psi(x) = 0$$

and insert $S^{-1}S = 1$.

$$\Rightarrow \quad S(i\gamma^\mu S^{-1} S \partial_\mu - m)\psi(x) = 0$$

$$\Rightarrow \quad i S\gamma^\mu S^{-1}\partial_\mu(S\psi(x)) - mS\psi(x) = 0$$

(note that $S(\Lambda)\partial_\mu = \partial_\mu S(\Lambda)$, since S is a 4×4 constant matrix)

$$\Rightarrow \quad (i S\gamma^\mu S^{-1}\partial_\mu - m)\psi'(x') = 0$$

So we have

$$S\gamma^\mu S^{-1}\partial_\mu = \gamma^\mu \partial'_\mu.$$

Since $\partial_\mu = \frac{\partial}{\partial x^\mu} = \frac{\partial x'^\nu}{\partial x^\mu}\frac{\partial}{\partial x'^\nu} = \Lambda^\nu{}_\mu \partial'_\nu$

then $S\gamma^\mu S^{-1}\Lambda^\nu{}_\mu \partial'_\nu = \gamma^\mu \partial'_\mu$

$$\Rightarrow \quad S\gamma^\mu S^{-1} \Lambda^\nu{}_\mu = \gamma^\nu$$

$$\Rightarrow \quad S^{-1} S \gamma^\mu S^{-1} S \Lambda^\nu{}_\mu = S^{-1}\gamma^\nu S$$

$$\Rightarrow \quad S^{-1}\gamma^\nu S = \Lambda^\nu{}_\mu \gamma^\mu.$$

This is the meaning of γ^μ behaves as a Lorentz vector.

Now consider an infinitesimal Lorentz transformation,

$\Lambda^\mu{}_\nu = \delta^\mu{}_\nu + \varepsilon^\mu{}_\nu$, and $\varepsilon_{\mu\nu} \equiv g_{\mu\lambda}\varepsilon^\lambda{}_\nu$

and we have shown before that $\varepsilon_{\mu\nu} = -\varepsilon_{\nu\mu}$.

To first order in $\varepsilon_{\mu\nu}$, $S(\Lambda)$ must have the form that

$$S(\Lambda) \simeq 1 - \frac{i}{4}\varepsilon^{\mu\nu}\sigma_{\mu\nu}$$

where $\sigma_{\mu\nu}$ are 4x4 matrices, $\sigma_{\mu\nu} = -\sigma_{\nu\mu}$.

the factor $(-\frac{i}{4})$ is introduced by convention.

Now let's find $\sigma_{\mu\nu}$.

Then from $S^{-1}\gamma^\nu S = \Lambda^\nu{}_\mu \gamma^\mu$, to first order in ε

$$\Rightarrow \quad \left(1 + \frac{i}{4}\varepsilon^{\alpha\beta}\sigma_{\alpha\beta}\right)\gamma^\nu\left(1 - \frac{i}{4}\varepsilon^{\rho\sigma}\sigma_{\rho\sigma}\right) \simeq (\delta^\nu{}_\mu + \varepsilon^\nu{}_\mu)\gamma^\mu$$

$$\Rightarrow \quad \left(\gamma^\nu + \frac{i}{4}\varepsilon^{\alpha\beta}\sigma_{\alpha\beta}\gamma^\nu\right)\left(1 - \frac{i}{4}\varepsilon^{\rho\sigma}\sigma_{\rho\sigma}\right) = \gamma^\nu + \varepsilon^\nu{}_\mu\gamma^\mu$$

$$\Rightarrow \quad \frac{i}{4}\varepsilon^{\alpha\beta}\sigma_{\alpha\beta}\gamma^\nu - \frac{i}{4}\gamma^\nu\varepsilon^{\rho\sigma}\sigma_{\rho\sigma} \simeq \varepsilon^\nu{}_\mu\gamma^\mu$$

$$\Rightarrow \quad -\frac{i}{4}\varepsilon^{\kappa\lambda}\left(\gamma^\nu\sigma_{\kappa\lambda} - \sigma_{\kappa\lambda}\gamma^\nu\right) = \varepsilon^\nu{}_\mu\gamma^\mu$$

the RHS $= \varepsilon^\nu{}_\mu\gamma^\mu = \varepsilon^{\nu\mu}\gamma_\mu = \varepsilon^{\kappa\lambda}(\delta^\nu_\kappa\gamma_\lambda)$

$$= \frac{1}{2}\varepsilon^{\kappa\lambda}\delta^\nu_\kappa\gamma_\lambda + \underbrace{\frac{1}{2}\varepsilon^{\kappa\lambda}\delta^\nu_\kappa\gamma_\lambda}_{-\frac{1}{2}\varepsilon^{\kappa\lambda}\delta^\nu_\lambda\gamma_\kappa}$$

$$\Rightarrow \quad -\frac{i}{4}\varepsilon^{\kappa\lambda}\left(\gamma^\nu\sigma_{\kappa\lambda} - \sigma_{\kappa\lambda}\gamma^\nu\right) = \frac{1}{2}\varepsilon^{\kappa\lambda}\left(\delta^\nu_\kappa\gamma_\lambda - \delta^\nu_\lambda\gamma_\kappa\right)$$

$$\Rightarrow \quad 2i\left(\delta^\nu_\kappa\gamma_\lambda - \delta^\nu_\lambda\gamma_\kappa\right) = [\gamma^\nu, \sigma_{\kappa\lambda}]$$

we can check that $\sigma_{\kappa\lambda} = \frac{i}{2}[\gamma_\kappa, \gamma_\lambda]$ is the solution to the above equation

check:

$$\text{RHS} = [\gamma^\nu, \tfrac{i}{2}[\gamma_\kappa, \gamma_\lambda]]$$

$$= \frac{i}{2}[\gamma^\nu, \gamma_\kappa\gamma_\lambda - \gamma_\lambda\gamma_\kappa]$$

$$= \frac{i}{2}(\gamma^\nu\gamma_\kappa\gamma_\lambda - \gamma^\nu\gamma_\lambda\gamma_\kappa - \gamma_\kappa\gamma_\lambda\gamma^\nu + \gamma_\lambda\gamma_\kappa\gamma^\nu)$$

use $\gamma_\mu\gamma_\nu + \gamma_\nu\gamma_\mu = 2g_{\mu\nu} \Rightarrow \gamma_\mu\gamma^\nu + \gamma^\nu\gamma_\mu = 2\delta^\nu_\mu$

$$\Rightarrow \text{RHS} = \frac{i}{2}\Big[(2\delta^\nu_\kappa - \gamma_\kappa\gamma^\nu)\gamma_\lambda - (2\delta^\nu_\lambda - \gamma_\lambda\gamma^\nu)\gamma_\kappa$$
$$- \gamma_\kappa(2\delta^\nu_\lambda - \gamma^\nu\gamma_\lambda) + \gamma_\lambda(2\delta^\nu_\kappa - \gamma^\nu\gamma_\kappa)\Big]$$

$$= \frac{i}{2}\Big[4\delta^\nu_\kappa\gamma_\lambda - 4\delta^\nu_\lambda\gamma_\kappa - \cancel{\gamma_\kappa\gamma^\nu\gamma_\lambda} + \cancel{\gamma_\lambda\gamma^\nu\gamma_\kappa}$$
$$+ \cancel{\gamma_\kappa\gamma^\nu\gamma_\lambda} - \cancel{\gamma_\lambda\gamma^\nu\gamma_\kappa}\Big]$$

$$= 2i(\delta^\nu_\kappa\gamma_\lambda - \delta^\nu_\lambda\gamma_\kappa)$$

$$= \text{LHS} \quad \checkmark$$

From $\sigma_{\mu\nu} = \frac{i}{2}[\gamma_\mu, \gamma_\nu] \Rightarrow \sigma_{\mu\nu}^{+} = -\frac{i}{2}(\gamma_\mu\gamma_\nu - \gamma_\nu\gamma_\mu)^{+} = -\frac{i}{2}(\gamma_\nu^{+}\gamma_\mu^{+} - \gamma_\mu^{+}\gamma_\nu^{+})$

$= \gamma_0 \sigma_{\mu\nu} \gamma_0$

$\Rightarrow \quad S^{+} = 1 + \frac{i}{4}\varepsilon^{\mu\nu}\sigma_{\mu\nu}^{+} = 1 + \frac{i}{4}\varepsilon^{\mu\nu}\gamma_0\sigma_{\mu\nu}\gamma_0$

while $S^{-1} = 1 + \frac{i}{4}\varepsilon^{\mu\nu}\sigma_{\mu\nu}, \quad \gamma_0^2 = 1$

$\Rightarrow \quad S^{+} = \gamma_0 S^{-1} \gamma_0$

Since $\psi'(x') = S\psi(x)$

then $\psi'^{+}(x') = \psi^{+}(x) S^{+}$

$\bar{\psi}'(x') = \psi'^{+}(x')\gamma^0 = \psi^{+}(x) S^{+}\gamma^0 = \psi^{+}(x)\gamma_0 S^{-1} = \bar{\psi}(x) S^{-1}$

So, Under Lorentz transformation

$$\bar{\psi}(x)\psi(x) \longrightarrow \bar{\psi}'(x')\psi'(x') = \bar{\psi}(x) S^{-1} S \psi(x) = \bar{\psi}(x)\psi(x)$$

$$i\bar{\psi}\gamma^\mu\partial_\mu\psi \longrightarrow i\bar{\psi}'(x')\gamma^\mu\partial'_\mu\psi'(x') = i\bar{\psi}(x) S^{-1}\gamma^\mu\partial'_\mu S\psi(x)$$

$$= i\bar{\psi}(x)\gamma^\nu\Lambda^\mu{}_\nu\partial'_\mu\psi(x) = i\bar{\psi}(x)\gamma^\mu\partial_\mu\psi(x)$$

Therefore, the Lagrangian

$$\mathcal{L} = i\bar{\psi}\gamma^\mu\partial_\mu\psi - m\bar{\psi}\psi$$

is invariant under Lorentz transformation.

$$\mathcal{L} = i\bar{\psi}\not{\partial}\psi - \bar{\psi} m \psi$$

$$= \bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi$$

$$\frac{\partial \mathcal{L}}{\partial \psi} = -m\bar{\psi}$$

$$\partial_\mu \frac{\partial \mathcal{L}}{\partial(\partial_\mu \psi)} = \partial_\mu(\bar{\psi} i \gamma^\mu) = i\gamma^\mu \partial_\mu \bar{\psi}$$

$$\Rightarrow \quad i\gamma^\mu \partial_\mu \bar{\psi} + m\bar{\psi} = 0$$

$$\Rightarrow \quad \bar{\psi}(i\overleftarrow{\not{\partial}} + m) = 0 \quad \checkmark$$

$$\frac{\partial \mathcal{L}}{\partial \bar{\psi}} = (i\gamma^\mu \partial_\mu - m)\psi$$

$$\partial_\mu \frac{\partial \mathcal{L}}{\partial(\partial_\mu \bar{\psi})} = \partial_\mu 0 = 0$$

$$\Rightarrow \quad (i\gamma^\mu \partial_\mu - m)\psi = 0. \quad \checkmark$$

if use ψ^+ as independent field, then

$$\mathcal{L} = i\psi^+ \gamma^0 \gamma^\mu \partial_\mu \psi - m\psi^+ \gamma^0 \psi$$

$$\frac{\partial \mathcal{L}}{\partial \psi^+} = i\gamma^0 \gamma^\mu \partial_\mu \psi - m\gamma^0 \psi$$

$$\frac{\partial \mathcal{L}}{\partial(\partial_\mu \psi^+)} = 0$$

$$\Rightarrow \gamma^0(i\gamma^\mu \partial_\mu \psi - m\psi) = 0$$

times γ^0 from the left and use $(\gamma^0)^2 = 1$

$$\Rightarrow \quad (i\gamma^\mu \partial_\mu - m)\psi = 0$$

spin of the Dirac field.

Do a Lorentz transformation

$$\psi'(x') = S(\Lambda)\,\psi(x)$$

$$\Rightarrow \psi'(x) = S(\Lambda)\,\psi(\Lambda^{-1}x)$$

(Recall for scalar field $\phi'(x') = \phi(x) \Rightarrow \phi'(x) = \phi(\Lambda^{-1}x)$

for vector field $A'^{\alpha}(x') = \Lambda^{\alpha}{}_{\beta} A^{\beta}(x) \Rightarrow A'^{\alpha}(x) = \Lambda^{\alpha}{}_{\beta} A^{\beta}(\Lambda^{-1}x)$)

For an infinitesimal transformation, $\Lambda^{\mu}{}_{\nu} \simeq \delta^{\mu}{}_{\nu} + \varepsilon^{\mu}{}_{\nu}$

where $\varepsilon_{\mu\nu} \equiv g_{\mu\lambda}\varepsilon^{\lambda}{}_{\nu}$ and $\varepsilon_{\delta\gamma} + \varepsilon_{\gamma\delta} = 0$

$$\Rightarrow \psi'(x) = \left(1 - \frac{i}{4}\varepsilon^{\mu\nu}\sigma_{\mu\nu}\right)\psi(x^{\rho} - \varepsilon^{\rho}{}_{\sigma}x^{\sigma})$$

$$= \left(1 - \frac{i}{4}\varepsilon^{\mu\nu}\sigma_{\mu\nu}\right)\left[\psi(x) - \frac{i}{2}\varepsilon^{\mu\nu}L_{\mu\nu}\psi(x)\right]$$

$$= \psi(x) - \frac{i}{2}\varepsilon^{\mu\nu}\left(L_{\mu\nu} + \frac{1}{2}\sigma_{\mu\nu}\right)\psi(x)$$

(recall for scalar field $\phi'(x) = \phi(\Lambda^{-1}x) = \phi\left((\delta^{\mu}{}_{\nu} - \varepsilon^{\mu}{}_{\nu})x^{\nu}\right) = \phi(x^{\mu} - \varepsilon^{\mu}{}_{\nu}x^{\nu})$

$$= \phi(x) - \varepsilon^{\mu}{}_{\nu}x^{\nu}\partial_{\mu}\phi(x)$$

$$= \phi(x) - \frac{i}{2}\varepsilon^{\mu\nu}L_{\mu\nu}\phi(x)$$

where $L_{\mu\nu} \equiv i(x_{\mu}\partial_{\nu} - x_{\nu}\partial_{\mu})$

$L_{\mu\nu}$ is antisymmetric in $\mu \leftrightarrow \nu$.

for vector field $A'^{\alpha}(x) = \Lambda^{\alpha}{}_{\beta}A^{\beta}(\Lambda^{-1}x) \simeq (\delta^{\alpha}{}_{\beta} + \varepsilon^{\alpha}{}_{\beta})(A^{\beta}(x) - \frac{i}{2}\varepsilon^{\mu\nu}L_{\mu\nu}A^{\beta}(x))$

$$= \left[\delta^{\alpha}{}_{\beta} - \frac{i}{2}\varepsilon^{\mu\nu}(\Sigma_{\mu\nu})^{\alpha}{}_{\beta}\right](A^{\beta}(x) - \frac{i}{2}\varepsilon^{\mu\nu}L_{\mu\nu}A^{\beta}(x))$$

$$= A^{\alpha}(x) - \frac{i}{2}\varepsilon^{\mu\nu}\left[L_{\mu\nu}\delta^{\alpha}{}_{\beta} + (\Sigma_{\mu\nu})^{\alpha}{}_{\beta}\right]A^{\beta}(x)$$

where $(\Sigma_{\mu\nu})^{\alpha}{}_{\beta} = i(\delta^{\alpha}{}_{\mu}g_{\nu\beta} - \delta^{\alpha}{}_{\nu}g_{\beta\mu})$

$\Sigma_{\mu\nu}$ is antisymmetric in $\mu \leftrightarrow \nu$.)

$$\Rightarrow \delta_0\psi(x) = -\frac{i}{2}\varepsilon^{\mu\nu}\left(L_{\mu\nu} + \frac{1}{2}\sigma_{\mu\nu}\right)\psi(x)$$

write the spinor indices explicitly,

$$\delta_0\psi_a(x) = -\frac{i}{2}\varepsilon^{\mu\nu}\left[L_{\mu\nu}\delta_{ab} + \frac{1}{2}(\sigma_{\mu\nu})_{ab}\right]\psi_b(x)$$

$\left(L_{\mu\nu} + \frac{1}{2}\sigma_{\mu\nu}\right)$ are the generators for infinitesimal Lorentz transformation of a Dirac field.

(note that $\sigma_{\mu\nu} = \frac{i}{2}[\gamma_{\mu}, \gamma_{\nu}]$ is antisymmetric in $\mu \leftrightarrow \nu$)

(recall for scalar field, $\delta_0 \phi(x) = \phi'(x) - \phi(x) = -\frac{i}{2}\varepsilon^{\mu\nu} L_{\mu\nu}\phi(x)$

for vector field, $\delta_0 A^\alpha(x) = -\frac{i}{2}\varepsilon^{\mu\nu}(L_{\mu\nu} + \Sigma_{\mu\nu})A^\alpha(x)$

where $\Sigma_{\mu\nu}A^\alpha \equiv (\Sigma_{\mu\nu})^\alpha{}_\beta A^\beta$, $L_{\mu\nu}A^\alpha \equiv L_{\mu\nu}\delta^\alpha{}_\beta A^\beta$

$L_{\mu\nu}$ are the generators of infinitesimal Lorentz transformation of a scalar field,

$(L_{\mu\nu} + \Sigma_{\mu\nu})$ are the generators of infinitesimal Lorentz transformation of a vector field.)

For the pure spatial part (i.e., for pure Lorentz rotation, no boost)

Define $L^k \equiv \frac{1}{2}\varepsilon^{ijk}L_{ij}$

$S^k \equiv \frac{1}{2}\varepsilon^{ijk}\frac{1}{2}\sigma_{ij}$

$\Rightarrow$ $L^1 = L_{23}$, $L^2 = L_{31}$, $L^3 = L_{12}$

$S^1 = \frac{1}{2}\sigma_{23}$, $S^2 = \frac{1}{2}\sigma_{31}$, $S^3 = \frac{1}{2}\sigma_{12}$

we have shown before (when we do scalar field)

$$[L^i, L^j] = i\varepsilon^{ijk}L^k$$

For $[S^i, S^j]$,

$$[S^1, S^1] = [S^2, S^2] = [S^3, S^3] = 0$$

use $\sigma_{ij} = \frac{i}{2}[\gamma_i, \gamma_j] = i\gamma_i\gamma_j$ for $i \neq j$.

$$\Rightarrow [S^1, S^2] = \frac{1}{4}[\sigma_{23}, \sigma_{31}] = \frac{1}{4}i^2[\gamma_2\gamma_3, \gamma_3\gamma_1]$$

$$= \frac{1}{4}i^2(\gamma_2\gamma_3\gamma_3\gamma_1 - \gamma_3\gamma_1\gamma_2\gamma_3)$$

$$= \frac{1}{4}i^2(-\gamma_2\gamma_1 + \gamma_1\gamma_2)$$

$$= \frac{1}{2}i^2\gamma_1\gamma_2$$

$$= \frac{i}{2}\sigma_{12} = i\varepsilon^{123}S^3$$

$$[S^2, S^3] = \frac{1}{4}[\sigma_{31}, \sigma_{12}] = \frac{1}{4} i^2 [\gamma_3\gamma_1, \gamma_1\gamma_2]$$

$$= \frac{1}{4} i^2 (\gamma_3\gamma_1\gamma_1\gamma_2 - \gamma_1\gamma_2\gamma_3\gamma_1)$$

$$= \frac{1}{4} i^2 (-\gamma_3\gamma_2 + \gamma_2\gamma_3)$$

$$= \frac{i^2}{2}\gamma_2\gamma_3$$

$$= \frac{i}{2}\sigma_{23}$$

$$= i\varepsilon^{231} S^1$$

$$[S^3, S^1] = \frac{1}{4}[\sigma_{12}, \sigma_{23}] = \frac{1}{4} i^2 [\gamma_1\gamma_2, \gamma_2\gamma_3]$$

$$= \frac{1}{4} i^2 (\gamma_1\gamma_2\gamma_2\gamma_3 - \gamma_2\gamma_3\gamma_1\gamma_2)$$

$$= \frac{1}{4} i^2 (-\gamma_1\gamma_3 + \gamma_3\gamma_1)$$

$$= \frac{1}{4} i^2 \, 2\gamma_3\gamma_1$$

$$= \frac{i}{2}\sigma_{31}$$

$$= i\varepsilon^{312} S^2$$

$$\Rightarrow \quad [S^i, S^j] = i\varepsilon^{ijk} S^k$$

Since $[\mathbb{1}, S^i] = 0$, then $[L^i, S^j] = 0$

Define $J^i \equiv L^i + S^i \Rightarrow [J^i, J^j] = i\varepsilon^{ijk} J^k$

So J^i is total angular momentum, L^i is orbital angular momentum, S^i is spin angular momentum.

Since $\sum_{i=1}^{3} S^i S^i = S^1S^1 + S^2S^2 + S^3S^3 = \frac{1}{4}(\sigma_{23}\sigma_{23} + \sigma_{31}\sigma_{31} + \sigma_{12}\sigma_{12})$

$$= \frac{1}{4} i^2 (\gamma_2\gamma_3\gamma_2\gamma_3 + \gamma_3\gamma_1\gamma_3\gamma_1 + \gamma_1\gamma_2\gamma_1\gamma_2)$$

$$= \frac{1}{4} i^2 (-1 - 1 - 1)$$

$$= \frac{3}{4}$$

$$= \frac{1}{2}\left(\frac{1}{2} + 1\right) \Rightarrow \text{spin is } \frac{1}{2}$$

recall for scalar field, $L^i \equiv \frac{1}{2}\varepsilon^{ijk} L_{jk} \Rightarrow [L^i, L^j] = i\varepsilon^{ijk} L^k$

for vector field, $L^i \equiv \frac{1}{2}\varepsilon^{ijk} L_{jk}$, $S^i = \frac{1}{2}\varepsilon^{ijk} \Sigma_{jk}$

that is, $(L^i)^m{}_n = \frac{i}{2}\varepsilon^{ijk}(x_j \partial_k - x_k \partial_j)\delta^m_n$

$(S^i)^m{}_n = -i\varepsilon^{imn}$

$\Rightarrow [L^i, L^j] = i\varepsilon^{ijk} L^k$, $[S^i, S^j] = i\varepsilon^{ijk} S^k$

$[L^i, S^j] = 0$ that is, $[S^i, S^j]^a{}_b = i\varepsilon^{ijk}(S^k)^a{}_b$

$[L^i + S^i, L^j + S^j] = 0$

$\sum_{k=1}^{3} (S^k S^k)^m{}_n = 2\delta^m_n = 1(1+1)\delta^m_n \Rightarrow$ spin is 1

Bilinear Covariants

use $\begin{cases} S(\Lambda)^{-1}\gamma^\mu S(\Lambda) = \Lambda^\mu{}_\nu \gamma^\nu, \\ \psi'(x') = S(\Lambda)\psi(x) \\ \bar\psi'(x') = \bar\psi(x) S^{-1} \end{cases}$

let's find out how $\bar\psi \Gamma \psi$ transform under Lorentz transformation, where Γ's are 1, γ^μ, $\sigma^{\mu\nu} = \frac{i}{2}[\gamma^\mu, \gamma^\nu]$, $\gamma_5\gamma^\mu$ and γ_5

$$\Rightarrow \quad \bar\psi'(x')\psi'(x') = \bar\psi(x) S^{-1} S \psi(x) = \bar\psi(x)\psi(x),$$ so $\bar\psi\psi$ transform as a scalar

$$\bar\psi'(x')\gamma^\mu\psi'(x') = \bar\psi(x) S^{-1}\gamma^\mu S\psi(x) = \Lambda^\mu{}_\nu \bar\psi(x)\gamma^\nu\psi(x)$$

so, $\bar\psi\gamma^\mu\psi$ transform as a vector

$$\bar\psi'(x')\sigma^{\mu\nu}\psi'(x') = \bar\psi'(x')\frac{i}{2}[\gamma^\mu, \gamma^\nu]\psi'(x')$$

$$= \bar\psi(x) S^{-1}\frac{i}{2}(\underbrace{\gamma^\mu\gamma^\nu}_{SS^{-1}} - \underbrace{\gamma^\nu\gamma^\mu}_{SS^{-1}}) S\psi(x)$$

$$= \frac{i}{2}\bar\psi(x)\left(\Lambda^\mu{}_\alpha\gamma^\alpha\Lambda^\nu{}_\beta\gamma^\beta - \Lambda^\nu{}_\alpha\gamma^\alpha\Lambda^\mu{}_\beta\gamma^\beta\right)\psi(x)$$

$$= \Lambda^\mu{}_\alpha\Lambda^\nu{}_\beta\bar\psi(x)\sigma^{\alpha\beta}\psi(x)$$

so, $\bar\psi\sigma^{\mu\nu}\psi$ transform as a tensor

use $\gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3$

$$= \frac{i}{4!}\varepsilon_{\mu\nu\rho\sigma}\gamma^\mu\gamma^\nu\gamma^\rho\gamma^\sigma$$

where $\varepsilon_{\mu\nu\rho\sigma} = \begin{cases} +1, & \text{for even permutation of } \varepsilon_{0123} \\ -1, & \text{for odd} \quad \text{- - - - - - - -} \\ 0, & \text{otherwise.} \end{cases}$

$$\Rightarrow \quad S^{-1}\gamma_5 S = S^{-1}(i\gamma^0\gamma^1\gamma^2\gamma^3)S = \frac{i}{4!}\varepsilon_{\mu\nu\rho\sigma} S^{-1}\gamma^\mu\underbrace{\gamma^\nu\gamma^\rho}_{SS^{-1}}\gamma^\sigma S$$

$$= \frac{i}{4!}\varepsilon_{\mu\nu\rho\sigma}\Lambda^\mu{}_\alpha\Lambda^\nu{}_\beta\Lambda^\rho{}_\varphi\Lambda^\sigma{}_\pi\gamma^\alpha\Lambda^\beta\Lambda^\varphi\Lambda^\pi$$

$$= i\,\varepsilon_{\mu\nu\rho\sigma}\Lambda^\mu{}_0\Lambda^\nu{}_1\Lambda^\rho{}_2\Lambda^\sigma{}_3\gamma^0\gamma^1\gamma^2\gamma^3$$

$$= \varepsilon_{\mu\nu\rho\sigma}\Lambda^\mu{}_0\Lambda^\nu{}_1\Lambda^\rho{}_2\Lambda^\sigma{}_3\gamma_5$$

$$= \mathrm{Det}(\Lambda)\gamma_5$$

$$\Rightarrow \quad \bar\psi'(x')\gamma_5\psi'(x') = \bar\psi(x) S^{-1}\gamma_5 S\psi(x) = \mathrm{Det}(\Lambda)\bar\psi(x)\gamma_5\psi(x)$$

Recall $\Lambda^\alpha{}_\gamma \Lambda^\beta{}_\delta \eta_{\alpha\beta} = \eta_{\gamma\delta}$ is the condition that a Lorentz transformation satisfies.

then
$$(\Lambda^T)_\gamma{}^\alpha \eta_{\alpha\beta} \Lambda^\beta{}_\delta = \eta_{\gamma\delta} \Rightarrow \Lambda^T \eta \Lambda = \eta$$
$$\Rightarrow \quad \mathrm{Det}(\Lambda^T \eta \Lambda) = \mathrm{Det}(\eta)$$

Since $\mathrm{Det}(A) = \mathrm{Det}(A^T)$ and $\mathrm{Det}(AB) = \mathrm{Det}(A)\,\mathrm{Det}(B)$

then $[\mathrm{Det}(\Lambda)]^2 = 1$

$$\Rightarrow \quad \mathrm{Det}(\Lambda) = \pm 1$$

Also, from $\Lambda^\alpha{}_\gamma \Lambda^\beta{}_\delta \eta_{\alpha\beta} = \eta_{\gamma\delta}$

$$\Rightarrow \quad \Lambda^0{}_0 \Lambda^0{}_0 \eta_{00} + \Lambda^i{}_0 \Lambda^i{}_0 \eta_{ii} = \eta_{00} = 1$$
$$\Rightarrow \quad (\Lambda^0{}_0)^2 - \sum_{i=1}^{3} (\Lambda^i{}_0)^2 = 1$$
$$\Rightarrow \quad (\Lambda^0{}_0)^2 = 1 + \sum_{i=1}^{3} (\Lambda^i{}_0)^2 \geq 1$$

While for identity matrix (i.e., no transformation)
$$\Lambda^\alpha{}_\beta = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$
$$\Rightarrow \quad \Lambda^0{}_0 = 1, \quad \mathrm{Det}(\Lambda) = 1.$$

The subclass of the Lorentz transformations satisfying $\Lambda^0{}_0 \geq 1$ & $\mathrm{Det} = 1$ forms a group called the proper inhomogeneous or homogenous Lorentz group.

$$x'^\alpha = \Lambda^\alpha{}_\beta x^\beta + \underset{\neq 0}{a^\alpha} \qquad x'^\alpha = \Lambda^\alpha{}_\beta x^\beta + \underset{0}{a^\alpha}$$

The space inversion $\Lambda^\alpha{}_\beta = \begin{pmatrix} 1 & & & \\ & -1 & & \\ & & -1 & \\ & & & -1 \end{pmatrix}$,

the time inversion $\Lambda^\alpha{}_\beta = \begin{pmatrix} -1 & & & \\ & 1 & & \\ & & 1 & \\ & & & 1 \end{pmatrix}$

and space-time inversion $\Lambda^\alpha{}_\beta = \begin{pmatrix} -1 & & & \\ & -1 & & \\ & & -1 & \\ & & & -1 \end{pmatrix}$

are not elements of proper Lorentz group.

$\Lambda^\alpha{}_\beta = \begin{pmatrix} 1 & & & \\ & -1 & & \\ & & -1 & \\ & & & -1 \end{pmatrix}$ is often called parity transformation, denoted as P

$\Lambda^\alpha{}_\beta = \begin{pmatrix} -1 & & & \\ & 1 & & \\ & & 1 & \\ & & & 1 \end{pmatrix}$ is often called time inversion transformation, denoted as T.

---

we say that $\bar{\psi}(x)\gamma_5\psi(x)$ transforms as a pseudoscalar under Lorentz transformation.

For $\bar{\psi}(x)\gamma_5\gamma^\mu\psi(x)$, it transform as

$$\bar{\psi}'(x')\gamma_5\gamma^\mu\psi'(x') = \bar{\psi}(x)S^{-1}\gamma_5\underbrace{\gamma^\mu S}_{SS^{-1}}\psi(x) = \Lambda^\mu{}_\nu \mathrm{Det}(\Lambda)\bar{\psi}(x)\gamma_5\gamma^\nu\psi(x)$$

we say that $\bar{\psi}(x)\gamma_5\gamma^\mu\psi(x)$ transform as a pseudovector (or, axial vector) under Lorentz transformation.

Notes:

① Because Parity is violated in weak interaction, but is not violated in electromagnetic interaction, the bilinear terms including γ_5 only appear in the former.

② Any 4×4 matrices can be constructed from the 16 linearly independent matrices, 1, γ^μ, γ_5, $\sigma^{\mu\nu}$, $\gamma_5\gamma^\mu$. (i.e., they form a set of basis)

③ In the bilinear combinations, $\bar{\psi}\psi$, $\bar{\psi}\gamma_5\psi$, $\bar{\psi}\gamma^\mu\psi$, $\bar{\psi}\gamma_5\gamma_\mu\psi$, $\bar{\psi}\sigma^{\mu\nu}\psi$, the $\bar{\psi}$ and ψ can be different Dirac spinor, i.e. $\bar{\psi}_1\psi_2$, $\bar{\psi}_1\gamma_5\psi_2$, $\bar{\psi}_1\gamma_5\gamma^\mu\psi_2$, $\bar{\psi}_1\sigma^{\mu\nu}\psi_2$, $\bar{\psi}_1\gamma^\mu\psi_2$

④ Since $(\bar{\psi}\gamma_5\psi)^\dagger = \psi^\dagger\gamma_5^\dagger\gamma_0^\dagger\psi = \psi^\dagger\gamma_5\gamma_0\psi = -\psi^\dagger\gamma_0\gamma_5\psi = -\bar{\psi}\gamma_5\psi$, sometimes it is more convinient to use $i\gamma_5$ as a basis, because $(\bar{\psi}i\gamma_5\psi)^\dagger = \bar{\psi}i\gamma_5\psi$