Survival Analysis

Chao Cheng

October 21, 2022

Contents

1 Basic knowledge

Let T denote the time to an event that we are interested in. Then we know the c.d.f.

$$F_T(t) = P(T \leq t),$$

and the corresponding p.d.f.

$$f_T(t) = \frac{\mathrm{d}}{\mathrm{d}t} F_T(t).$$

Here to simplify the discussion, we assume T is a continuous random variable. In the context of survival analysis, the *event* often refers to death. Then T represents the lifespan of the subject. So $F_T(t)$ represents the probability that the death occurs before t. In another word, we know the probability that the subject survives passes t is

$$S_T(t) = 1 - F_T(t) = P(T > t).$$

$S_T(t)$ is often called the survival function? and clearly

$$f_T(t) = -\frac{\mathrm{d}}{\mathrm{d}t} S_T(t).$$

The **hazard function** $h(t)$ is defined as

$$h(t) = \lim_{\Delta \to 0} \frac{P(T \leq t + \Delta | T > t)}{\Delta} = \lim_{\Delta \to 0} \frac{F_T(t + \Delta) - F_T(t)}{\Delta \cdot S_T(t)} = \frac{f_T(t)}{S_T(t)}.$$

$h(t)$ represents the instant hazard? unified probability? that the subject will be dead instantly after t given the fact that it's alive at t. And the **cummulative hazard function** is

$$H(t) = \int_0^t h(x)\,\mathrm{d}x = \int_0^t \frac{f_T(x)}{S_T(x)}\mathrm{d}x = \int_0^t \frac{-\mathrm{d}S_T(x)}{S_X(t)} = -\log(S_T(x))|_0^t = -\log(S_T(t)).$$

References