

# Cox Proportional Hazard Model

Chao Cheng

December 26, 2022

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Estimation</b>	<b>2</b>
2.1	What is $L_1(\beta)$	3
2.1.1	Intuition: profile likelihood perspective	3
2.1.2	Intuition: conditional distribution perspective	4
2.2	What if there is censoring?	4
2.3	What if there are tied event times?	4
2.3.1	Exact partial likelihood	4
2.3.2	Breslow's approximation	5
2.3.3	Efron's approximation	5
<b>3</b>	<b>Inference</b>	<b>5</b>

## 1 Introduction

In this note we will talk about the Cox's proportional hazards (Cox's PH) model. Suppose we observe some non-informatively right-censored data  $(U, \delta)$  with covariate vector  $Z$ . That is, for subject  $i$ , the covariate vector is  $Z_i$ , survival time  $T_i$  and censoring time  $C_i$ . The observed data is  $(U_i, \delta_i)$  where  $U_i = \min(T_i, C_i)$  and  $\delta_i = 1(T_i \leq C_i)$ . Also  $T_i \perp C_i | Z_i$ .

And now we want to model the relationship between  $Z$  and  $T$ . One way to do that is to incorporate  $Z$  into the hazard function  $h(\cdot)$ , e.g.,

$$T \sim \text{Exp}(\lambda_Z) \implies h(t) = \lambda_Z \stackrel{\Delta}{=} e^{\alpha + \beta Z} = \lambda_0 e^{\beta Z},$$

where  $\lambda_0 = e^\alpha$  can be viewed as a baseline hazard. If  $\beta = 0$  then  $Z$  is not associated with  $T$ .

We can generalize this idea as

$$h(t|Z) = h_0(t) \times g(Z).$$

So the hazard can be factorized and this model is sometimes called a “multiplicative intensive model” or “multiplicative hazard model” or “proportional hazard model” because this factorization implies that

$$\frac{h(t|Z = z_1)}{h(t|Z = z_2)} = \frac{g(z_1)}{g(z_2)}.$$

The hazard ratio is constant with respect to  $t$ , hence the (constant) proportional hazard. So in our previous model (the exponential survival time), the hazard ratio is

$$\frac{h(t|Z = z_1)}{h(t|Z = z_2)} = e^{\beta(z_1 - z_2)}.$$

Also this exponential form of  $g(Z)$

$$h(t|Z) = h_0(t) \cdot e^{\beta Z} \quad (1)$$

is the **Cox's PH** model.

## 2 Estimation

(1) implies that

$$\begin{aligned} S(t|Z) &= \exp(-H(t|Z)) \\ &= \exp\left(-\int_0^t h(u|Z) du\right) \\ &= \exp\left(-\int_0^t h_0(u) du \cdot g(Z)\right) \\ &= (S_0(t))^{g(Z)} = (S_0(t))^{\exp(\beta Z)}, \end{aligned}$$

where  $S_0(t) = \exp\left(-\int_0^t h_0(u) du\right)$ , the survival function for  $Z = 0$ , hence  $S(t|Z = 0)$ . Also remember that  $f(t|Z) = h(t|Z) S(t|Z)$ . Thus, given  $n$  independent data  $(u_i, \delta_i, z_i)$ , the likelihood ([one can refer to our previous notes about survival analysis.](#)) is

$$\begin{aligned} L(\beta, h_0(\cdot)) &= \prod_{i=1}^n (f(u_i|z_i))^{\delta_i} (S(u_i|z_i))^{1-\delta_i} = \prod_{i=1}^n h(u_i|z_i)^{\delta_i} S(u_i|z_i) \\ &= \prod_{i=1}^n (h_0(u_i) e^{\beta z_i})^{\delta_i} \left( \exp\left(-\int_0^{u_i} h_0(t) dt\right) \right)^{\exp(\beta z_i)} \\ &= \text{function}(data, h_0(\cdot), \beta). \end{aligned} \quad (2)$$

If  $h_0(\cdot)$  is allowed to be “arbitrary”, then the “parameter space “ is

$$\mathcal{H} \times \mathcal{R}^p = \left\{ (h(\cdot), \beta) \left| h_0(\cdot) \geq 0, \int_0^\infty h_0(t) dt = \infty, \beta \in \mathcal{R}^p \right. \right\},$$

where  $\int_0^\infty h_0(t) dt = \infty$  ensures that  $S_0(\infty) = 0$ .

In general this likelihood is hard to maximize. And Cox proposed this idea: to factor  $L(\beta, h_0(\cdot))$  as

$$L(\beta, h_0(\cdot)) = L_1(\beta) \times L_2(\beta, h_0(\cdot)),$$

where  $L_1$  only depends on  $\beta$  and its maximization ( $\hat{\beta}$ ) enjoys nice properties such as consistency and asymptotic normality while  $L_2$  contains relatively little information about  $\beta$ . And this  $L_1$  is called a **partial likelihood**.

## 2.1 What is $L_1(\beta)$

In this section we introduce the  $L_1$  proposed by Cox. First let's assume there are **NO tied** nor censoring observations. And define the distinct times of failure  $\tau_1 < \tau_2 < \dots$ . Denote

$$R_j = \{i | U_i \geq \tau_j\} = \text{risk set at } \tau_j,$$

and

$$Z_{(j)} = \text{value of } Z \text{ for the subject who fails at } \tau_j.$$

we can reconstruct the data from  $\{\tau_j\}$ ,  $\{R_j\}$  and  $\{Z_{(j)}\}$ . And  $L_1$  is defined as

$$L_1(\beta) \triangleq \prod_j \left\{ \frac{e^{\beta Z_{(j)}}}{\sum_{l \in R_j} e^{\beta Z_l}} \right\} \quad (3)$$

Note that under this setting (no tie, no censor), the full likelihood (2) becomes

$$L(\beta, h_0(\cdot)) = \prod_{i=1}^n h_0(u_i) e^{\beta z_i} \left( \exp \left( - \int_0^{u_i} h_0(t) dt \right) \right)^{\exp(\beta z_i)}.$$

Furthermore, we can assume  $u_i = \tau_i$ , i.e. the data has been sorted based on survival time. And use the KM idea, i.e. assume the survival function is **discrete** with baseline hazard value  $h_j$  at  $u_j$ . Then this likelihood becomes

$$L(\beta, h_1, \dots, h_n) = \prod_{i=1}^n h_i e^{\beta z_i} \exp \left( - \sum_{j=1}^i h_j \right)^{\exp(\beta z_i)}. \quad (4)$$

Note that, in previous notes we have deduct that in discrete case, for any  $t \in [v_j, v_{j+1})$ :

$$H(t) = \sum_{i=1}^j h_i \quad S(t) = \prod_{i=1}^j (1 - h_i).$$

Here in (4) we use the approximation that  $e^{-h_j} \approx 1 - h_j$  when  $h_j$  is close to 0.

### 2.1.1 Intuition: profile likelihood perspective

We can use the method of profile likelihood: That is, for any given  $\beta$ , we maximize  $L$  (or equivalently,  $\log L$ ) over  $h_j$ s so the result is a function of  $\beta$ . Taking derivative, we have

$$\frac{\partial \log L}{\partial h_j} = \frac{1}{h_j} - \sum_{i \leq j} \exp(\beta z_i), \quad j = 1, \dots, n.$$

Set them to 0 we have  $\hat{h}_j = 1 / \sum_{i \leq j} \exp(\beta z_i)$ . And the log profile likelihood of  $\beta$  is

$$\begin{aligned}
\log L_{profile}(\beta) &= \log \left\{ \prod_{i=1}^n \frac{\exp(\beta z_i)}{\sum_{k \leq i} \exp(\beta z_k)} \exp \left( - \sum_{j=1}^i \frac{1}{\sum_{k \leq j} \exp(\beta z_k)} \right)^{\exp(\beta z_i)} \right\} \\
&= \sum_{i=1}^n \left\{ \log \left\{ \frac{\exp(\beta z_i)}{\sum_{k \leq i} \exp(\beta z_k)} \exp \left( - \sum_{j=1}^i \frac{1}{\sum_{k \leq j} \exp(\beta z_k)} \right)^{\exp(\beta z_i)} \right\} \right\} \\
&= \sum_{i=1}^n \left\{ \log \left( \frac{\exp(\beta z_i)}{\sum_{k \leq i} \exp(\beta z_k)} \right) - \exp(\beta z_i) \cdot \left( \sum_{j=1}^i \frac{1}{\sum_{k \leq j} \exp(\beta z_k)} \right) \right\} \\
&= \sum_{i=1}^n \left\{ \log \left( \frac{\exp(\beta z_i)}{\sum_{k \leq i} \exp(\beta z_k)} \right) \right\} - \sum_{i=1}^n \sum_{j=1}^i \frac{\exp(\beta z_i)}{\sum_{k \leq j} \exp(\beta z_k)},
\end{aligned} \tag{5}$$

where the second part of last equation can be reduced to  $-n$ , which means

$$L_{profile}(\beta) \propto \prod_{i=1}^n \frac{\exp(\beta z_i)}{\sum_{k \leq i} \exp(\beta z_k)}.$$

And this is what Cox uses as  $L_1(\beta)$ .

### 2.1.2 Intuition: conditional distribution perspective

To be added.

## 2.2 What if there is censoring?

Then (3) is still used.

## 2.3 What if there are tied event times?

Through the probability of tie existence is 0 in the continuous time case, in real life it is pretty common.

### 2.3.1 Exact partial likelihood

The exact partial likelihood considers all the possible rankings for the tied observations. Specifically

$$L_1(\beta) = \prod_{j=1}^K \left\{ \sum_{(k_1, \dots, k_{d_j}) = (1, 2, \dots, d_j)} \prod_{i=1}^{d_j} \left\{ \frac{\exp(\beta z_{(j)}^{k_i})}{\sum_{l \in R_j} \exp(\beta z_l) - \sum_{s=i}^{d_j} \exp(\beta z_{(j)}^{(s)})} \right\} \right\}.$$

The computation of the exact partial likelihood gets out of hand pretty quickly as  $d_j$  increases. So some modification/approximation methods are proposed.

2.3.2 Breslow's approximation

2.3.3 Efron's approximation

## 3 Inference

## References