

Only One Event in Cox Model

Chao Cheng

September 27, 2026

Contents

1 Introduction	1
2 Only One Event Observed	1

1 Introduction

Let's consider the Cox model and there are **No tied** observations. And define the distinct times of failure (or censoring) $\tau_1 < \tau_2 < \dots < \tau_n$. Denote

$$R_j = \{i | U_i \geq \tau_j\} = \text{risk set at } \tau_j,$$

where $U_i = \min(T_i, C_i)$ is the observed time and

$$Z_{(j)} = \text{value of } Z \text{ for the subject who fails at } \tau_j.$$

The partial likelihood $L_1(\beta)$ is defined as

$$L_1(\beta) \triangleq \prod_j \left\{ \frac{e^{\beta Z_{(j)}}}{\sum_{l \in R_j} e^{\beta Z_l}} \right\} \quad (1)$$

In this notes, we will consider the clinical trial scenario. So $Z_i \in \{0, 1\}$ for $i = 1, \dots, n$. $Z_i = 0$ denotes control group and $Z_i = 1$ denotes the treatment group. And we will consider the case when there's only one event observed.

2 Only One Event Observed

If there's only one event observed, then (1) can be simplified as

$$L_1(\beta) = \frac{e^{\beta Z_j}}{\sum_{l \in R_j} e^{\beta Z_l}} = \frac{e^{\beta Z_j}}{n_{trt} e^{\beta \times 1} + n_{ctrl} e^{\beta \times 0}} = \frac{e^{\beta Z_j}}{n_{trt} e^{\beta} + n_{ctrl}}, \quad (2)$$

where n_{trt} is the number of observations in R_j that belongs to treatment group and n_{ctrl} is the number of observations in R_j that belongs to the control group.

To maximize (2), we can see

$$L_1(\beta) = \begin{cases} \frac{e^\beta}{n_{trt}e^\beta + n_{ctrl}} = \frac{1}{n_{trt}} - \frac{n_{ctrl}}{n_{trt}} \frac{1}{n_{trt}e^\beta + n_{ctrl}} & \text{if } Z_j = 1 \\ \frac{1}{n_{trt}e^\beta + n_{ctrl}} & \text{if } Z_j = 0 \end{cases}$$

If $Z_j = 1$, (2) is maximized at $\beta = \infty$. If $Z_j = 0$, (2) is maximized at $\beta = -\infty$.

References