

# Solution to Assignment 1

1.8: We have

$$\liminf A_n = \bigcup_n \bigcap_{k \geq n} A_k = \bigcup_n BC = BC$$

and

$$\limsup A_n = \bigcap_n \bigcup_{k \geq n} A_k = \bigcap_n (B \cup C) = B \cup C.$$

1.18: Since  $\Omega \in \mathcal{F}$ , if  $A \in \mathcal{F}$  then  $A^c = \Omega \setminus A \in \mathcal{F}$ . Thus  $\mathcal{F}$  is closed under complements. If  $A$  and  $B$  lie in  $\mathcal{F}$ , then by deMorgan's law

$$A \cup B = (A^c B^c)^c = (A^c \setminus B)^c,$$

which lies in  $\mathcal{F}$  as  $\mathcal{F}$  is closed under complements and differences. Thus  $\mathcal{F}$  is closed under unions.

1.24: Clearly  $\Omega \in \mathcal{A}$ , and  $\mathcal{A}$  is closed under complements by the symmetry of the definition. Suppose  $A$  and  $B$  lie in  $\mathcal{A}$ . If  $\#A < \infty$  and  $\#B < \infty$ , then  $\#(A \cup B) < \#A + \#B < \infty$  and so  $A \cup B \in \mathcal{A}$ . If  $\#A^c < \infty$ , then  $\#(A \cup B)^c = \#(A^c B^c) \leq \#A^c < \infty$  and so  $A \cup B \in \mathcal{A}$ . Similarly, if  $\#B^c < \infty$  then  $A \cup B \in \mathcal{A}$ . As one of these cases must hold,  $\mathcal{A}$  is closed under finite unions and must be a field. But  $\mathcal{A}$  is not closed under countable unions for if  $A_n = \{2n\} \in \mathcal{A}$ ,  $n \geq 1$ , then

$$\bigcup_n A_n = \{\text{even natural numbers}\} \notin \mathcal{A}.$$

So  $\mathcal{A}$  is not a  $\sigma$ -field.

1.34: Let  $\mathcal{C}$  be the class of all sets of the given form. Since  $\sigma$ -fields are closed under unions, intersections, and complements, if  $B, B' \in \mathcal{B}$  then

$$AB \cup A^c B' \in \sigma(\mathcal{B} \cup \{A\}),$$

and hence  $\mathcal{C} \subset \sigma(\mathcal{B} \cup \{A\})$ . So the desired result will follow if we show that  $\mathcal{C}$  is a  $\sigma$ -field. Since  $\Omega = A\Omega \cup A^c\Omega$ , we have  $\Omega \in \mathcal{C}$ . Next, if  $B, B' \in \mathcal{B}$ , then

$$(AB \cup A^c B')^c = AB^c \cup A^c B'^c \in \mathcal{C}.$$

Thus  $\mathcal{C}$  is closed under complements. Finally, if  $AB_n \cup A^c B'_n$ ,  $n \geq 1$ , are sets in  $\mathcal{C}$ , with  $B_n, B'_n$ ,  $n \geq 1$ , in  $\mathcal{B}$ , then

$$\bigcup_n (AB_n \cup A^c B'_n) = (A \cup_n B_n) \cup (A^c \cup_n B'_n),$$

which lies in  $\mathcal{C}$  as  $\cup_n B_n \in \mathcal{B}$  and  $\cup_n B'_n \in \mathcal{B}$ . Thus  $\mathcal{C}$  is a  $\sigma$ -field.

2.4: By monotonicity, the value  $P_1$  assigns to  $A$  must satisfy

$$P_1(A) \leq P_1(B) = P(B), \quad \forall B \in \mathcal{B} \text{ with } B \supset A.$$

So the largest possible value for  $P_1(A)$  will be

$$\pi^* = \inf_{B \in \mathcal{B}, B \supset A} P(B),$$

the outer measure of  $A$ . We will seek an extension with  $P_1(A) = \pi^*$ . Let  $A_n$ ,  $n \geq 1$ , be a sequence of sets in  $\mathcal{B}$  with  $A_n \supset A$  and  $P(A_n) \rightarrow \pi^*$ , and define

$$\tilde{A} = \bigcap_n A_n.$$

Since  $\tilde{A} \supset A$  and  $\tilde{A} \in \mathcal{B}$ , we have  $P(\tilde{A}) \geq \pi^*$ . By continuity, since  $\cap_{k=1}^n A_k \uparrow \tilde{A}$ ,

$$P(A_n) \leq P\left(\bigcap_{k=1}^n A_k\right) \rightarrow P(\tilde{A}),$$

and so

$$P(\tilde{A}) = \liminf_{n \rightarrow \infty} P\left(\bigcap_{k=1}^n A_k\right) \leq \liminf_{n \rightarrow \infty} P(A_n) = \pi^*.$$

Thus  $P(\tilde{A}) = \pi^*$ . So it seems we can make  $P_1(A)$  as large as possible by making  $A$  and  $\tilde{A}$  equivalent. Specifically, if  $B, B' \in \mathcal{B}$ , the extension will be given by

$$P_1(AB \cup A^c B') = P(\tilde{A}B) + P(\tilde{A}^c B').$$

To show that  $P_1$  is a probability measure, we first note that if  $A$  and  $B \in \mathcal{B}$  are disjoint, then  $P(\tilde{A}B) = 0$ . To see this, note that since  $A \subset \tilde{A}B^c$ , we have  $P(\tilde{A}B^c) \geq \pi^*$ , and so

$$\pi^* = P(\tilde{A}) = P(\tilde{A}B) + P(\tilde{A}B^c) \geq P(\tilde{A}B) + \pi^*.$$

Using this, if  $AB$  and  $AB'$  are disjoint with  $B, B' \in \mathcal{B}$ , then  $A$  and  $BB'$  are disjoint, and by the addition law

$$\begin{aligned} P(\tilde{A}B \cup \tilde{A}B') &= P(\tilde{A}B) + P(\tilde{A}B') - P(\tilde{A}BB') \\ &= P(\tilde{A}B) + P(\tilde{A}B'). \end{aligned}$$

Iterating this, if  $AB_k$ ,  $1 \leq k \leq n$ , are disjoint, then

$$P\left(\bigcup_{k=1}^n \tilde{A}B_k\right) = \sum_{k=1}^n P(\tilde{A}B_k).$$

Using this, we next show that  $P_1$  is  $\sigma$ -additive. Let  $D_n$ ,  $n \geq 1$ , be disjoint sets in  $\mathcal{B}_1$ , and write

$$D_n = AB_n \cup A^c B'_n, \text{ with } B_n, B'_n \in \mathcal{B}.$$

Then

$$D = \bigcup_n D_n = (A \cup_n B_n) \cup (A^c \cup_n B'_n),$$

and so

$$P_1(D) = P(\cup_n \tilde{A}B_n) + P(\cup_n \tilde{A}^c B'_n).$$

Since  $A^c B'_n$ ,  $n \geq 1$ , are disjoint and  $\tilde{A}^c B'_n \subset A^c B'_n$ , the sets  $\tilde{A}^c B'_n$ ,  $n \geq 1$  are disjoint, and so

$$P(\cup_n \tilde{A}^c B'_n) = \sum_n P(\tilde{A}^c B'_n).$$

By continuity and the identity above,

$$P(\cup_n \tilde{A}B_n) = \lim_{n \rightarrow \infty} P(\cup_{k=1}^n \tilde{A}B_k) = \lim_{n \rightarrow \infty} \sum_{k=1}^n P(\tilde{A}B_k) = \sum_n P(\tilde{A}B_n).$$

So

$$P_1(D) = \sum_n [P(\tilde{A}B_n) + P(\tilde{A}^c B'_n)] = \sum_n P_1(D_n).$$

For the other axioms, from the definition  $P_1(B) \geq 0$  for all  $B \in \mathcal{B}_1$ , and

$$P_1(\Omega) = P(\tilde{A}) + P(\tilde{A}^c) = 1.$$

Thus  $P_1$  is a probability measure on  $(\Omega, \mathcal{B}_1)$ .

2.8: In the example given in the problem,

$$\begin{aligned} P_1(\{a, b\}) &= P_1(\{d, c\}) = P_1(\{a, c\}) = P_1(\{b, d\}) \\ &= P_2(\{a, b\}) = P_2(\{d, c\}) = P_2(\{a, c\}) = P_2(\{b, d\}) = 1/2, \end{aligned}$$

and so  $P_1$  and  $P_2$  agree on  $\mathcal{C}$ . But  $\sigma(\mathcal{C}) = 2^{\{a,b,c,d\}}$ , and  $P_1$  and  $P_2$  do not agree on  $\sigma(\mathcal{C})$ , because, for instance,  $P_1(\{a\}) \neq P_2(\{a\})$ . Note that  $\mathcal{C}$  is not a  $\pi$ -system as  $\{a, b\}\{a, c\} = \{a\} \notin \mathcal{C}$ .

2.11: Using Boole's and D'Morgan's laws,

$$P((\cap_n B_n)^c) = P(\cup_n B_n^c) \leq \sum_n P(B_n^c) = 0,$$

and so  $P(\cap_n B_n) = 1$ .

2.12: If  $B \in \mathcal{C}$ , then  $\mathcal{C}_B = \{B\} \subset \mathcal{C}$  is a countable class with  $B \in \sigma(\mathcal{C}_B)$ , and so  $\mathcal{C} \subset \mathcal{G}$ . And  $\Omega \in \mathcal{G}$  as  $\Omega \in \mathcal{C}_B$  for any  $\mathcal{C}_B$ . If  $B \in \mathcal{G}$  with  $B \in \mathcal{C}_B$  for some countable  $\mathcal{C}_B$ , then  $B^c \in \mathcal{G}$  because  $B^c \in \mathcal{C}_B$ . So  $\mathcal{G}$  is closed under complements. Finally, let  $B_n$ ,  $n \geq 1$ , be sets in  $\mathcal{G}$ . Then there exist countable classes  $\mathcal{C}_{B_n} \subset \mathcal{C}$ ,  $n \geq 1$ , with  $B_n \in \sigma(\mathcal{C}_{B_n})$ . But then  $\mathcal{C}_B \stackrel{\text{def}}{=} \cup_n \mathcal{C}_{B_n} \subset \mathcal{C}$  is a countable class and  $B \stackrel{\text{def}}{=} \cup_n B_n \in \sigma(\mathcal{C}_B)$ , and so  $B \in \mathcal{G}$ . Thus  $\mathcal{G}$  is a  $\sigma$ -field and must contain  $\sigma(\mathcal{C})$ . (In fact,  $\mathcal{G} = \sigma(\mathcal{C})$  as all of the sets of  $\mathcal{G}$  lie in  $\sigma(\mathcal{C})$ .) The desired result follows.