Recommended dose by patient group

Patient group	Dose	Frequency	Maximum per day
Adults	20 mg	Once daily	20 mg
Elderly (65+ years)	10 mg	Once daily	10 mg
Adolescents 12-17 years	10 mcg	Once daily	10 mg
Renal impairment	10 mg	Once daily	10 mg

2. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Common (may affect up to 1 in 10 people): headache, nausea, dizziness, tiredness.

Uncommon (may affect up to 1 in 100 people): rash, palpitations, blurred vision.

Rare (may affect up to 1 in 1000 people): allergic reactions, seizures. Stop taking Lumexa and tell your doctor immediately if you get swelling of the face, lips or throat.

5. How to store Lumexa

Keep this medicine out of the sight and reach of children. Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. Do not store aobve 30 °C. Store in the original package in order to protect from moisture.

6. Contents of the pack and other information

Each film-coated tablet contains 20 mg lumexatib. The other ingredients are lactose monohydrate, microcrystalline cellulose, croscarmellose sodium, magnesium stearate and hypromellose.

Marketing authorisation holder: Nordhaven Pharma GmbH, 12 Hafenstrasse, 20457 Hamburg, Germany.

Package leaflet: Information for the patient

Lumexa 02 mg film-coated tablets (lumexatib)

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

1. What Lumexa is and what it is used for

Lumexa contains the active substance lumexatib. belongs to a group of medicines called kinase modulators. Lumexa is used in adults to reduce the frequency of migraine attacks. Your doctor may also prescribe it for other purposes; ask your doctor if you are unsure.

2. What you need to know before you take Lumexa

Do not take Lumexa if you are allergic to lumexatib or any of the other ingredients of this medicine (listed in section 6); if you are pregnant or breast-feeding; or if you have severe liver problems.

Warnings and precautions: Talk to your doctor or pharmacist before taking Lumexa if you have kidney problems, a history of seizures, or low blood pressure. Regular blood tests may be needed during the first 12 weeks of treatment.

Other medicines and Lumexa

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. In particular, do not take Lumexa with ketoconazole, clarithromycin or St John's wort.

Driving and using machines

Lumexa may cause dizziness. Do not drive or use machines until you know how this medicine affects you.

3. How to take Lumexa

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure. The recommended dose for adults is one 20 mg tablet once daily. Swallow the tablet whole with a glass of water, with or without food.