

Diffusion processes in the extracellular space

Fredrik E. Pettersen

June 16, 2013

Contents

- 1 Introduction
- 2 Mathematical models
- 3 Measurement techniques
- 4 Other possible modeling approaches
- 5 Outlook

Brief introduction to the brain




Cells in the brain

Neurons:


- Signal processing

Structure of a Typical Neuron



Neuroglia:

- Janitorial tasks



The extracellular space

- Space surrounding neurons and neuroglia
- Accounting for $\sim 20\%$ of total brain volume
- Important for transport of nutrients, medicines etc.




Figure: Extracellular space marked as dark grey.

Basic diffusion

The basic diffusion equation reads

$$\frac{\partial C}{\partial t} = D \nabla^2 C \quad (1)$$

Einstein relations

$$D = \frac{k_B T}{6\pi\eta r} \quad (2)$$

$$\langle r^2 \rangle = 2dDt \quad (3)$$

Diffusion in ECS

Network simulations:

- Verification against experimental results
- Local field potential
- Extracellular conductance

$$\sigma = \frac{cq}{k_B T} D \quad (4)$$

$$\nabla \cdot (\sigma(\mathbf{r}) \nabla \phi(\mathbf{r}, t)) = -C(\mathbf{r}, t)$$

Optical measurements

TMA⁺ measurements

Radiotracer measurements

Diffusion Tensor Imaging

Molecular dynamics

Random walks

Random walks

