

Algorithms II Cheat Sheet

Tips

Apply an algorithm you know in a clever way, don't write a new algorithm.

Set notation

$A \in [10] \equiv A \in [1..10]$

Big O

Notation	Intuitive meaning	Analogue
$f(n) \in O(g(n))$	f grows at most as fast as g	\leq
$f(n) \in \Omega(g(n))$	f grows at least as fast as g	\geq
$f(n) \in \Theta(g(n))$	f at the same rate as g	$=$
$f(n) \in o(g(n))$	f grows strictly less fast than g	$<$
$f(n) \in \omega(g(n))$	f grows strictly faster than g	$>$

Notation	Formal definition
$f(n) \in O(g(n))$	$\exists C, n_0: \forall n \geq n_0: f(n) \leq C \cdot g(n)$
$f(n) \in \Omega(g(n))$	$\exists c, n_0: \forall n \geq n_0: f(n) \geq c \cdot g(n)$
$f(n) \in \Theta(g(n))$	$\exists c, C, n_0: \forall n \geq n_0: c \cdot g(n) \leq f(n) \leq C \cdot g(n)$
$f(n) \in o(g(n))$	$\forall C: \exists n_0: \forall n \geq n_0: f(n) \leq C \cdot g(n)$
$f(n) \in \omega(g(n))$	$\forall c: \exists n_0: \forall n \geq n_0: f(n) \geq c \cdot g(n)$

Interval Scheduling

A **request** is a pair of integers (s, f) with $0 \leq s \leq f$.
We call s the **start time** and f the **finish time**.

A set A of requests is **compatible** if for all distinct $(s, f), (s', f') \in A$, either $s' \geq f$ or $s \geq f'$ — that is, the requests' time intervals don't overlap.

Interval Scheduling Problem

Input: An array \mathcal{R} of n requests $(s_1, f_1), \dots, (s_n, f_n)$.

Desired Output: A compatible subset of \mathcal{R} of maximum possible size.

Algorithm: GREEDYSCHEDULE

Input: An array \mathcal{R} of n requests.

Output: A maximum compatible subset of \mathcal{R} .

```

1 begin
2   Sort  $\mathcal{R}$ 's entries so that  $\mathcal{R} \leftarrow [(s_1, f_1), \dots, (s_n, f_n)]$  where  $f_1 \leq \dots \leq f_n$ .
3   Initialise  $A \leftarrow []$ , lastf  $\leftarrow 0$ .
4   foreach  $i \in \{1, \dots, n\}$  do
5     if  $s_i \geq \text{lastf}$  then
6       Append  $(s_i, f_i)$  to  $A$  and update lastf  $\leftarrow f_i$ .
7   Return  $A$ .
```

Complexity:

Step 2 takes $O(n \log n)$

Steps 3–6 all take $O(1)$ time and are executed at most n times.

$\therefore \text{totalrunningtime} = O(n \log n) + O(n)O(1) = O(n \log n)$.

Interval Scheduling

Formal GreedySchedule

$A^+ := \text{argmin} \{f : (s, f) \in R, A \cup \{(s, f)\} \text{ is compatible}\}$ for all $A \subseteq R$,

$A_0 := \emptyset, \quad A_{i+1} := A_i \cup \{A_i^+\}$

$t := \max\{i: A_i \text{ is defined}\}$

Interval Scheduling Proofs

Lemma: Greedy Schedule always outputs A_t

Proof: By induction form the following loop invariant. At the start of the i 'th iteration of 4-7:

- A is equal to $A_t \cap \{(s_1, f_1), \dots, (s_{i-1}, f_{i-1})\}$
- lastf is equal to the latest finish time of any request in A (or 0 if $A = []$)

Lemma: A_t is a compatible set

Proof: Instant by induction; A_0 is compatible, and if A_i is compatible then so is $A_{i+1} = A_i \cup A^+$ by the definition of A_i^+

Lemma: A_t is a maximum compatible subset of the Array R (look in pseudocode)

Proof:

Base case for $i = 1$: A_0^+ is the fastest finishing request in R by definition

Inductive step: Suppose A_i finishes faster than B_i .

Let B_i^+ be the $(i+1)$ 'st fastest-finishing element of B . Since A_i finishes faster than B_i , $A_i \cup \{B_i^+\}$ is compatible. Hence by definition, A_i^+ exists and finishes no later than B_i^+

ODEs

1st Order Linear

Use integrating factor,
 $I = e^{\int P(x)dx}$

Separable:

$\int P(y)dy/dx = \int Q(x)$

HomogEnEous:

$dy/dx = f(x, y) = f(x, y/x)$
sub $y = xV$ solve, then sub
 $V = y/x$

Exact:

If $M(x, y) + N(x, y)dy/dx = 0$ and $M_y = N_x$ i.e.
 $\langle M, N \rangle = \nabla F$ then $\int_x M + \int_y N = F$

Order Reduction

Let $v = dy/dx$ then check other types
If purely a function of y ,
 $\frac{dv}{dx} = v \frac{dv}{dy}$

Variation of Parameters:

When $y'' + a_1 y' + a_2 y = F(x)$
 F contains $\ln x$, $\sec x$, $\tan x$,
 \div

Bernoulli

$y' + P(x)y = Q(x)y^n$
 $\div y^n$
 $y^{-n}y' + P(x)y^{1-n} = Q(x)$
Let $U(x) = y^{1-n}(x)$
 $\frac{dU}{dx} = (1-n)y^{-n} \frac{dy}{dx}$
 $\frac{1}{1-n} \frac{dU}{dx} + P(x)U(x) = Q(x)$
solve as a 1st order

Cauchy-Euler

$x^n y^n + a_1 x^{n-1} y^{n-1} + \dots + a_{n-1} y^{n-2} + a_n y = 0$
guess $y = x^r$

3 Cases:

1) Distinct real roots

$y = ax^{r_1} + bx^{r_2}$

2) Repeated real roots

$y = Ax^r + y_2$
Guess $y_2 = x^r u(x)$
Solve for $u(x)$ and choose one ($A = 1, C = 0$)

3) Distinct complex roots

$y = B_1 x^a \cos(b \ln x) + B_2 x^a \sin(b \ln x)$

Laplace Transforms

$L[f](s) = \int_0^\infty e^{-sx} f(x) dx$

$f(t) = t^n, n \geq 0$

$F(s) = \frac{n!}{s^{n+1}}, s > 0$

$f(t) = e^{at}, a \text{ constant}$

$F(s) = \frac{1}{s-a}, s > a$

$f(t) = \sin bt, b \text{ constant}$

$F(s) = \frac{b}{s^2 + b^2}, s > 0$

$f(t) = \cos bt, b \text{ constant}$

$F(s) = \frac{s}{s^2 + b^2}, s > 0$

$f(t) = t^{-1/2}$

$F(s) = \frac{\pi}{s^{1/2}}, s > 0$

$f(t) = \delta(t - a)$

$F(s) = e^{-as}$

f'

$L[f'] = sL[f] - f(0)$

f''

$L[f''] = s^2 L[f] - sf(0) - f'(0)$

$L[e^{at} f(t)]$

$L[f](s-a)$

Vector Spaces

$$v_1, v_2 \in V$$

$$1. v_1 + v_2 \in V$$

$$2. k \in \mathbb{F}, kv_1 \in V$$

$$3. v_1 + v_2 = v_2 + v_1$$

$$4. (v_1 + v_2) + v_3 = v_1 + (v_2 + v_3)$$

$$5. \forall v \in V, 0 \in V \mid 0 + v_1 = v_1 + 0 = v_1$$

$$6. \forall v \in V, \exists -v \in V \mid v + (-v) = (-v) + v = 0$$

$$7. \forall v \in V, 1 \in \mathbb{F} \mid 1 * v = v$$

$$8. \forall v \in V, k, l \in \mathbb{F}, (kl)v = k(lv)$$

$$9. \forall k \in \mathbb{F}, k(v_1 + v_2) = kv_1 + kv_2$$

$$10. \forall v \in V, k, l \in \mathbb{F}, (k + l)v = kv + lv$$