
Example 1: PL



Each run first gives one overall error across the six parameters:


$$E_N^{(r)} = \|\widehat{\boldsymbol{\theta}}_N^{(r)} - \boldsymbol{\theta}^\star\|_2 = \left[\sum_{j=1}^{6} (\widehat{\theta}_{j,N}^{(r)} - \theta_j^\star)^2\right]^{1/2}.$$

The plotted means and pointwise approximate 95% Monte Carlo confidence intervals across retained runs are

$$\overline{E}_N = \frac{1}{R_N} \sum_{r \in \mathcal{I}_N} E_N^{(r)}, \qquad \overline{E}_N \pm 1.96\, \frac{s_{E,N}}{\sqrt{R_N}},$$

where $s_{E,N}$ is the sample standard deviation of the per-run errors. These intervals describe uncertainty in the mean overall error; they are not intervals for an individual run or simultaneous bands. All six parameters enter on their original scale, with equal weights. Finite stalled and iteration-limited outputs are retained; failed fits are excluded. Linear vertical axes start at zero and are capped at one, with tighter ranges retained. Points and interval portions outside the display range are clipped; boundary triangles mark clipped means.

Example 1: FSM, 128 coefficients



Each run first gives one overall error across the six parameters:


$$E_N^{(r)} = \|\widehat{\boldsymbol{\theta}}_N^{(r)} - \boldsymbol{\theta}^\star\|_2 = \left[\sum_{j=1}^{6} (\widehat{\theta}_{j,N}^{(r)} - \theta_j^\star)^2\right]^{1/2}.$$

The plotted means and pointwise approximate 95% Monte Carlo confidence intervals across retained runs are

$$\overline{E}_N = \frac{1}{R_N} \sum_{r \in \mathcal{I}_N} E_N^{(r)}, \qquad \overline{E}_N \pm 1.96\, \frac{s_{E,N}}{\sqrt{R_N}},$$

where $s_{E,N}$ is the sample standard deviation of the per-run errors. These intervals describe uncertainty in the mean overall error; they are not intervals for an individual run or simultaneous bands. All six parameters enter on their original scale, with equal weights. Finite stalled and iteration-limited outputs are retained; failed fits are excluded. Linear vertical axes start at zero and are capped at one, with tighter ranges retained. Points and interval portions outside the display range are clipped; boundary triangles mark clipped means. For GHAM-2-I, the retained counts are shown in record-length order; the final point summarizes only 7 of 100 runs.

CSTR: PL



Each run first gives one overall error across the six parameters:


$$E_N^{(r)} = \|\widehat{\boldsymbol{\theta}}_N^{(r)} - \boldsymbol{\theta}^\star\|_2 = \left[\sum_{j=1}^{6} (\widehat{\theta}_{j,N}^{(r)} - \theta_j^\star)^2\right]^{1/2}.$$

The plotted means and pointwise approximate 95% Monte Carlo confidence intervals across retained runs are

$$\overline{E}_N = \frac{1}{R_N} \sum_{r \in \mathcal{I}_N} E_N^{(r)}, \qquad \overline{E}_N \pm 1.96\, \frac{s_{E,N}}{\sqrt{R_N}},$$

where $s_{E,N}$ is the sample standard deviation of the per-run errors. These intervals describe uncertainty in the mean overall error; they are not intervals for an individual run or simultaneous bands. All six parameters enter on their original scale, with equal weights. Finite stalled and iteration-limited outputs are retained; failed fits are excluded. Linear vertical axes start at zero and are capped at one, with tighter ranges retained. Points and interval portions outside the display range are clipped; boundary triangles mark clipped means.

CSTR: FSM, 128 coefficients



Each run first gives one overall error across the six parameters:


$$E_N^{(r)} = \|\widehat{\boldsymbol{\theta}}_N^{(r)} - \boldsymbol{\theta}^\star\|_2 = \left[\sum_{j=1}^{6} (\widehat{\theta}_{j,N}^{(r)} - \theta_j^\star)^2\right]^{1/2}.$$

The plotted means and pointwise approximate 95% Monte Carlo confidence intervals across retained runs are

$$\overline{E}_N = \frac{1}{R_N} \sum_{r \in \mathcal{I}_N} E_N^{(r)}, \qquad \overline{E}_N \pm 1.96\, \frac{s_{E,N}}{\sqrt{R_N}},$$

where $s_{E,N}$ is the sample standard deviation of the per-run errors. These intervals describe uncertainty in the mean overall error; they are not intervals for an individual run or simultaneous bands. All six parameters enter on their original scale, with equal weights. Finite stalled and iteration-limited outputs are retained; failed fits are excluded. Linear vertical axes start at zero and are capped at one, with tighter ranges retained. Points and interval portions outside the display range are clipped; boundary triangles mark clipped means.

Example 1: FSM, 256 coefficients



Each run first gives one overall error across the six parameters:


$$E_N^{(r)} = \|\widehat{\boldsymbol{\theta}}_N^{(r)} - \boldsymbol{\theta}^\star\|_2 = \left[\sum_{j=1}^{6} (\widehat{\theta}_{j,N}^{(r)} - \theta_j^\star)^2\right]^{1/2}.$$

The plotted means and pointwise approximate 95% Monte Carlo confidence intervals across retained runs are

$$\overline{E}_N = \frac{1}{R_N} \sum_{r \in \mathcal{I}_N} E_N^{(r)}, \qquad \overline{E}_N \pm 1.96\, \frac{s_{E,N}}{\sqrt{R_N}},$$

where $s_{E,N}$ is the sample standard deviation of the per-run errors. These intervals describe uncertainty in the mean overall error; they are not intervals for an individual run or simultaneous bands. All six parameters enter on their original scale, with equal weights. Finite stalled and iteration-limited outputs are retained; failed fits are excluded. Linear vertical axes start at zero and are capped at one, with tighter ranges retained. Points and interval portions outside the display range are clipped; boundary triangles mark clipped means. For GHAM-2-I, the final point summarizes only 7 of 100 runs.

CSTR: FSM, 256 coefficients



Each run first gives one overall error across the six parameters:

$$E_N^{(r)} = \|\widehat{\boldsymbol{\theta}}_N^{(r)} - \boldsymbol{\theta}^\star\|_2 = \left[\sum_{j=1}^{6} (\widehat{\theta}_{j,N}^{(r)} - \theta_j^\star)^2\right]^{1/2}.$$

The plotted means and pointwise approximate 95% Monte Carlo confidence intervals across retained runs are

$$\overline{E}_N = \frac{1}{R_N} \sum_{r \in \mathcal{I}_N} E_N^{(r)}, \qquad \overline{E}_N \pm 1.96\, \frac{s_{E,N}}{\sqrt{R_N}},$$

where $s_{E,N}$ is the sample standard deviation of the per-run errors. These intervals describe uncertainty in the mean overall error; they are not intervals for an individual run or simultaneous bands. All six parameters enter on their original scale, with equal weights. Finite stalled and iteration-limited outputs are retained; failed fits are excluded. Linear vertical axes start at zero and are capped at one, with tighter ranges retained. Points and interval portions outside the display range are clipped; boundary triangles mark clipped means.