

- [1]. AGA, H., SHIBUYA, T., SUGIMOTO, T., KURIMOTO, M., NAKAJIMA, S. ISOLATION AND IDENTIFICATION OF ANTIMICROBIAL COMPOUNDS IN BRAZILIAN PROPOLIS. BIOSCIENCE BIOTECHNOLOGY AND BIOCHEMISTRY. 1994 doi 10.1271/bbb.58.945
- [2]. ASTUDILLO S., L., AVILA R., F., MORRISON A., R., GUTIERREZ C., M., BASTIDA, J., CODINA, C., et al. BIOLOGICALLY ACTIVE COMPOUNDS FROM CHILEAN PROPOLIS. Boletín de la Sociedad Chilena de Química. 2000
- [3]. Abu-Mellal A, K.N., Duke RK, T.V., CC., D. Prenylated cinnamate and stilbenes from Kangaroo Island propolis and their antioxidant activity. Phytochemistry. 2012 doi 10.1016/j.phytochem.2012.01.012
- [4]. Adelman J, P.M., Breyer DH, S.M., Lenz C, L.N., Lanças FM, F.J. Exotic flora dependence of an unusual Brazilian propolis: the pinocembrin biomarker by capillary techniques. J Pharm Biomed Anal. 2007 doi 10.1016/j.jpba.2006.07.014
- [5]. Agüero MB, G.M., Lima B, S.L., Sánchez M, Z.S., Feresin GE, S.H.G., Palermo J, W.D., A., T. Argentinean propolis from *Zuccagnia punctata* Cav. (Caesalpinieae) exudates: phytochemical characterization and antifungal activity. J Agric Food Chem. 2010 doi 10.1021/jf902991t
- [6]. Ahn MR, K.S., Hamasaka T, B.K., T., N. Antioxidant activity and constituents of propolis collected in various areas of Korea. J Agric Food Chem. 2004 doi 10.1021/jf048726s
- [7]. Ahn, M.R., Kumazawa, S., Usui, Y., Nakamura, J., Matsuka, M., Zhu, F., et al. Antioxidant activity and constituents of propolis collected in various areas of China. FOOD CHEMISTRY. 2007 doi 10.1016/j.foodchem.2006.03.045
- [8]. Akao Y, M.H., Matsumoto K, O.K., Nishizawa K, S.T., Araki Y, M.S., Y., N. Cell growth inhibitory effect of cinnamic acid derivatives from propolis on human tumor cell lines. Biol Pharm Bull. 2003 doi 10.1248/bpb.26.1057
- [9]. Alday E, V.D., Garibay-Escobar A, D.E.Z., Piccinelli AL, R.L., Monribot-Villanueva J, G.A.J., Robles-Zepeda RE, H.J., C., V. Plant origin authentication of Sonoran Desert propolis: an antiproliferative propolis from a semi-arid region. Naturwissenschaften. 2019 doi 10.1007/s00114-019-1620-2
- [10]. Alencar SM, O.T., Castro ML, C.I., Costa-Neto CM, C.J., Rosalen PL, I.M. Chemical composition and biological activity of a new type of Brazilian propolis: red propolis. J Ethnopharmacol. 2007 doi 10.1016/j.jep.2007.06.005
- [11]. Alencar, S.M., Aguiar, C.L., Paredes-Guzmán, J., Park, Y.K. Composição química de *Baccharis dracunculifolia*, fonte botânica das própolis dos estados de São Paulo e Minas Gerais. Ciência Rural. 2005
- [12]. Aliboni, A., D'Andrea, A., Massanisso, P. Treatment of propolis specimens from Central Italy to yield a product with a lower charge of allergenic species. SEPARATION AND PURIFICATION TECHNOLOGY. 2011 doi 10.1016/j.seppur.2011.08.022

- [13]. Aliyazicioglu, R., Sahin, H., Erturk, O., Ulusoy, E., Kolayli, S. Properties of Phenolic Composition and Biological Activity of Propolis from Turkey. *INTERNATIONAL JOURNAL OF FOOD PROPERTIES*. 2013 doi 10.1080/10942912.2010.551312
- [14]. Amoros M, S.C., Girre L, S.F., M., C. Synergistic effect of flavones and flavonols against herpes simplex virus type 1 in cell culture. Comparison with the antiviral activity of propolis. *J Nat Prod*. 1992 doi 10.1021/np50090a003
- [15]. Asgharpour F, M.A., Motalebnejad M, N.H. Propolis attenuates lipopolysaccharide-induced inflammatory responses through intracellular ROS and NO levels along with downregulation of IL-1 $\beta$  and IL-6 expressions in murine RAW 264.7 macrophages. *J Food Biochem*. 2019 doi 10.1111/jfbc.12926
- [16]. Awale S, L.F., Onozuka H, E.H., Tezuka Y, K.S. Constituents of Brazilian red propolis and their preferential cytotoxic activity against human pancreatic PANC-1 cancer cell line in nutrient-deprived condition. *Bioorg Med Chem*. 2008 doi 10.1016/j.bmc.2007.10.004
- [17]. Awale S, S.S., Tezuka Y, U.J., Matsushige K, K.S. Neoflavonoids and related constituents from Nepalese propolis and their nitric oxide production inhibitory activity. *J Nat Prod*. 2005 doi 10.1021/np050009k
- [18]. BANKOVA, V., DYULGEROV, A., POPOV, S., EVSTATIEVA, L., KULEVA, L., PUREB, O., et al. PROPOLIS PRODUCED IN BULGARIA AND MONGOLIA - PHENOLIC-COMPOUNDS AND PLANT-ORIGIN. *APIDOLOGIE*. 1992 doi 10.1051/apido:19920109
- [19]. BONVEHI, J., COLL, F. PHENOLIC COMPOSITION OF PROPOLIS FROM CHINA AND FROM SOUTH-AMERICA. *ZEITSCHRIFT FUR NATURFORSCHUNG C-A JOURNAL OF BIOSCIENCES*. 1994
- [20]. BONVEHI, J., COLL, F., JORDA, R. THE COMPOSITION, ACTIVE COMPONENTS AND BACTERIOSTATIC ACTIVITY OF PROPOLIS IN DIETETICS. *JOURNAL OF THE AMERICAN OIL CHEMISTS SOCIETY*. 1994 doi 10.1007/BF02540666
- [21]. Bankova V, C.R., Kujumgiev A, M.M., S., P. Chemical composition and antibacterial activity of Brazilian propolis. *Z Naturforsch C J Biosci*. 1995 doi 10.1515/znc-1995-3-402
- [22]. Bankova V, P.M., Bogdanov S, S.A. Chemical composition of European propolis: expected and unexpected results. *Z Naturforsch C J Biosci*. 2002 doi 10.1515/znc-2002-5-622
- [23]. Bankova, V., Boudourova-Krasteva, G., Popov, S., Sforcin, J., Funari, S. Seasonal variations of the chemical composition of Brazilian propolis. *APIDOLOGIE*. 1998 doi 10.1051/apido:19980406
- [24]. Bankova, V., Boudourova-Krasteva, G., Sforcin, J., Frete, X., Kujumgiev, A., Maimoni-Rodella, R., et al. Phytochemical evidence for the plant origin of Brazilian propolis from Sao Paulo state. *ZEITSCHRIFT FUR NATURFORSCHUNG C-A JOURNAL OF BIOSCIENCES*. 1999

- [25]. Bankova, V., Christov, R., Marcucci, C., Popov, S. Constituents of Brazilian geopropolis. ZEITSCHRIFT FUR NATURFORSCHUNG SECTION C-A JOURNAL OF BIOSCIENCES. 1998
- [26]. Bankova, V., Nikolova, N., Marcucci, M. New lignan from Brazilian propolis. ZEITSCHRIFT FUR NATURFORSCHUNG C-A JOURNAL OF BIOSCIENCES. 1996 doi 10.1515/znc-1996-9-1019
- [27]. Banskota AH, N.T., Sumioka LY, T.Y., Awale S, M.K., Matsushige K, K.S. Antiproliferative activity of the Netherlands propolis and its active principles in cancer cell lines. J Ethnopharmacol. 2002 doi 10.1016/s0378-8741(02)00022-3
- [28]. Banskota AH, T.Y., Midorikawa K, M.K., S., K. Two novel cytotoxic benzofuran derivatives from Brazilian propolis. J Nat Prod. 2000 doi 10.1021/np000143z
- [29]. Barbosa da Silva Cunha, I., Lucia Tiveron Rodrigues, M., Cesar Meurer, E., Bankova, V.S., Cristina Marcucci, M., Nogueira Eberlin, M., et al. Effect of the maceration time on chemical composition of extracts of Brazilian propolis. JOURNAL OF APICULTURAL RESEARCH. 2006
- [30]. Barrientos L, H.C., Montenegro G, O.X., Veloz J, A.M., Cuevas A, S.N., LA., S. Chemical and botanical characterization of Chilean propolis and biological activity on cariogenic bacteria Streptococcus mutans and Streptococcus sobrinus. Braz J Microbiol. 2013 doi 10.1590/S1517-83822013000200038
- [31]. Barth, O.M., De Freitas, A.S., Matsuda, A.H., De Almeida-Muradian, L.B. Botanical origin and Artepillin-C content of Brazilian propolis samples. GRANA. 2013 doi 10.1080/00173134.2012.747561
- [32]. Barud Hda S, A.J.A., Saska S, M.L., Campos JA, F.R., Ferreira NU, N.A., Miguel FG, V.M., Barizon EA, M.O.F., et al. Antimicrobial Brazilian Propolis (EPP-AF) Containing Biocellulose Membranes as Promising Biomaterial for Skin Wound Healing. Evid Based Complement Alternat Med. 2013 doi 10.1155/2013/703024
- [33]. Bayram, N.E., Gercek, Y.C., Bayram, S., Togar, B. Effects of processing methods and extraction solvents on the chemical content and bioactive properties of propolis. JOURNAL OF FOOD MEASUREMENT AND CHARACTERIZATION. 2020 doi 10.1007/s11694-019-00340-z
- [34]. Bener, M., Ozyurek, M., Guclu, K., Apak, R. Polyphenolic contents of natural dyes produced from industrial plants assayed by HPLC and novel spectrophotometric methods. INDUSTRIAL CROPS AND PRODUCTS. 2010 doi 10.1016/j.indcrop.2010.06.020
- [35]. Benhanifia, M., Mohamed, W.M., Bellik, Y., Benbarek, H. Antimicrobial and antioxidant activities of different propolis samples from north-western Algeria. INTERNATIONAL JOURNAL OF FOOD SCIENCE AND TECHNOLOGY. 2013 doi 10.1111/ijfs.12244

- [36]. Berretta AA, C.P., Cavaleiro AH, F.V., Bom VP, N.A., Marquele-Oliveira F, P.V., Ramalho LN, G.G. Evaluation of Mucoadhesive Gels with Propolis (EPP-AF) in Preclinical Treatment of Candidiasis Vulvovaginal Infection. *Evid Based Complement Alternat Med*. 2013 doi 10.1155/2013/641480
- [37]. Bertelli D, P.G., Bortolotti L, M.G., M., P. <sup>1</sup>H-NMR simultaneous identification of health-relevant compounds in propolis extracts. *Phytochem Anal*. 2012 doi 10.1002/pca.1352
- [38]. Bertrams, J., Mueller, M.B., Kunz, N., Kammerer, D.R., Stintzing, F.C. Phenolic compounds as marker compounds for botanical origin determination of German propolis samples based on TLC and TLC-MS. *JOURNAL OF APPLIED BOTANY AND FOOD QUALITY*. 2013 doi 10.5073/JABFQ.2013.086.020
- [39]. Bilikova, K., Popova, M., Trusheva, B., Bankova, V. New anti-Paenibacillus larvae substances purified from propolis. *APIDOLOGIE*. 2013 doi 10.1007/s13592-012-0178-1
- [40]. Biscaia, D., Ferreira, S.R.S. Propolis extracts obtained by low pressure methods and supercritical fluid extraction. *JOURNAL OF SUPERCRITICAL FLUIDS*. 2009 doi 10.1016/j.supflu.2009.07.011
- [41]. Biu, C., Li-Ping, L., Yan, L., Yuan-Jun, X., Yu-Xin, F., Yin-Yu, G. Analyses of Phenolic Compounds of Water Extract from Propolis by Reversed Phase High Performance Liquid Chromatography. *CHINESE JOURNAL OF ANALYTICAL CHEMISTRY*. 2009
- [42]. Bolfa P, V.R., Petruta A, D.D., Stan L, V.L., Damian G, C.C., Filip A, C.S. Photoprotective effects of Romanian propolis on skin of mice exposed to UVB irradiation. *Food Chem Toxicol*. 2013 doi 10.1016/j.fct.2013.08.078
- [43]. Bouaroura A, S.N., Diaz JG, B.C., Akkal S, R.S. Preliminary analysis of the chemical composition, antioxidant and anticholinesterase activities of Algerian propolis. *Nat Prod Res*. 2019 doi 10.1080/14786419.2018.1556658
- [44]. Buchta, V., Cerny, J., Opletalova, V. In vitro antifungal activity of propolis samples of Czech and Slovak origin. *CENTRAL EUROPEAN JOURNAL OF BIOLOGY*. 2011 doi 10.2478/s11535-010-0112-3
- [45]. Bueno-Silva B, K.H., Falsetta ML, A.S., Ikegaki M, R.P. Effect of neovestitol-vestitol containing Brazilian red propolis on accumulation of biofilm in vitro and development of dental caries in vivo. *Biofouling*. 2013 doi 10.1080/08927014.2013.834050
- [46]. Buratti S, B.S., MS., C. Evaluation of the antioxidant power of honey, propolis and royal jelly by amperometric flow injection analysis. *Talanta*. 2007 doi 10.1016/j.talanta.2006.07.006
- [47]. Bursal, E., Gulcin, I. Polyphenol contents and in vitro antioxidant activities of lyophilised aqueous extract of kiwifruit (*Actinidia deliciosa*). *FOOD RESEARCH INTERNATIONAL*. 2011 doi 10.1016/j.foodres.2011.03.031

- [48]. Búfalo MC, F.I., Costa G, F.V., Liberal J, C.M., Lopes MC, B.M., JM., S. Propolis and its constituent caffeic acid suppress LPS-stimulated pro-inflammatory response by blocking NF- $\kappa$ B and MAPK activation in macrophages. *J Ethnopharmacol.* 2013 doi 10.1016/j.jep.2013.06.004
- [49]. CHI, H., HSIEH, A., NG, C., LEE, H., LI, S. DETERMINATION OF COMPONENTS IN PROPOLIS BY CAPILLARY ELECTROPHORESIS AND PHOTODIODE-ARRAY DETECTION. *JOURNAL OF CHROMATOGRAPHY A.* 1994 doi 10.1016/0021-9673(94)85159-X
- [50]. CHRISTOV, R., BANKOVA, V. GAS-CHROMATOGRAPHIC ANALYSIS OF UNDERIVATIZED PHENOLIC CONSTITUENTS FROM PROPOLIS USING AN ELECTRON-CAPTURE DETECTOR. *JOURNAL OF CHROMATOGRAPHY.* 1992 doi 10.1016/0021-9673(92)85316-L
- [51]. Cabral, I.S.R., Oldoni, T.L.C., Prado, A., Bezerra, R.M.N., Alencar, S.M., Ikegaki, M., et al. Composição fenólica, atividade antibacteriana e antioxidante da própolis vermelha brasileira. *Química Nova.* 2009
- [52]. Cai W, X.J., Li G, L.T., Guo X, W.H., L., L. Ethanol extract of propolis prevents high-fat diet-induced insulin resistance and obesity in association with modulation of gut microbiota in mice. *Food Res Int.* 2020 doi 10.1016/j.foodres.2019.108939
- [53]. Campo Fernández M, C.R.O., Rosado Perez A, M.O.P.R., Márquez Hernández I, P.A., L., R. GC-MS determination of isoflavonoids in seven red Cuban propolis samples. *J Agric Food Chem.* 2008 doi 10.1021/jf801870f
- [54]. Canogullari, S., Baylan, M., Sahinler, N., Sahin, A. Effects of propolis and pollen supplementations on growth performance and body components of Japanese quails (*Coturnix coturnix japonica*). *ARCHIV FUR GEFLUGELKUNDE.* 2009
- [55]. Cao, W., Liu, H.F., Cheng, N., Gao, H., Wang, B.N., Zheng, J.B. LC with Electrochemical Detection for Analysis of Caffeic Acid and Caffeic Acid Phenyl Ester in Propolis. *CHROMATOGRAPHIA.* 2011 doi 10.1007/s10337-010-1864-7
- [56]. Carvalho, A.A., Finger, D., Machado, C.S., Schmidt, E.M., da Costa, P.M., Negreiros Nunes Alves, A.P., et al. In vivo antitumoural activity and composition of an oil extract of Brazilian propolis. *FOOD CHEMISTRY.* 2011 doi 10.1016/j.foodchem.2010.12.035
- [57]. Cassina Lopez, B.G., Bataglion, G.A., Paz Jara, J.L., Pacheco Filho, E.F., de Mendonca Melo, L.S., Cardoso, J.C., et al. Variation of the phytoestrogen composition of red propolis throughout the year. *JOURNAL OF APICULTURAL RESEARCH.* 2020 doi 10.1080/00218839.2020.1714192
- [58]. Castro ML, V.W., Zauli RC, I.M., Rehder VL, F.M., de Alencar SM, R.P. Bioassay guided purification of the antimicrobial fraction of a Brazilian propolis from Bahia state. *BMC Complement Altern Med.* 2009 doi 10.1186/1472-6882-9-25

- [59]. Castro, M.L., Cury, J.A., Rosalen, P.L., Alencar, S.M., Ikegaki, M., Duarte, S., et al. Propolis from southeastern and northeastern of Brazil: The influence of seasonality in antibacterial activity and phenolic composition. *QUIMICA NOVA*. 2007 doi 10.1590/S0100-40422007000700003
- [60]. Castro, M.L., Cury, J.A., Rosalen, P.L., Alencar, S.M., Ikegaki, M., Duarte, S., et al. Própolis do sudeste e nordeste do Brasil: influência da sazonalidade na atividade antibacteriana e composição fenólica. *Química Nova*. 2007
- [61]. Catchpole, O., Grey, J., Mitchell, K., Lan, J. Supercritical antisolvent fractionation of propolis tincture. *JOURNAL OF SUPERCRITICAL FLUIDS*. 2004 doi 10.1016/S0896-8446(03)00033-0
- [62]. Celemlí, O.G., Hatjina, F., Charistos, L., Schiesser, A., Ozkirim, A. More Insight into the Chemical Composition of Greek Propolis; Differences and Similarities with Turkish Propolis. *ZEITSCHRIFT FÜR NATURFORSCHUNG SECTION C-A JOURNAL OF BIOSCIENCES*. 2013 doi 10.5560/ZNC.2013.68c0429
- [63]. Chaillou, L.L., Nazareno, M.A. Bioactivity of propolis from Santiago del Estero, Argentina, related to their chemical composition. *LWT-FOOD SCIENCE AND TECHNOLOGY*. 2009 doi 10.1016/j.lwt.2009.03.002
- [64]. Chaillou, L.L., Nazareno, M.A. Chemical variability in propolis from Santiago del Estero, Argentina, related to the arboreal environment as the sources of resins. *JOURNAL OF THE SCIENCE OF FOOD AND AGRICULTURE*. 2009 doi 10.1002/jsfa.3542
- [65]. Chang, R., Pilo-Veloso, D., Morais, S.A.L., Nascimento, E.A. Analysis of a Brazilian green propolis from *Baccharis dracunculifolia* by HPLC-APCI-MS and GC-MS. *REVISTA BRASILEIRA DE FARMACOGNOSIA-BRAZILIAN JOURNAL OF PHARMACOGNOSY*. 2008 doi 10.1590/S0102-695X2008000400009
- [66]. Chen CN, H.C., Lee SS, G.J., Chiang PC, H.C., WJ., H. Chemical modification and anticancer effect of prenylated flavanones from Taiwanese propolis. *Nat Prod Res*. 2012 doi 10.1080/14786419.2010.535146
- [67]. Chen CN, W.C., Shy HS, L.J. Cytotoxic prenylflavanones from Taiwanese propolis. *J Nat Prod*. 2003 doi 10.1021/np0203180
- [68]. Chen CN, W.M., Wu CL, L.J. Comparison of Radical Scavenging Activity, Cytotoxic Effects and Apoptosis Induction in Human Melanoma Cells by Taiwanese Propolis from Different Sources. *Evid Based Complement Alternat Med*. 2004 doi 10.1093/ecam/neh034
- [69]. Chen F, C.L., Wang Q, Z.J., Xue X, Z.J. Determination of organochlorine pesticides in propolis by gas chromatography-electron capture detection using double column series solid-phase extraction. *Anal Bioanal Chem*. 2009 doi 10.1007/s00216-008-2474-1

- [70]. Chen J, L.Y., Han M, W.T., Chen Q, W.R. Water-soluble derivative of propolis mitigates scopolamine-induced learning and memory impairment in mice. *Pharmacol Biochem Behav.* 2008 doi 10.1016/j.pbb.2008.03.029
- [71]. Chen W, T.X., Wu D, G.Z., Wu S, H.S. Comparison of the Partition Efficiencies of Multiple Phenolic Compounds Contained in Propolis in Different Modes of Acetonitrile-Water-Based Homogenous Liquid-Liquid Extraction. *Molecules.* 2019 doi 10.3390/molecules24030442
- [72]. Chen, C.R., Lee, Y.N., Chang, C.M.J., Lee, M.R., Wei, I.C. Hot-pressurized fluid extraction of flavonoids and phenolic acids from Brazilian propolis and their cytotoxic assay in vitro. *JOURNAL OF THE CHINESE INSTITUTE OF CHEMICAL ENGINEERS.* 2007 doi 10.1016/j.jcice.2007.04.004
- [73]. Chen, Y.W., Wu, S.W., Ho, K.K., Lin, S.B., Huang, C.Y., Chen, C.N. Characterisation of Taiwanese propolis collected from different locations and seasons. *JOURNAL OF THE SCIENCE OF FOOD AND AGRICULTURE.* 2008 doi 10.1002/jsfa.3101
- [74]. Cheng, H., Qin, Z.H., Guo, X.F., Hu, X.S., Wu, J.H. Geographical origin identification of propolis using GC-MS and electronic nose combined with principal component analysis. *FOOD RESEARCH INTERNATIONAL.* 2013 doi 10.1016/j.foodres.2013.01.053
- [75]. Chernetsova ES, B.M., Scheibner O, M.G. DART-Orbitrap MS: a novel mass spectrometric approach for the identification of phenolic compounds in propolis. *Anal Bioanal Chem.* 2012 doi 10.1007/s00216-012-5800-6
- [76]. Chi Y, L.L., Cui M, H.Y., Liu T, H.X., X., G. Chemical Composition and Antioxidant Activity of Essential Oil of Chinese Propolis. *Chem Biodivers.* 2020 doi 10.1002/cbdv.201900489
- [77]. Choi, S.J., Shimomura, K., Kumazawa, S., Ahn, M.R. Antioxidant Properties and Phenolic Composition of Propolis from Diverse Geographic Regions in Korea. *FOOD SCIENCE AND TECHNOLOGY RESEARCH.* 2013 doi 10.3136/fstr.19.211
- [78]. Choudhari MK, P.S., Ranade RV, P.K. Antimicrobial activity of stingless bee (*Trigona* sp.) propolis used in the folk medicine of Western Maharashtra, India. *J Ethnopharmacol.* 2012 doi 10.1016/j.jep.2012.02.047
- [79]. Christov R, T.B., Popova M, B.V., M., B. Chemical composition of propolis from Canada, its antiradical activity and plant origin. *Nat Prod Res.* 2005 doi 10.1080/14786410512331328159
- [80]. Christov R, T.B., Popova M, B.V., M., B. Chemical composition of propolis from Canada, its antiradical activity and plant origin. *Nat Prod Res.* 2006 doi 10.1080/14786410500056918
- [81]. Christov, R., Bankova, V., Hegazi, A., Abd El Hady, F., Popov, S. Chemical composition of Egyptian propolis. *ZEITSCHRIFT FÜR NATURFORSCHUNG C-A JOURNAL OF BIOSCIENCES.* 1998

- [82]. Cigut T, P.T., Gašperlin L, R.P., P., J. Antioxidative activity of propolis extract in yeast cells. *J Agric Food Chem.* 2011 doi 10.1021/jf2022258
- [83]. Cole N, S.P., Ngo A, T.K., Severino JA, A.S., Duke CC, R.V. Topical 'Sydney' propolis protects against UV-radiation-induced inflammation, lipid peroxidation and immune suppression in mouse skin. *Int Arch Allergy Immunol.* 2010 doi 10.1159/000265530
- [84]. Conti BJ, B.M., Golim Mde A, B.V., JM., S. Cinnamic Acid is partially involved in propolis immunomodulatory action on human monocytes. *Evid Based Complement Alternat Med.* 2013 doi 10.1155/2013/109864
- [85]. Corrêa JL, V.F., Jarros IC, C.M., Castilho PF, O.K., Rosseto HC, B.M., Svidzinski TIE, N.M. Propolis extract has bioactivity on the wall and cell membrane of *Candida albicans*. *J Ethnopharmacol.* 2020 doi 10.1016/j.jep.2020.112791
- [86]. Costa AG Jr, Y.N., Garcez WS, P.R., Matos MFC, G.F. Metabolomics Approach Expands the Classification of Propolis Samples from Midwest Brazil. *J Nat Prod.* 2020 doi 10.1021/acs.jnatprod.9b00783
- [87]. Costa da Silva, E.C., Muniz, M.P., Saraiva Nunomura, R.C., Nunomura, S.M., Carvalho Zilse, G.A. PHENOLIC CONSTITUENTS AND ANTIOXIDANT ACTIVITY OF GEOPROPOLIS FROM TWO SPECIES OF AMAZONIAN STINGLESS BEES. *QUIMICA NOVA.* 2013
- [88]. Cuesta-Rubio O, F.U.B., Ramírez-Apan T, C.J. Polyisoprenylated benzophenones in cuban propolis; biological activity of nemorosone. *Z Naturforsch C J Biosci.* 2002 doi 10.1515/znc-2002-3-429
- [89]. Cuesta-Rubio O, P.A., Fernandez MC, H.I., Rosado A, R.L. Chemical characterization of Cuban propolis by HPLC-PDA, HPLC-MS, and NMR: the brown, red, and yellow Cuban varieties of propolis. *J Agric Food Chem.* 2007 doi 10.1021/jf071296w
- [90]. Cueto, A.P., Alves, S.H., Pilau, M., Weiblen, R., Kubica, T.F., Lovato, L.T. Antiviral activity of propolis extracts against feline calicivirus, canine adenovirus 2, and bovine viral diarrhea virus. *CIENCIA RURAL.* 2011 doi 10.1590/S0103-84782011001000021
- [91]. Curti V, Z.V., Tsetegho Sokeng AJ, D.M., Masiello I, M.A., D'Antona G, D.M. Bioavailability and In Vivo Antioxidant Activity of a Standardized Polyphenol Mixture Extracted from Brown Propolis. *Int J Mol Sci.* 2019 doi 10.3390/ijms20051250
- [92]. Daikh, A., Segueni, N., Dogan, N.M., Arslan, S., Mutlu, D., Kivrak, I., et al. Comparative study of antibiofilm, cytotoxic activity and chemical composition of Algerian propolis. *JOURNAL OF APICULTURAL RESEARCH.* 2020 doi 10.1080/00218839.2019.1701777
- [93]. Daleprane JB, F.V.S., Pacheco A, R.M., Faine LA, D.F., Ikegaki M, S., Ong TP, A.D. Anti-atherogenic and anti-angiogenic activities of polyphenols from propolis. *J Nutr Biochem.* 2012 doi 10.1016/j.jnutbio.2011.02.012



- [94]. Dario Pierini, G., Marcelo Granero, A., Susana Di Nezio, M., Eugenia Centurion, M., Alicia Zon, M., Fernandez, H. Development of an electroanalytical method for the determination of lead in Argentina raw propolis based on bismuth electrodes. MICROCHEMICAL JOURNAL. 2013 doi 10.1016/j.microc.2012.05.015
- [95]. Darwish, R.M., Abu Fares, R.J., Abu Zarga, M.H., Nazer, I.K. Antibacterial effect of Jordanian propolis and isolated flavonoids against human pathogenic bacteria. AFRICAN JOURNAL OF BIOTECHNOLOGY. 2010
- [96]. Dausch A, M.C., Fort P, P.Y. Brazilian red propolis--chemical composition and botanical origin. Evid Based Complement Alternat Med. 2008 doi 10.1093/ecam/nem057
- [97]. Denli, M., Cankaya, S., Silici, S., Okan, F., Uluocak, A. Effect of dietary addition of Turkish propolis on the growth performance, carcass characteristics and serum variables of quail (*Coturnix coturnix japonica*). ASIAN-AUSTRALASIAN JOURNAL OF ANIMAL SCIENCES. 2005 doi 10.5713/ajas.2005.848
- [98]. Duarte S, R.P., Hayacibara MF, C.J., Bowen WH, M.R., Rehder VL, S.A., Ikegaki M, K.H. The influence of a novel propolis on mutans streptococci biofilms and caries development in rats. Arch Oral Biol. 2006 doi 10.1016/j.archoralbio.2005.06.002
- [99]. Díaz-Carballo D, M.S., Bardenheuer W, F.M., H., P.R. The contribution of plukenetione A to the anti-tumoral activity of Cuban propolis. Bioorg Med Chem. 2008 doi 10.1016/j.bmc.2008.10.019
- [100]. Díaz-Carballo D, U.K., Kleff V, E.S., Malak S, F.M., Somogyi S, K.C., Bardenheuer W, S.D. Antiretroviral activity of two polyisoprenylated acylphloroglucinols, 7-epi-nemorosone and plukenetione A, isolated from Caribbean propolis. Int J Clin Pharmacol Ther. 2010 doi 10.5414/cpp48670
- [101]. Eda, M., Hayashi, Y., Kinoshita, K., Koyama, K., Takahashi, K., Akutu, K. Anti-emetic principles of water extract of Brazilian propolis. PHARMACEUTICAL BIOLOGY. 2005 doi 10.1080/13880200590919546
- [102]. El-Bassuony A, A.S. A new prenylated flavanoid with antibacterial activity from propolis collected in Egypt. Nat Prod Commun. 2010
- [103]. Eraslan G, K.M., Silici S, A.S., M., K. Effects of cypermethrin on some biochemical changes in rats: the protective role of propolis. Exp Anim. 2008 doi 10.1538/expanim.57.453
- [104]. Eraslan, G., Kanbur, M., Silici, S. Evaluation of propolis effects on some biochemical parameters in rats treated with sodium fluoride. PESTICIDE BIOCHEMISTRY AND PHYSIOLOGY. 2007 doi 10.1016/j.pestbp.2007.01.002
- [105]. FERRERES, F., ORTIZ, A., SILVA, C., GARCIAVIGUERA, C., TOMASBARBERAN, F., TOMASLORENTE, F. FLAVONOIDS OF LA ALCARRIA HONEY - A STUDY OF THEIR BOTANICAL ORIGIN. ZEITSCHRIFT FUR LEBENSMITTEL-UNTERSUCHUNG UND-FORSCHUNG. 1992 doi 10.1007/BF01190185

- [106]. Falcao, S.I., Tomas, A., Vale, N., Gomes, P., Freire, C., Vilas-Boas, M. Phenolic quantification and botanical origin of Portuguese propolis. *INDUSTRIAL CROPS AND PRODUCTS*. 2013 doi 10.1016/j.indcrop.2013.07.021
- [107]. Falcão SI, C.R., Touzani S, L.B., Ferreira ICFR, V.B.M. In Vitro Interactions of Moroccan Propolis Phytochemical's on Human Tumor Cell Lines and Anti-Inflammatory Properties. *Biomolecules*. 2019 doi 10.3390/biom9080315
- [108]. Falcão SI, V.B.M., Estevinho LM, B.C., Domingues MR, C.S. Phenolic characterization of Northeast Portuguese propolis: usual and unusual compounds. *Anal Bioanal Chem*. 2010 doi 10.1007/s00216-009-3232-8
- [109]. Fernandes-Silva, C.C., Salatino, A., Salatino, M.L.F., Breyer, E.D.H., Negri, G. CHEMICAL PROFILING OF SIX SAMPLES OF BRAZILIAN PROPOLIS. *QUIMICA NOVA*. 2013 doi 10.1590/S0100-40422013000200006
- [110]. Ferreira FB, T.S., Rosa OP, F.C., Garcia RB, M.M., BP., G. Antimicrobial effect of propolis and other substances against selected endodontic pathogens. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2007 doi 10.1016/j.tripleo.2007.05.019
- [111]. Finger, D., Machado, C.S., Torres, Y.R., Quinaia, S.P., Godot Thomaz, A.C., Gobbo, A.R., et al. ANTIFUNGAL BIOASSAY-GUIDED FRACTIONATION OF AN OIL EXTRACT OF PROPOLIS. *JOURNAL OF FOOD QUALITY*. 2013 doi 10.1111/jfq.12039
- [112]. Fischer G, P.N., Marcucci MC, S.B., Munhoz LS, F.P., Vargas GD, H.S., Vidor T, R.P. Green propolis phenolic compounds act as vaccine adjuvants, improving humoral and cellular responses in mice inoculated with inactivated vaccines. *Mem Inst Oswaldo Cruz*. 2010 doi 10.1590/s0074-02762010000700012
- [113]. Fontana, J., Passos, M., dos Santos, M., Fontana, C., Oliveira, B., Schause, L., et al. Profiling propolis flavonoids by means of micellar electrokinetic capillary chromatography, capillary gas chromatography and bactericidal action. *CHROMATOGRAPHIA*. 2000 doi 10.1007/BF02490445
- [114]. Freitas AS, C.A., Cardoso SM, O.R., C., A.A. Constancy of the bioactivities of propolis samples collected on the same apiary over four years. *Food Res Int*. 2019 doi 10.1016/j.foodres.2018.10.040
- [115]. Freitas, M.O., Ponte, F.A.E., Lima, M.A.S., Silveira, E.R. Flavonoids and triterpenes from the nest of the stingless bee *Trigona spinipes*. *JOURNAL OF THE BRAZILIAN CHEMICAL SOCIETY*. 2008
- [116]. Frión-Herrera Y, G.D., Díaz-García A, C.R.O., M., C. Chemosensitizing activity of Cuban propolis and nemorosone in doxorubicin resistant human colon carcinoma cells. *Fitoterapia*. 2019 doi 10.1016/j.fitote.2019.104173

- [117]. Furukawa S, T.N., Ikeda T, O.M., Nafady AM, N.T., Sugimoto H, D.S., H., Y. Two novel long-chain alkanolic acid esters of lupeol from alecrim-propolis. *Chem Pharm Bull (Tokyo)*. 2002 doi 10.1248/cpb.50.439
- [118]. GARCIAVIGUERA, C., GREENAWAY, W., WHATLEY, F. COMPOSITION OF PROPOLIS FROM 2 DIFFERENT SPANISH REGIONS. *ZEITSCHRIFT FUR NATURFORSCHUNG C-A JOURNAL OF BIOSCIENCES*. 1992 doi 10.1515/znc-1992-7-824
- [119]. GIL, M., FERRERES, F., ORTIZ, A., SUBRA, E., TOMASBARBERAN, F. PLANT PHENOLIC METABOLITES AND FLORAL ORIGIN OF ROSEMARY HONEY. *JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY*. 1995 doi 10.1021/jf00059a012
- [120]. GREENAWAY, W., MAY, J., SCAYSBROOK, T., WHATLEY, F. IDENTIFICATION BY GAS-CHROMATOGRAPHY MASS-SPECTROMETRY OF 150 COMPOUNDS IN PROPOLIS. *ZEITSCHRIFT FUR NATURFORSCHUNG C-A JOURNAL OF BIOSCIENCES*. 1991
- [121]. GREENAWAY, W., SCAYSBROOK, T., WHATLEY, F. COMPOSITION OF PROPOLIS IN OXFORDSHIRE, UK AND ITS RELATION TO POPLAR BUD EXUDATE. *ZEITSCHRIFT FUR NATURFORSCHUNG C-A JOURNAL OF BIOSCIENCES*. 1988 doi 10.1515/znc-1988-3-423
- [122]. Gabrys J, K.J., Krol W, S.S., J., S. Free amino acids in bee hive product (propolis) as identified and quantified by gas-liquid chromatography. *Pharmacol Res Commun*. 1986 doi 10.1016/0031-6989(86)90146-3
- [123]. Gardana C, B.A., Simonetti P, G.S. Biotransformation strategy to reduce allergens in propolis. *Appl Environ Microbiol*. 2012 doi 10.1128/AEM.00811-12
- [124]. Gardana C, S.M., Pietta P, S.P. Analysis of the polyphenolic fraction of propolis from different sources by liquid chromatography-tandem mass spectrometry. *J Pharm Biomed Anal*. 2007 doi 10.1016/j.jpba.2007.06.022
- [125]. Gardana C, S.P., Berti C, P.P. Evaluation of propolis polyphenols absorption in humans by liquid chromatography/tandem mass spectrometry. *Rapid Commun Mass Spectrom*. 2007 doi 10.1002/rcm.3285
- [126]. González M, G.M., Slanis A, B.A., Fiedler S, F.L., Dellacassa E, C.C., Lorenzo D, R.M., ML., T. Phytochemical Findings Evidencing Botanical Origin of New Propolis Type from North-West Argentina. *Chem Biodivers*. 2019 doi 10.1002/cbdv.201800442
- [127]. Graikou, K., Chinou, I., Gortzi, O., Lalas, S. Chemical composition - biological activities of selected samples of propolis from South Greece. *PLANTA MEDICA*. 2010
- [128]. Graikou, K., Kapeta, S., Aligiannis, N., Sotiroidis, G., Chondrogianni, N., Gonos, E., et al. Chemical analysis of Greek pollen - Antioxidant, antimicrobial and proteasome activation properties. *CHEMISTRY CENTRAL JOURNAL*. 2011 doi 10.1186/1752-153X-5-33

- [129]. Grecka K, K.P., Okińczyc P, W.R., Walkusz J, S.P. The Anti-Staphylococcal Potential of Ethanolic Polish Propolis Extracts. *Molecules*. 2019 doi 10.3390/molecules24091732
- [130]. Gregoris E, S.R. Correlations between polyphenolic composition and antioxidant activity of Venetian propolis. *Food Chem Toxicol*. 2010 doi 10.1016/j.fct.2009.09.018
- [131]. Grunberger D, B.R., Eisinger K, O.E., Efros L, C.M., Estevez V, N.K. Preferential cytotoxicity on tumor cells by caffeic acid phenethyl ester isolated from propolis. *Experientia*. 1988 doi 10.1007/bf01941717
- [132]. Guo X, C.B., Luo L, Z.X., Dai X, G.S. Chemical compositions and antioxidant activities of water extracts of Chinese propolis. *J Agric Food Chem*. 2011 doi 10.1021/jf202818p
- [133]. Gómez-Romero M, A.R.D., Moreno-Torres R, G.S.P., Segura-Carretero A, F.G.A. Antioxidant compounds of propolis determined by capillary electrophoresis-mass spectrometry. *J Sep Sci*. 2007 doi 10.1002/jssc.200600354
- [134]. Gülçin I, B.E., Sehitoglu MH, B.M., AC., G. Polyphenol contents and antioxidant activity of lyophilized aqueous extract of propolis from Erzurum, Turkey. *Food Chem Toxicol*. 2010 doi 10.1016/j.fct.2010.05.053
- [135]. Hamasaka, T., Kumazawa, S., Fujimoto, T., Nakayama, T. Antioxidant activity and constituents of propolis collected in various areas of Japan. *FOOD SCIENCE AND TECHNOLOGY RESEARCH*. 2004 doi 10.3136/fstr.10.86
- [136]. Han, M.S., Lee, I.K., Kim, Y.S., Kim, J.T., Choe, K.R., Yun, B.S. Flavonoids from Propolis Inhibit DNA Single Strand Breakage by the Fenton Reaction. *JOURNAL OF THE KOREAN SOCIETY FOR APPLIED BIOLOGICAL CHEMISTRY*. 2010 doi 10.3839/jksabc.2010.078
- [137]. Hashem, N.M., Abd El-Hady, A., Hassan, O. Effect of vitamin E or propolis supplementation on semen quality, oxidative status and hemato-biochemical changes of rabbit bucks during hot season. *LIVESTOCK SCIENCE*. 2013 doi 10.1016/j.livsci.2013.09.003
- [138]. Hattori H, O.K., Murase T, S.Y., Narise K, S.G., H., N. Isolation, identification, and biological evaluation of HIF-1-modulating compounds from Brazilian green propolis. *Bioorg Med Chem*. 2011 doi 10.1016/j.bmc.2011.07.060
- [139]. Hegazi AG, A.E.H.F. Egyptian propolis: 3. Antioxidant, antimicrobial activities and chemical composition of propolis from reclaimed lands. *Z Naturforsch C J Biosci*. 2002 doi 10.1515/znc-2002-3-432
- [140]. Hernandez J, G.F., Quintero J, A.A., Castañeda M, D.Z., Robles R, V.M.L., Velazquez EF, A.H., Lugo E, V.C. Sonoran propolis: chemical composition and antiproliferative activity on cancer cell lines. *Planta Med*. 2007 doi 10.1055/s-2007-990244
- [141]. Hernández IM, F.M., Cuesta-Rubio O, P.A., L., R. Polyprenylated benzophenone derivatives from Cuban propolis. *J Nat Prod*. 2005 doi 10.1021/np0495884

- [142]. Herrera, C.L., Alvear, M., Barrientos, L., Montenegro, G., Salazar, L.A. The antifungal effect of six commercial extracts of Chilean propolis on *Candida* spp. *Ciencia e investigación agraria*. 2010
- [143]. Hilhorst, M., Somsen, G., de Jong, G. Potential of capillary electrophoresis for the profiling of propolis. *HRC-JOURNAL OF HIGH RESOLUTION CHROMATOGRAPHY*. 1998
- [144]. Hirota M, M.T., Fujiwara T, S.H., S., M. Enhanced cytotoxicity in a Z-photoisomer of a benzopyran derivative of propolis. *J Nat Prod*. 2000 doi 10.1021/np990463m
- [145]. Hrobonova, K., Lehotay, J., Cizmarik, J. Determination of Organic Acids in Propolis by HPLC Using Two Columns with an On-Line SPE System. *JOURNAL OF LIQUID CHROMATOGRAPHY & RELATED TECHNOLOGIES*. 2009 doi 10.1080/10826070802548721
- [146]. Hrobonova, K., Lehotay, J., Cizmarik, J., Sadecka, J. COMPARISON HPLC AND FLUORESCENCE SPECTROMETRY METHODS FOR DETERMINATION OF COUMARIN DERIVATIVES IN PROPOLIS. *JOURNAL OF LIQUID CHROMATOGRAPHY & RELATED TECHNOLOGIES*. 2013 doi 10.1080/10826076.2012.660724
- [147]. Ichi I, H.H., Takashima Y, A.N., Kataoka R, O.K., Hashimoto K, K.S. The beneficial effect of propolis on fat accumulation and lipid metabolism in rats fed a high-fat diet. *J Food Sci*. 2009 doi 10.1111/j.1750-3841.2009.01147.x
- [148]. Inouye S, T.M., S., A. Composition, antifungal and radical scavenging activities of 4 propolis. *Med Mycol J*. 2011 doi 10.3314/mmj.52.305
- [149]. Inui S, H.T., Shimamura Y, M.S., Ogawa T, K.H., Shirafuji K, M.R., Kozone I, S.Y.K., S., K. Solophenols B-D and solomonin: new prenylated polyphenols isolated from propolis collected from the Solomon Islands and their antibacterial activity. *J Agric Food Chem*. 2012 doi 10.1021/jf303516w
- [150]. Inui S, S.Y., Masuda S, S.K., Moli RT, K.S. A new prenylflavonoid isolated from propolis collected in the Solomon Islands. *Biosci Biotechnol Biochem*. 2012 doi 10.1271/bbb.120021
- [151]. Ioshida, M.M., Young, M.C.M., Lago, J.H.G. Chemical Composition and Antifungal Activity of Essential Oil from Brazilian Propolis. *JOURNAL OF ESSENTIAL OIL BEARING PLANTS*. 2010 doi 10.1080/0972060X.2010.10643873
- [152]. Isla MI, P.G.J., Nieva-Moreno MI, K.H., YK., P. Some chemical composition and biological activity of northern Argentine propolis. *J Agric Food Chem*. 2005 doi 10.1021/jf040130h
- [153]. Ito J, C.F., Wang HK, P.Y., Ikegaki M, K.N., KH., L. Anti-AIDS agents. 48.(1) Anti-HIV activity of moronic acid derivatives and the new melliferone-related triterpenoid isolated from Brazilian propolis. *J Nat Prod*. 2001 doi 10.1021/np010211x

- [154]. Ivanauskas L, J.V., Radusiene J, L.A., A., B. Evaluation of phenolic acids and phenylpropanoids in the crude drugs. *Medicina (Kaunas)*. 2008
- [155]. Izuta H, N.Y., Shimazawa M, M.S., Kondo S, H.H. 1,1-diphenyl-2-picrylhydrazyl radical scavenging activity of bee products and their constituents determined by ESR. *Biol Pharm Bull*. 2009 doi 10.1248/bpb.32.1947
- [156]. Izuta H, S.M., Tazawa S, A.Y., Mishima S, H.H. Protective effects of Chinese propolis and its component, chrysin, against neuronal cell death via inhibition of mitochondrial apoptosis pathway in SH-SY5Y cells. *J Agric Food Chem*. 2008 doi 10.1021/jf8014206
- [157]. Jasprica I, B.M., Mornar A, B.E., Bucan K, M.S.M. Evaluation of antioxidative activity of Croatian propolis samples using DPPH\* and ABTS\*+ stable free radical assays. *Molecules*. 2007 doi 10.3390/12051006
- [158]. Jasprica I, M.A., Debeljak Z, S.B.A., Medić-Sarić M, M.L., Romić Z, B.K., Balog T, S.S., V., S. In vivo study of propolis supplementation effects on antioxidative status and red blood cells. *J Ethnopharmacol*. 2007 doi 10.1016/j.jep.2006.10.023
- [159]. Jiang L, F.G., Zhang Y, C.G., S., W. Analysis of flavonoids in propolis and Ginkgo biloba by micellar electrokinetic capillary chromatography. *J Agric Food Chem*. 2008 doi 10.1021/jf802567r
- [160]. Jiang X, T.J., Zheng Y, Z.Y., Wu Y, Z.C., Zheng H, H.F. A New Propolis Type from Changbai Mountains in North-east China: Chemical Composition, Botanical Origin and Biological Activity. *Molecules*. 2019 doi 10.3390/molecules24071369
- [161]. Johnson KS, E.F., DE., G. Chemical composition of North American bee propolis and biological activity towards larvae of greater wax moth (Lepidoptera: Pyralidae). *J Chem Ecol*. 1994 doi 10.1007/BF02059899
- [162]. Jorge, R., Furtado, N.A.J.C., Sousa, J.P.B., da Silva Fiho, A.A., Gregorio, L.E.J., Martins, C.H.G., et al. Brazilian Propolis: Seasonal Variation of the Prenylated p-Coumaric Acids and Antimicrobial Activity. *PHARMACEUTICAL BIOLOGY*. 2008 doi 10.1080/13880200802370373
- [163]. Juliano, C., Pala, C.L., Cossu, M. Preparation and characterisation of polymeric films containing propolis. *JOURNAL OF DRUG DELIVERY SCIENCE AND TECHNOLOGY*. 2007 doi 10.1016/S1773-2247(07)50033-X
- [164]. Kalogeropoulos, N., Konteles, S.J., Troullidou, E., Mourtzinis, I., Karathanos, V.T. Chemical composition, antioxidant activity and antimicrobial properties of propolis extracts from Greece and Cyprus. *FOOD CHEMISTRY*. 2009 doi 10.1016/j.foodchem.2009.02.060
- [165]. Kanbur M, E.G., S., S. Antioxidant effect of propolis against exposure to propetamphos in rats. *Ecotoxicol Environ Saf*. 2009 doi 10.1016/j.ecoenv.2007.12.018

- [166]. Kartal M, K.S., S., K. GC-MS analysis of propolis samples from two different regions of Turkey. *Z Naturforsch C J Biosci.* 2002 doi 10.1515/znc-2002-9-1025
- [167]. Kharsany K, V.A., Leonard C, V.S. The new buzz: Investigating the antimicrobial interactions between bioactive compounds found in South African propolis. *J Ethnopharmacol.* 2019 doi 10.1016/j.jep.2019.111867
- [168]. Koc, N.A., Silic, S. Comparative study of in vitro methods used to analyse the antifungal activity of propolis against *Trichophyton rubrum* and *Trichophyton mentagrophytes*. *ANNALS OF MICROBIOLOGY.* 2008 doi 10.1007/BF03175555
- [169]. Koo, M., Park, Y. Investigation of flavonoid aglycones in propolis collected by two different varieties of bees in the same region. *BIOSCIENCE BIOTECHNOLOGY AND BIOCHEMISTRY.* 1997 doi 10.1271/bbb.61.367
- [170]. Koru O, T.F., Acikel CH, T.Y., Baysallar M, U.G.A., Akca E, O.T.A., Sorkun K, T.M., B., S. In vitro antimicrobial activity of propolis samples from different geographical origins against certain oral pathogens. *Anaerobe.* 2007 doi 10.1016/j.anaerobe.2007.02.001
- [171]. Kumazawa S, H.K., Kajiya K, I.T., Hamasaka T, N.T. Studies of the constituents of Uruguayan propolis. *J Agric Food Chem.* 2002 doi 10.1021/jf020279y
- [172]. Kumazawa S, N.J., Murase M, M.M., Ahn MR, F.S. Plant origin of Okinawan propolis: honeybee behavior observation and phytochemical analysis. *Naturwissenschaften.* 2008 doi 10.1007/s00114-008-0383-y
- [173]. Kumazawa S, U.R., Hamasaka T, F.S., Fujimoto T, N.T. Antioxidant prenylated flavonoids from propolis collected in Okinawa, Japan. *J Agric Food Chem.* 2007 doi 10.1021/jf071187h
- [174]. Kumazawa S, Y.M., Shibata I, K.J., Hamasaka T, N.T. Direct evidence for the plant origin of Brazilian propolis by the observation of honeybee behavior and phytochemical analysis. *Chem Pharm Bull (Tokyo).* 2003 doi 10.1248/cpb.51.740
- [175]. Kumazawa, S., Hamasaka, T., Nakayama, T. Antioxidant activity of propolis of various geographic origins. *FOOD CHEMISTRY.* 2004 doi 10.1016/S0308-8146(03)00216-4
- [176]. Kusumoto T, M.T.R., Doi S, S.H., H., Y. Isolation and structures of two new compounds from the essential oil of Brazilian propolis. *Chem Pharm Bull (Tokyo).* 2001 doi 10.1248/cpb.49.1207
- [177]. Li F, A.S., Tezuka Y, K.S. Cytotoxic constituents from Brazilian red propolis and their structure-activity relationship. *Bioorg Med Chem.* 2008 doi 10.1016/j.bmc.2008.04.016
- [178]. Li F, A.S., Tezuka Y, K.S. Cytotoxic constituents of propolis from Myanmar and their structure-activity relationship. *Biol Pharm Bull.* 2009 doi 10.1248/bpb.32.2075
- [179]. Li F, A.S., Tezuka Y, K.S. Cytotoxicity of constituents from Mexican propolis against a panel of six different cancer cell lines. *Nat Prod Commun.* 2010

- [180]. Li F, A.S., Zhang H, T.Y., Esumi H, K.S. Chemical constituents of propolis from Myanmar and their preferential cytotoxicity against a human pancreatic cancer cell line. *J Nat Prod.* 2009 doi 10.1021/np9002433
- [181]. Li YJ, X.H., Shou QY, Z.Z., Lu X, H.F. Therapeutic effects of propolis essential oil on anxiety of restraint-stressed mice. *Hum Exp Toxicol.* 2012 doi 10.1177/0960327111412805
- [182]. Li, H., Kapur, A., Yang, J.X., Srivastava, S., McLeod, D.G., Paredes-Guzman, J.F., et al. Antiproliferation of human prostate cancer cells by ethanolic extracts of Brazilian propolis and its botanical origin. *INTERNATIONAL JOURNAL OF ONCOLOGY.* 2007
- [183]. Lima B, T.A., Luna L, F.M., Schmeda-Hirschmann G, P.N., Wunderlin DA, F.G. Main flavonoids, DPPH activity, and metal content allow determination of the geographical origin of propolis from the Province of San Juan (Argentina). *J Agric Food Chem.* 2009 doi 10.1021/jf803866t
- [184]. Lima, V.H.M., Almeida, K.C.R., Alves, C.C.F., Rodrigues, M.L., Crotti, A.E.M., Souza, J.M., et al. Biological properties of volatile oil from Brazilian brown propolis. *Revista Brasileira de Farmacognosia.* 2020 doi 10.1016/j.bjp.2019.07.004
- [185]. Lirdprapamongkol K, S.H., Abdelhamed S, Y.S., Maruyama T, A.S., Viriyaroj A, A.S., Yagita H, R.S., Svasti J, S.I. A flavonoid chrysin suppresses hypoxic survival and metastatic growth of mouse breast cancer cells. *Oncol Rep.* 2013 doi 10.3892/or.2013.2667
- [186]. Lombardo-Agüí M, G.C.A., Gámiz-Gracia L, C.B.C. Determination of quinolones of veterinary use in bee products by ultra-high performance liquid chromatography-tandem mass spectrometry using a QuEChERS extraction procedure. *Talanta.* 2012 doi 10.1016/j.talanta.2012.02.011
- [187]. Londono Orozco, A., Avila Acevedo, J.G., Canales Martinez, M.M., Hernandez Delgado, C.T., Rocio Serrano, P., Flores Ortiz, C.M., et al. Antibacterial Comparative Study Between Extracts of Mexican Propolis and of Three Plants Which Use *Apis mellifera* for its Production. *JOURNAL OF ANIMAL AND VETERINARY ADVANCES.* 2010
- [188]. Lotti C, C.F.M., Piccinelli AL, C.R.O., Márquez Hernández I, R.L. Chemical constituents of red Mexican propolis. *J Agric Food Chem.* 2010 doi 10.1021/jf100070w
- [189]. Lotti C, P.A., Arevalo C, R.I., Migliani De Castro GM, F.R.S.L., Tesis AC, F.P.A., L., R. Constituents of Hondurian propolis with inhibitory effects on *Saccharomyces cerevisiae* multidrug resistance protein Pdr5p. *J Agric Food Chem.* 2012 doi 10.1021/jf302578r
- [190]. Lu, Y., Wu, C., Yuan, Z. Determination of hesperetin, cinnamic acid and nicotinic acid in propolis with micellar electrokinetic capillary chromatography. *FITOTERAPIA.* 2004 doi 10.1016/j.fitote.2003.12.026
- [191]. Luo, C., Zou, X., Li, Y., Sun, C., Jiang, Y., Wu, Z. Determination of flavonoids in propolis-rich functional foods by reversed phase high performance liquid chromatography with diode array detection. *FOOD CHEMISTRY.* 2011 doi 10.1016/j.foodchem.2011.01.006



- [192]. MAMONTOVA, N., TIKHONOV, A., SOBOLEVA, V. CHEMICAL-COMPOSITION OF PHENOL-POLYSACCHARIDE PROPOLIS PREPARATION. KHIMIYA PRIRODNYKH SOEDINENII. 1988
- [193]. Maciejewicz, W. Isolation of flavonoid aglycones from propolis by a column chromatography method and their identification by GC-MS and TLC methods. JOURNAL OF LIQUID CHROMATOGRAPHY & RELATED TECHNOLOGIES. 2001 doi 10.1081/JLC-100103439
- [194]. Marcucci MC, F.F., García-Viguera C, B.V., De Castro SL, D.A., Valente PH, P.N. Phenolic compounds from Brazilian propolis with pharmacological activities. J Ethnopharmacol. 2001 doi 10.1016/s0378-8741(00)00326-3
- [195]. Marcucci, M., Rodriguez, J., Ferreres, F., Bankova, V., Groto, R., Popov, S. Chemical composition of Brazilian propolis from Sao Paulo State. ZEITSCHRIFT FUR NATURFORSCHUNG C-A JOURNAL OF BIOSCIENCES. 1998
- [196]. Markham, K., Mitchell, K., Wilkins, A., Daldy, J., Lu, Y. HPLC and GC-MS identification of the major organic constituents in New Zealand propolis. PHYTOCHEMISTRY. 1996 doi 10.1016/0031-9422(96)83286-9
- [197]. Marostica, M.R.J., Dausch, A., Moraes, C.S., Queiroga, C.L., Pastore, G.M., Park, Y.K. Comparison of volatile and polyphenolic compounds in Brazilian green propolis and its botanical origin *Baccharis dracunculifolia*. CIENCIA E TECNOLOGIA DE ALIMENTOS. 2008 doi 10.1590/S0101-20612008000100026
- [198]. Martos, I., Cossentini, M., Ferreres, F., TomasBarberan, F. Flavonoid composition of Tunisian honeys and propolis. JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY. 1997 doi 10.1021/jf9609284
- [199]. Maruyama H, S.Y., Sakamoto T, A.Y., H., H. Antihypertensive effects of flavonoids isolated from brazilian green propolis in spontaneously hypertensive rats. Biol Pharm Bull. 2009 doi 10.1248/bpb.32.1244
- [200]. Mascheroni, E., Guillard, V., Nalin, F., Mora, L., Piergiovanni, L. Diffusivity of propolis compounds in Polylactic acid polymer for the development of anti-microbial packaging films. JOURNAL OF FOOD ENGINEERING. 2010 doi 10.1016/j.jfoodeng.2009.12.028
- [201]. Matsui T, E.S., Fujise T, A.K., Doi S, Y.H., K., M. Strong antihyperglycemic effects of water-soluble fraction of Brazilian propolis and its bioactive constituent, 3,4,5-tri-O-caffeoylquinic acid. Biol Pharm Bull. 2004 doi 10.1248/bpb.27.1797
- [202]. Matsuno T, M.Y., Saito M, M.J. Isolation and characterization of cytotoxic diterpenoid isomers from propolis. Z Naturforsch C J Biosci. 1997 doi 10.1515/znc-1997-9-1020
- [203]. Mauri P, P.P. Electrospray characterization of selected medicinal plant extracts. J Pharm Biomed Anal. 2000 doi 10.1016/s0731-7085(00)00264-8

- [204]. Mavri A, A.H., Polak T, B.J., Jamnik P, S.M.S., B., J. Chemical properties and antioxidant and antimicrobial activities of Slovenian propolis. *Chem Biodivers.* 2012 doi 10.1002/cbdv.201100337
- [205]. Medana C, C.F., Aigotti R, A.G., C., B. Selective analysis of phenolic compounds in propolis by HPLC-MS/MS. *Phytochem Anal.* 2008 doi 10.1002/pca.1010
- [206]. Medic-Saric, M., Bojic, M., Rastija, V., Cvek, J. Polyphenolic Profiling of Croatian Propolis and Wine. *FOOD TECHNOLOGY AND BIOTECHNOLOGY.* 2013
- [207]. Melliou E, C.I. Chemical analysis and antimicrobial activity of Greek propolis. *Planta Med.* 2004 doi 10.1055/s-2004-827150
- [208]. Melliou, E., Stratis, E., Chinou, I. Volatile constituents of propolis from various regions of Greece - Antimicrobial activity. *FOOD CHEMISTRY.* 2007 doi 10.1016/j.foodchem.2006.07.033
- [209]. Mello, B.C.B.S., Cunha Petrus, J.C., Hubinger, M.D. Concentration of flavonoids and phenolic compounds in aqueous and ethanolic propolis extracts through nanofiltration. *JOURNAL OF FOOD ENGINEERING.* 2010 doi 10.1016/j.jfoodeng.2009.08.040
- [210]. Memmedov H, O.L., Durmaz B, G.N., Yı Ldırım HK, S.E. Propolis prevents inhibition of apoptosis by potassium bromate in CCD 841 human colon cell. *Cell Biochem Funct.* 2020 doi 10.1002/cbf.3475
- [211]. Mendiola JA, M.A.P., Señoráns FJ, R.G., Capodicasa A, N.F., Sada A, C.A., E., I. Design of natural food antioxidant ingredients through a chemometric approach. *J Agric Food Chem.* 2010 doi 10.1021/jf901951z
- [212]. Menghinello, P., Cucchiari, L., Palma, F., Agostini, D., Dacha, M., Stocchi, V. Simultaneous analysis of flavonoid aglycones in natural products using an RP-HPLC method. *JOURNAL OF LIQUID CHROMATOGRAPHY & RELATED TECHNOLOGIES.* 1999 doi 10.1081/JLC-100102074
- [213]. Meto A, C.B., Meto A, B.G., Pinetti D, M.L., Benvenuti S, P.F., E., B. Propolis Affects *Pseudomonas aeruginosa* Growth, Biofilm Formation, eDNA Release and Phenazine Production: Potential Involvement of Polyphenols. *Microorganisms.* 2020 doi 10.3390/microorganisms8020243
- [214]. Midorikawa K, B.A., Tezuka Y, N.T., Matsushige K, M.D., Huertas AA, K.S. Liquid chromatography-mass spectrometry analysis of propolis. *Phytochem Anal.* 2001 doi 10.1002/pca.605
- [215]. Miguel MG, A.M. Is propolis safe as an alternative medicine?. *J Pharm Bioallied Sci.* 2011 doi 10.4103/0975-7406.90101
- [216]. Ming-Da, Z., Wei, M., Dong-Dong, C., Tao, P., Shu-Juan, L., Guang-Hua, Z., et al. Simultaneous Determination of Active Components and Anti-diabetic Drugs in Propolis Health

Foods by Ultra Performance Liquid Chromatography Tandem Mass Spectrometry. CHINESE JOURNAL OF ANALYTICAL CHEMISTRY. 2010 doi 10.3724/SP.J.1096.2010.00169

[217]. Mishima S, N.Y., Chikamatsu S, I.Y., Ohta S, Y.C., Araki Y, A.Y., Suzuki KM, N.Y. Effects of propolis on cell growth and gene expression in HL-60 cells. J Ethnopharmacol. 2005 doi 10.1016/j.jep.2005.02.005

[218]. Mohammadzadeh, S., Shariatpanahi, M., Hamed, M., Ahmadkhaniha, R., Samadi, N., Ostad, S.N. Chemical composition, oral toxicity and antimicrobial activity of Iranian propolis. FOOD CHEMISTRY. 2007 doi 10.1016/j.foodchem.2006.10.006

[219]. Moncla BJ, G.P., Wallace JA, M.M., Nor JE, B.W. The inhibitory activity of typified propolis against Enterococcus species. Z Naturforsch C J Biosci. 2012 doi 10.1515/znc-2012-5-603

[220]. Moraes, C.S., Dausch, A., Li, H., Rhim, J.S., Park, Y.K. Comparative antiproliferation of human prostate cancer cells by ethanolic extracts of two groups of Brazilian propolis. CIENCIA E TECNOLOGIA DE ALIMENTOS. 2010 doi 10.1590/S0101-20612010000200038

[221]. Márquez Hernández I, C.R.O., Campo Fernández M, R.P.A., Montes de Oca Porto R, P.A., L., R. Studies on the constituents of yellow Cuban propolis: GC-MS determination of triterpenoids and flavonoids. J Agric Food Chem. 2010 doi 10.1021/jf904527n

[222]. Nafady AM, E.S.M., Mohamed MH, H.H., Nohara T, Y.H., Ono M, S.H., Doi S, S.K., H., K. Cyclodextrin-enclosed substances of Brazilian propolis. Chem Pharm Bull (Tokyo). 2003 doi 10.1248/cpb.51.984

[223]. Nagaoka T, B.A., Tezuka Y, M.K., Matsushige K, K.S. Caffeic acid phenethyl ester (CAPE) analogues: potent nitric oxide inhibitors from the Netherlands propolis. Biol Pharm Bull. 2003 doi 10.1248/bpb.26.487

[224]. Naik, D.G., Mujumdar, A.M., Vaidya, H.S. Anti-inflammatory activity of propolis from Maharashtra, India. JOURNAL OF APICULTURAL RESEARCH. 2013 doi 10.3896/IBRA.1.52.2.06

[225]. Nani BD, S.J., Lazarini JG, S.D., Massariolli AP, C.T., de Alencar SM, F.M., PL., R. Anti-inflammatory and anti-Candida Effects of Brazilian Organic Propolis, a Promising Source of Bioactive Molecules and Functional Food. J Agric Food Chem. 2020 doi 10.1021/acs.jafc.8b07304

[226]. Narbona, E., Garcia-Garcia, E., Vazquez-Araujo, L., Carbonell-Barrachina, A.A. Volatile composition of functional 'a la Piedra'turron with propolis. INTERNATIONAL JOURNAL OF FOOD SCIENCE AND TECHNOLOGY. 2010 doi 10.1111/j.1365-2621.2009.02167.x

[227]. Nascimento, E.A., Chang, R., Morais, S.A.L., Pilo-Veloso, D., Reis, D.C. An easily detectable chemical marker for the Baccharis dracunculifolia propolis. REVISTA BRASILEIRA DE FARMACOGNOSIA-BRAZILIAN JOURNAL OF PHARMACOGNOSY. 2008 doi 10.1590/S0102-695X2008000300012

- [228]. Navarro-Navarro, M., Ruiz-Bustos, P., Valencia, D., Robles-Zepeda, R., Ruiz-Bustos, E., Virues, C., et al. Antibacterial Activity of Sonoran Propolis and Some of Its Constituents Against Clinically Significant *Vibrio* Species. *FOODBORNE PATHOGENS AND DISEASE*. 2013 doi 10.1089/fpd.2012.1318
- [229]. Negri, G., Marcucci, M., Salatino, A., Salatino, M. Comb and propolis waxes from Brazil: Triterpenoids in propolis waxes. *JOURNAL OF APICULTURAL RESEARCH*. 2000 doi 10.1080/00218839.2000.11101026
- [230]. Negri, G., Salatino, M., Salatino, A. 'Green propolis': unreported constituents and a novel compound from chloroform extracts. *JOURNAL OF APICULTURAL RESEARCH*. 2003 doi 10.1080/00218839.2003.11101087
- [231]. Negri, G., Salatino, M., Salatino, A. Unusual chemical composition of a sample of Brazilian propolis, as assessed by analysis of a chloroform extract. *JOURNAL OF APICULTURAL RESEARCH*. 2003 doi 10.1080/00218839.2003.11101092
- [232]. Netíková L, B.P., P., H. Czech ethanol-free propolis extract displays inhibitory activity against a broad spectrum of bacterial and fungal pathogens. *J Food Sci*. 2013 doi 10.1111/1750-3841.12230
- [233]. Nie P, X.Z., Sun DW, H.Y. Application of visible and near infrared spectroscopy for rapid analysis of chrysin and galangin in Chinese propolis. *Sensors (Basel)*. 2013 doi 10.3390/s130810539
- [234]. Nieva Moreno MI, Z.I., Ordóñez RM, J.G., Vattuone MA, I.M. Evaluation of the cytotoxicity, genotoxicity, mutagenicity, and antimutagenicity of propolis from Tucuman, Argentina. *J Agric Food Chem*. 2005 doi 10.1021/jf0513359
- [235]. Nolkemper S, R.J., Sensch KH, S.P. Mechanism of herpes simplex virus type 2 suppression by propolis extracts. *Phytomedicine*. 2010 doi 10.1016/j.phymed.2009.07.006
- [236]. Nunes CA, G.M. Characterization of Brazilian green propolis throughout the seasons by headspace GC/MS and ESI-MS. *J Sci Food Agric*. 2012 doi 10.1002/jsfa.4596
- [237]. Nunes, L.C.C., Galindo, A.B., Deus, A.S.O., Rufino, D.A., Randau, K.P., Xavier, H.S., et al. Variabilidade sazonal dos constituintes da própolis vermelha e bioatividade em *Artemia salina*. *Revista Brasileira de Farmacognosia*. 2009
- [238]. Olegário LS, A.J., Andrade GRS, D.M., Cavalcanti RL, S.M., N., N. Chemical characterization of four Brazilian brown propolis: An insight in tracking of its geographical location of production and quality control. *Food Res Int*. 2019 doi 10.1016/j.foodres.2019.04.004
- [239]. Oliveira, A.P., Franca, H.S., Kuster, R.M., Teixeira, L.A., Rocha, L.M. Chemical composition and antibacterial activity of Brazilian propolis essential oil. *JOURNAL OF VENOMOUS ANIMALS AND TOXINS INCLUDING TROPICAL DISEASES*. 2010 doi 10.1590/S1678-91992010005000007

- [240]. Opoka W, S.M., Maślanka A, P.R., Baś B, W.E., J., K. Optimization of method for zinc analysis in several bee products on renewable mercury film silver based electrode. *Acta Pol Pharm.* 2013
- [241]. Oroian M, D.F., F., U. Comparative evaluation of maceration, microwave and ultrasonic-assisted extraction of phenolic compounds from propolis. *J Food Sci Technol.* 2020 doi 10.1007/s13197-019-04031-x
- [242]. Oroian M, U.F., F., D. Influence of ultrasonic amplitude, temperature, time and solvent concentration on bioactive compounds extraction from propolis. *Ultrason Sonochem.* 2020 doi 10.1016/j.ultsonch.2020.105021
- [243]. Orsi RO, S.J., Funari SR, B.V. Effects of Brazilian and Bulgarian propolis on bactericidal activity of macrophages against *Salmonella Typhimurium*. *Int Immunopharmacol.* 2005 doi 10.1016/j.intimp.2004.10.003
- [244]. Orsolić N, B.I. Water-soluble derivative of propolis and its polyphenolic compounds enhance tumoricidal activity of macrophages. *J Ethnopharmacol.* 2005 doi 10.1016/j.jep.2005.05.036
- [245]. Oršolić N, S.D., Gajski G, G.V.V., Jazvinščak Jembrek M, K.I. Assessment of DNA damage and lipid peroxidation in diabetic mice: effects of propolis and epigallocatechin gallate (EGCG). *Mutat Res.* 2013 doi 10.1016/j.mrgentox.2013.04.022
- [246]. Ozen, T., Kilic, A., Bedir, O., Koru, O., Sorkun, K., Tanyuksel, M., et al. In Vitro Activity of Turkish Propolis Samples Against Anaerobic Bacteria Causing Oral Cavity Infections. *KAFKAS UNIVERSITESI VETERINER FAKULTESI DERGISI.* 2010
- [247]. Ozkok, D., Iscan, K.M., Silici, S. Effects of Dietary Propolis Supplementation on Performance and Egg Quality in Laying Hens. *JOURNAL OF ANIMAL AND VETERINARY ADVANCES.* 2013
- [248]. PARK, Y.K., IKEGAKI, M., ABREU, J.A.S., ALCICI, N.M.F. ESTUDO DA PREPARAÇÃO DOS EXTRATOS DE PRÓPOLIS E SUAS APLICAÇÕES. *Food Science and Technology.* 1998
- [249]. Pacheco, A., Daleprane, J.B., Freitas, V.S., Ferderbar, S., Hirabara, S., Cuevas, A., et al. Effect of Chilean Propolis on Glucose Metabolism in Diabetic Mice. *INTERNATIONAL JOURNAL OF MORPHOLOGY.* 2011 doi 10.4067/S0717-95022011000300015
- [250]. Pagliarone AC, O.C., Búfalo MC, M.F., Bachiega TF, J.J., JM., S. Propolis effects on pro-inflammatory cytokine production and Toll-like receptor 2 and 4 expression in stressed mice. *Int Immunopharmacol.* 2009 doi 10.1016/j.intimp.2009.08.005
- [251]. Pamplona-Zomenhan LC, P.B., da Silva CB, M.M., LM., M. Evaluation of the in vitro antimicrobial activity of an ethanol extract of Brazilian classified propolis on strains of *Staphylococcus aureus*. *Braz J Microbiol.* 2011 doi 10.1590/S1517-83822011000400002

- [252]. Papotti G, B.D., Bortolotti L, P.M. Chemical and functional characterization of Italian propolis obtained by different harvesting methods. J Agric Food Chem. 2012 doi 10.1021/jf205179d
- [253]. Papotti, G., Bertelli, D., Plessi, M., Rossi, M.C. Use of HR-NMR to classify propolis obtained using different harvesting methods. INTERNATIONAL JOURNAL OF FOOD SCIENCE AND TECHNOLOGY. 2010 doi 10.1111/j.1365-2621.2010.02310.x
- [254]. Park YK, A.S., CL., A. Botanical origin and chemical composition of Brazilian propolis. J Agric Food Chem. 2002 doi 10.1021/jf011432b
- [255]. Park YK, F.I., Ashida H, N.S., Yoshida K, D.A., Sato HH, P.G. Suppressive effects of ethanolic extracts from propolis and its main botanical origin on dioxin toxicity. J Agric Food Chem. 2005 doi 10.1021/jf058111a
- [256]. Park YK, P.G.J., Aguiar CL, A.S., FY., F. Chemical constituents in *Baccharis dracunculifolia* as the main botanical origin of southeastern Brazilian propolis. J Agric Food Chem. 2004 doi 10.1021/jf021060m
- [257]. Park, Y.K., Alencar, S.M., Scamparini, A.R.P., Aguiar, C.L. Própolis produzida no sul do Brasil, Argentina e Uruguai: evidências fitoquímicas de sua origem vegetal. Ciência Rural. 2002
- [258]. Paulino N, T.C., Martins R, S.A., Dirsch VM, V.A., Abreu SR, C.S., MC., M. Evaluation of the analgesic and anti-inflammatory effects of a Brazilian green propolis. Planta Med. 2006 doi 10.1055/s-2006-947185
- [259]. Paulino Zunini, M., Rojas, C., De Paula, S., Elingold, I., Alvareda Migliaro, E., Casanova, M.B., et al. PHENOLIC CONTENTS AND ANTIOXIDANT ACTIVITY IN CENTRAL-SOUTHERN URUGUAYAN PROPOLIS EXTRACTS. JOURNAL OF THE CHILEAN CHEMICAL SOCIETY. 2010
- [260]. Paviani, L.C., Dariva, C., Marcucci, M.C., Cabral, F.A. SUPERCRITICAL CARBON DIOXIDE SELECTIVITY TO FRACTIONATE PHENOLIC COMPOUNDS FROM THE DRY ETHANOLIC EXTRACT OF PROPOLIS. JOURNAL OF FOOD PROCESS ENGINEERING. 2010 doi 10.1111/j.1745-4530.2008.00256.x
- [261]. Pavlovic R, B.G., Leoni V, G.L., Ceciliani G, S.S., Bassoli A, G.A. Effectiveness of Different Analytical Methods for the Characterization of Propolis: A Case of Study in Northern Italy. Molecules. 2020 doi 10.3390/molecules25030504
- [262]. Pazin WM, S.S., Queiroz SCN, B., Soares AEE, M.I., AS., I. Bioactivity and action mechanism of green propolis against *Pythium aphanidermatum*. An Acad Bras Cienc. 2019 doi 10.1590/0001-3765201920180598
- [263]. Peev, C., Vlase, L., Dehelean, C., Soica, C., Feflea, S., Alexa, E. HPLC COMPARATIVE ANALYSIS OF POLYPHENOLIC CONTENT OF PROPOLIS AND BLACK POPLAR FOLIAR BUD EXTRACTS. ACTUAL TASKS ON AGRICULTURAL ENGINEERING, PROCEEDINGS. 2009

- [264]. Pellati F, P.F., S., B. Headspace solid-phase microextraction-gas chromatography-mass spectrometry characterization of propolis volatile compounds. *J Pharm Biomed Anal.* 2013 doi 10.1016/j.jpba.2013.05.045
- [265]. Peng, Y. Study on Capillary Electrophoresis-Amperometric Detection Profiles from Propolis and Its Medicinal Preparations. *ADVANCED ENGINEERING MATERIALS III*, PTS 1-3. 2013 doi 10.4028/www.scientific.net/AMR.750-752.1617
- [266]. Pereira AS, N.E., NF., A. Lupeol alkanoates in Brazilian propolis. *Z Naturforsch C J Biosci.* 2002 doi 10.1515/znc-2002-7-829
- [267]. Pereira AS, N.M., Cardoso JN, A.N.F., MF., R. Rapid screening of polar compounds in Brazilian propolis by high-temperature high-resolution gas chromatography-mass spectrometry. *J Agric Food Chem.* 2000 doi 10.1021/jf000594p
- [268]. Pereira, A., Bicalho, B., Radler, F., Neto, D. Comparison of propolis from *Apis mellifera* and *Tetragonisca angustula*. *APIDOLOGIE.* 2003 doi 10.1051/apido:2003023
- [269]. Pereira, A., Ramos, M., Pocas, E., Dias, P., dos Santos, E., da Silva, J., et al. Study of propolis by high temperature high resolution gas chromatography-mass spectrometry. *ZEITSCHRIFT FUR NATURFORSCHUNG SECTION C-A JOURNAL OF BIOSCIENCES.* 1999
- [270]. Permana AD, U.R., Courtenay AJ, M.M., Donnelly RF, R.L. Phytosomal nanocarriers as platforms for improved delivery of natural antioxidant and photoprotective compounds in propolis: An approach for enhanced both dissolution behaviour in biorelevant media and skin retention profiles. *J Photochem Photobiol B.* 2020 doi 10.1016/j.jphotobiol.2020.111846
- [271]. Picolotto A, P.D., Pereira GP, M.K., da Silva Barud H, R.E.M., Gonzalez MH, T.L., Figueiredo JG, M.S. Bacterial cellulose membrane associated with red propolis as phytomodulator: Improved healing effects in experimental models of diabetes mellitus. *Biomed Pharmacother.* 2019 doi 10.1016/j.biopha.2019.108640
- [272]. Pietta PG, G.C., AM., P. Analytical methods for quality control of propolis. *Fitoterapia.* 2002 doi 10.1016/s0367-326x(02)00186-7
- [273]. Popova M, B.V., Butovska D, P.V., Nikolova-Damyanova B, S.A., Marcazzan GL, B.S. Validated methods for the quantification of biologically active constituents of poplar-type propolis. *Phytochem Anal.* 2004 doi 10.1002/pca.777
- [274]. Popova M, C.C., Chen PY, H.C., V., B. A validated spectrophotometric method for quantification of prenylated flavanones in pacific propolis from Taiwan. *Phytochem Anal.* 2010 doi 10.1002/pca.1176
- [275]. Popova M, D.R., Al-Lawati HT, T.I., Najdenski H, B.V. Omani propolis: chemical profiling, antibacterial activity and new propolis plant sources. *Chem Cent J.* 2013 doi 10.1186/1752-153X-7-158

- [276]. Popova M, S.S., Kaftanoglu O, B.V. Antibacterial activity of Turkish propolis and its qualitative and quantitative chemical composition. *Phytomedicine*. 2005 doi 10.1016/j.phymed.2003.09.007
- [277]. Popova M, T.B., Cutajar S, A.D., Mifsud D, F.C., V., B. Identification of the plant origin of the botanical biomarkers of Mediterranean type propolis. *Nat Prod Commun*. 2012
- [278]. Popova MP, C.I., Marekov IN, B.V. Terpenes with antimicrobial activity from Cretan propolis. *Phytochemistry*. 2009 doi 10.1016/j.phytochem.2009.07.025
- [279]. Popova MP, G.K., Chinou I, B.V. GC-MS profiling of diterpene compounds in Mediterranean propolis from Greece. *J Agric Food Chem*. 2010 doi 10.1021/jf903841k
- [280]. Popova MP, T.B., Nedialkov PT, T.I., Pardo-Mora DP, N.H., Torres-García OA, S.J., VS., B. New  $\Delta$ -tocotrienol derivatives from Colombian propolis. *Nat Prod Res*. 2019 doi 10.1080/14786419.2019.1590710
- [281]. Popova, M., Trusheva, B., Khismatullin, R., Gavrilova, N., Legotkina, G., Lyapunov, J., et al. The Triple Botanical Origin of Russian Propolis from the Perm Region, Its Phenolic Content and Antimicrobial Activity. *NATURAL PRODUCT COMMUNICATIONS*. 2013
- [282]. Qian, W.L., Khan, Z., Watson, D.G., Fearnley, J. Analysis of sugars in bee pollen and propolis by ligand exchange chromatography in combination with pulsed amperometric detection and mass spectrometry. *JOURNAL OF FOOD COMPOSITION AND ANALYSIS*. 2008 doi 10.1016/j.jfca.2007.07.001
- [283]. Quiroga EN, S., Soberón JR, S.M., MA., V. Propolis from the northwest of Argentina as a source of antifungal principles. *J Appl Microbiol*. 2006 doi 10.1111/j.1365-2672.2006.02904.x
- [284]. Ramanauskienė, K., Savickas, A., Inkenienė, A., Vitkevicius, K., Kasparaviciene, G., Briedis, V., et al. Analysis of content of phenolic acids in Lithuanian propolis using high-performance liquid chromatography technique. *MEDICINA-LITHUANIA*. 2009 doi 10.3390/medicina45090093
- [285]. Reis JHO, B.G., Cerqueira JC, A.J., Andrade LN, P.F., Druzian JI, M.B. Evaluation of the antioxidant profile and cytotoxic activity of red propolis extracts from different regions of northeastern Brazil obtained by conventional and ultrasound-assisted extraction. *PLoS One*. 2019 doi 10.1371/journal.pone.0219063
- [286]. Ristivojević P, S.T., Starović M, P.S., Özcan MM, B.T., I., D. Phenolic composition and biological activities of geographically different type of propolis and black cottonwood resins against oral streptococci, vaginal microbiota and phytopathogenic *Fusarium* species. *J Appl Microbiol*. 2020 doi 10.1111/jam.14633
- [287]. Rivero-Cruz JF, G.P.J., Pedraza-Chaverri J, P.R.J., Kumar-Passari A, D.R.G., BE., R.C. Phytochemical Constituents, Antioxidant, Cytotoxic, and Antimicrobial Activities of the Ethanolic Extract of Mexican Brown Propolis. *Antioxidants (Basel)*. 2020 doi 10.3390/antiox9010070



- [288]. Rodrigues DM, S.M., Arruda C, P.R., JK., B. The Role of *Baccharis dracunculifolia* and its Chemical Profile on Green Propolis Production by *Apis mellifera*. J Chem Ecol. 2020 doi 10.1007/s10886-019-01141-w
- [289]. Russo A, T.N., Sanchez F, G.J., A., V. Propolis protects human spermatozoa from DNA damage caused by benzoapyrene and exogenous reactive oxygen species. Life Sci. 2006 doi 10.1016/j.lfs.2004.10.085
- [290]. Sahinler N, K.O. Natural product propolis: chemical composition. Nat Prod Res. 2005 doi 10.1080/14786410410001704877
- [291]. Salomão K, D.A., Borba CM, C.L., Machado DG, A.N.F., SL., C. Chemical composition and microbicidal activity of extracts from Brazilian and Bulgarian propolis. Lett Appl Microbiol. 2004 doi 10.1111/j.1472-765x.2003.01458.x
- [292]. Salomão K, P.P., Campos LC, B.C., Cabello PH, M.M., SL., C. Brazilian propolis: correlation between chemical composition and antimicrobial activity. Evid Based Complement Alternat Med. 2008 doi 10.1093/ecam/nem058
- [293]. Santos, H.C., Vieira, D.S., Yamamoto, S.M., Costa, M.M., Sá, M.C., Silva, E.M., et al. Antimicrobial activity of propolis extract fractions against *Staphylococcus* spp. isolated from goat mastitis. Pesquisa Veterinária Brasileira. 2020 doi 10.1590/1678-5150-pvb-5940
- [294]. Sawaya, A., Souza, K., Marcucci, M., Cunha, I., Shimizu, M. Analysis of the composition of Brazilian propolis extracts by chromatography and evaluation of their in vitro activity against Gram-positive bacteria. BRAZILIAN JOURNAL OF MICROBIOLOGY. 2004 doi 10.1590/S1517-83822004000100017
- [295]. Schnitzler P, N.A., Nolkemper S, Z.C., Nowack H, S.K., J., R. Antiviral activity and mode of action of propolis extracts and selected compounds. Phytother Res. 2010 doi 10.1002/ptr.2868
- [296]. Seibert JB, B.S.J., Amparo TR, P.A., Pervier P, S.A.J., Azevedo MC, S.B., Brandão GC, S.G., de Medeiros Teixeira LF, S.O. Development of propolis nanoemulsion with antioxidant and antimicrobial activity for use as a potential natural preservative. Food Chem. 2019 doi 10.1016/j.foodchem.2019.02.078
- [297]. Sha N, H.H., Zhang JQ, C.G., Tao SJ, Y.M., Li XN, L.P., DA., G. Simultaneous quantification of eight major bioactive phenolic compounds in Chinese propolis by high-performance liquid chromatography. Nat Prod Commun. 2009
- [298]. Shaheen SA, Z.M., Nazer IK, D.R., HI., A.J. Chemical constituents of Jordanian propolis. Nat Prod Res. 2011 doi 10.1080/14786419.2010.509060
- [299]. Shi H, Y.H., Zhang X, S.Y., Huang H, Y.L. Isolation and characterization of five glycerol esters from Wuhan propolis and their potential anti-inflammatory properties. J Agric Food Chem. 2012 doi 10.1021/jf302601m

- [300]. Shimizu K, A.H., Matsuura Y, K.K. Antioxidative bioavailability of artemisinin C in Brazilian propolis. Arch Biochem Biophys. 2004 doi 10.1016/j.abb.2004.02.021
- [301]. Shrestha SP, N.Y., T., T. Chemical constituents of Nepalese propolis (II). Chem Pharm Bull (Tokyo). 2007 doi 10.1248/cpb.55.926
- [302]. Shrestha, S.P., Narukawa, Y., Takeda, T. Chemical constituents of Nepalese propolis: isolation of new dalbergiones and related compounds. JOURNAL OF NATURAL MEDICINES. 2007 doi 10.1007/s11418-006-0024-8
- [303]. Siess, M., LeBon, A., Canivenc-Lavier, M., Amiot, M., Sabatier, S., Aubert, S., et al. Flavonoids of honey and propolis: Characterization and effects on hepatic drug-metabolizing enzymes and benzo(a)pyrene-DNA binding in rats. JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY. 1996 doi 10.1021/jf9504733
- [304]. Siheri W, E.G., Igoli JO, G.A., Biddau M, A.P., Alenezi S, A.M., Edrada-Ebel R, M.S., Lawrence CE, F.J., et al. Isolation of a Novel Flavanonol and an Alkylresorcinol with Highly Potent Anti-Trypanosomal Activity from Libyan propolis. Molecules. 2019 doi 10.3390/molecules24061041
- [305]. Silici S, K.S. Chemical composition and antibacterial activity of propolis collected by three different races of honeybees in the same region. J Ethnopharmacol. 2005 doi 10.1016/j.jep.2005.01.046
- [306]. Silici S, Ü.M., G., V.Ü. Antibacterial activity and phytochemical evidence for the plant origin of Turkish propolis from different regions. World J Microbiol Biotechnol. 2007 doi 10.1007/s11274-007-9430-7
- [307]. Silva BB, R.P., Cury JA, I.M., Souza VC, E.A., SM., A. Chemical composition and botanical origin of red propolis, a new type of Brazilian propolis. Evid Based Complement Alternat Med. 2008 doi 10.1093/ecam/nem059
- [308]. Silva, M.S.S., Citó, A.M.G.L., Chaves, M.H., Lopes, J.A.D. Triterpenóides tipo cicloartano de própolis de Teresina - PI. Química Nova. 2005
- [309]. Silva, M.S.S., De Lima, S.G., Oliveira, E.H., Lopes, J.A.D., Chaves, M.H., Reis, F.A.M., et al. Anacardic acid derivatives from Brazilian propolis and their antibacterial activity. Eclética Química. 2008
- [310]. Simões LM, G., Da Silva Filho AA, S.M., Azzolini AE, B.J., YM., L.V. Effect of Brazilian green propolis on the production of reactive oxygen species by stimulated neutrophils. J Ethnopharmacol. 2004 doi 10.1016/j.jep.2004.04.026
- [311]. Siripatrawan, U., Vitchayakitti, W., Sanguandeekul, R. Antioxidant and antimicrobial properties of Thai propolis extracted using ethanol aqueous solution. INTERNATIONAL JOURNAL OF FOOD SCIENCE AND TECHNOLOGY. 2013 doi 10.1111/j.1365-2621.2012.03152.x

- [312]. Skaba D, M.T., Tanasiewicz M, M.A., Bobela E, S.E., Skucha-Nowak M, D.M., Yamamoto R, I.S., Makita Y, R.M., et al. Influence of the toothpaste with brazilian ethanol extract propolis on the oral cavity health. *Evid Based Complement Alternat Med*. 2013 doi 10.1155/2013/215391
- [313]. Slavov, A., Trifonov, A., Peychev, L., Dimitrova, S., Peycheva, S., Gotcheva, V., et al. BIOLOGICALLY ACTIVE COMPOUNDS WITH ANTITUMOR ACTIVITY IN PROPOLIS EXTRACTS FROM DIFFERENT GEOGRAPHIC REGIONS. *BIOTECHNOLOGY & BIOTECHNOLOGICAL EQUIPMENT*. 2013 doi 10.5504/BBEQ.2013.0034
- [314]. Sobocanec S, S.V., Balog T, S.A., Rusak G, L.S., Kusić B, K.V., Radić S, M.T. Oxidant/antioxidant properties of Croatian native propolis. *J Agric Food Chem*. 2006 doi 10.1021/jf0612023
- [315]. Sobocanec, S., Balog, T., Saric, A., Macak-Safranko, Z., Stroser, M., Zarkovic, K., et al. Antitumor effect of Croatian propolis as a consequence of diverse sex-related dihydropyrimidine dehydrogenase (DPD) protein expression. *PHYTOMEDICINE*. 2011 doi 10.1016/j.phymed.2011.01.015
- [316]. Sousa, J.P., Furtado, N.A., Jorge, R., Soares, A.E., Bastos, J.K. Perfis físico-químico e cromatográfico de amostras de própolis produzidas nas microrregiões de Franca (SP) e Passos (MG), Brasil. *Revista Brasileira de Farmacognosia*. 2007
- [317]. Souza, J.P.B., Tacon, L.A., Correia, C.C., Bastos, J.K., Freitas, L.A.P. Spray-dried propolis extract, II: Prenylated components of green propolis. *PHARMAZIE*. 2007 doi 10.1691/ph.2007.7.6286
- [318]. Souza, R.M., de Souza, M.C., Patitucci, M.L., Silva, J.F.M. Evaluation of antioxidant and antimicrobial activities and characterization of bioactive components of two Brazilian propolis samples using a pK(a)-guided fractionation. *ZEITSCHRIFT FÜR NATURFORSCHUNG SECTION C-A JOURNAL OF BIOSCIENCES*. 2007 doi 10.1515/znc-2007-11-1205
- [319]. Strehl E, V.R., EF., E. Biochemical activities of propolis-extracts. III. Inhibition of dihydrofolate reductase. *Z Naturforsch C J Biosci*. 1994 doi 10.1515/znc-1994-1-207
- [320]. Sun S, L.M., He J, L.K., Zhang X, Y.G. Identification and Determination of Seven Phenolic Acids in Brazilian Green Propolis by UPLC-ESI-QTOF-MS and HPLC. *Molecules*. 2019 doi 10.3390/molecules24091791
- [321]. Szliszka E, K.A., Sokół-Łętowska A, M.A., Czuba ZP, K.W. Chemical Composition and Anti-Inflammatory Effect of Ethanolic Extract of Brazilian Green Propolis on Activated J774A.1 Macrophages. *Evid Based Complement Alternat Med*. 2013 doi 10.1155/2013/976415
- [322]. Szliszka E, S.Ł.A., Kucharska AZ, J.D., Czuba ZP, K.W. Ethanolic Extract of Polish Propolis: Chemical Composition and TRAIL-R2 Death Receptor Targeting Apoptotic Activity against Prostate Cancer Cells. *Evid Based Complement Alternat Med*. 2013 doi 10.1155/2013/757628

- [323]. TOMASBARBERAN, F., GARCIAVIGUERA, C., VITOLIVIER, P., FERRERES, F., TOMASLORENTE, F. PHYTOCHEMICAL EVIDENCE FOR THE BOTANICAL ORIGIN OF TROPICAL PROPOLIS FROM VENEZUELA. PHYTOCHEMISTRY. 1993 doi 10.1016/S0031-9422(00)90804-5
- [324]. Tani H, H.K., Tatefuji T, H.K., Koshino H, T.S. Inhibitory activity of Brazilian green propolis components and their derivatives on the release of cys-leukotrienes. Bioorg Med Chem. 2010 doi 10.1016/j.bmc.2009.11.007
- [325]. Tavares DC, L.W., Santini CB, T.C., JK., B. Effects of propolis crude hydroalcoholic extract on chromosomal aberrations induced by doxorubicin in rats. Planta Med. 2007 doi 10.1055/s-2007-993737
- [326]. Teixeira EW, N.G., Meira RM, M.D., A., S. Plant Origin of Green Propolis: Bee Behavior, Plant Anatomy and Chemistry. Evid Based Complement Alternat Med. 2005 doi 10.1093/ecam/neh055
- [327]. Teixeira, T., Message, D., Negri, G., Salatino, A. Bauer-7-en-3 beta-yl acetate: A major constituent of unusual samples of Brazilian propolis. QUIMICA NOVA. 2006
- [328]. Tomazzoli, M.M., Zeggio, A.R.S., Pai Neto, R.D., Specht, L., Costa, C., Rocha, M., et al. Botanical source investigation and evaluation of the effect of seasonality on Brazilian propolis from *Apis mellifera* L. Scientia Agricola. 2020 doi 10.1590/1678-992x-2018-0258
- [329]. Tomažević T, J.J. A double blind randomised placebo controlled study of propolis (bee glue) effectiveness in the treatment of severe oral mucositis in chemotherapy treated children. Complement Ther Med. 2013 doi 10.1016/j.ctim.2013.04.002
- [330]. Torres, R.N.S., Lopes, J.A.D., Moita Neto, J.M., Citó, A.M.G.L. Constituintes voláteis de própolis piauiense. Química Nova. 2008 doi 10.1590/S0100-40422008000300003
- [331]. Tozuka, Y., Imono, M., Uchiyama, H., Tahara, K., Tazawa, S., Araki, Y., et al. Dry powder formulation with alpha-glycosyltransferase-treated stevia for the effective absorption of hydrophobic bioactive compounds in crude drugs. POWDER TECHNOLOGY. 2013 doi 10.1016/j.powtec.2012.07.014
- [332]. Tran VH, D.R., Abu-Mellal A, D.C. Propolis with high flavonoid content collected by honey bees from *Acacia paradoxa*. Phytochemistry. 2012 doi 10.1016/j.phytochem.2012.06.002
- [333]. Trusheva B, P.M., Bankova V, S.S., Marcucci MC, M.P., da Rocha Pasin F, T.I. Bioactive constituents of brazilian red propolis. Evid Based Complement Alternat Med. 2006 doi 10.1093/ecam/nel006
- [334]. Trusheva B, P.M., Naydenski H, T.I., Gregorio Rodriguez J, B.V. New polyisoprenylated benzophenones from Venezuelan propolis. Fitoterapia. 2004 doi 10.1016/j.fitote.2004.08.001

- [335]. Tukmechi A, O.A., A., M. In vitro antibacterial activities of ethanol extract of iranian propolis (EEIP) against fish pathogenic bacteria (*Aeromonas hydrophila*, *Yersinia ruckeri* & *Streptococcus iniae*). *Braz J Microbiol*. 2010 doi 10.1590/S1517-838220100004000030
- [336]. Tukmechi, A., Ownagh, A., Mohebbat, A. In vitro antibacterial activities of ethanol extract of iranian propolis (EEIP) against fish pathogenic bacteria (*Aeromonas hydrophila*, *Yersinia ruckeri* & *Streptococcus iniae*). *Brazilian Journal of Microbiology*. 2010
- [337]. Usia T, B.A., Tezuka Y, M.K., Matsushige K, K.S. Constituents of Chinese propolis and their antiproliferative activities. *J Nat Prod*. 2002 doi 10.1021/np010486c
- [338]. Uzel A, S.K., Onçağ O, C.D., Gençay O, S.B. Chemical compositions and antimicrobial activities of four different Anatolian propolis samples. *Microbiol Res*. 2005 doi 10.1016/j.micres.2005.01.002
- [339]. Velikova M, B.V., Tsvetkova I, K.A., MC., M. Antibacterial ent-kaurene from Brazilian propolis of native stingless bees. *Fitoterapia*. 2000 doi 10.1016/s0367-326x(00)00213-6
- [340]. Volpi N, B.G. Analysis of flavonoids from propolis by on-line HPLC-electrospray mass spectrometry. *J Pharm Biomed Anal*. 2006 doi 10.1016/j.jpba.2006.04.017
- [341]. Wang T, L.Q., Wang M, Z.L. Metabolomics Reveals Discrimination of Chinese Propolis from Different Climatic Regions. *Foods*. 2020 doi 10.3390/foods9040491
- [342]. Wang, S.P., Fu, M.D., Wang, M.H. Separation mechanism and determination of flavanones with capillary electrophoresis and high-performance liquid chromatography. *JOURNAL OF CHROMATOGRAPHY A*. 2007 doi 10.1016/j.chroma.2007.06.072
- [343]. Watson DG, P.E., Zheng L, L.D., Seidel V, J.B., Parkinson JA, F.J. Application of principal components analysis to <sup>1</sup>H-NMR data obtained from propolis samples of different geographical origin. *Phytochem Anal*. 2006 doi 10.1002/pca.921
- [344]. Weng MS, L.C., Chen CN, W.C., JK., L. Propolin H from Taiwanese propolis induces G1 arrest in human lung carcinoma cells. *J Agric Food Chem*. 2007 doi 10.1021/jf070201n
- [345]. Wojtyczka RD, D.A., Idzik D, K.M., Kubina R, K.D.A., Smoleń-Dzirba J, S.J., Sajewicz M, W.T. Susceptibility of *Staphylococcus aureus* clinical isolates to propolis extract alone or in combination with antimicrobial drugs. *Molecules*. 2013 doi 10.3390/molecules18089623
- [346]. Xuan, H., Zhao, J., Miao, J., Li, Y., Chu, Y., Hu, F. Effect of Brazilian propolis on human umbilical vein endothelial cell apoptosis. *FOOD AND CHEMICAL TOXICOLOGY*. 2011 doi 10.1016/j.fct.2010.09.034
- [347]. Yang, L., Yan, Q.H., Ma, J.Y., Wang, Q., Zhang, J.W., Xi, G.X. High Performance Liquid Chromatographic Determination of Phenolic Compounds in Propolis. *TROPICAL JOURNAL OF PHARMACEUTICAL RESEARCH*. 2013 doi 10.4314/tjpr.v12i5.17

- [348]. Yang, S.Z., Peng, L.T., Su, X.J., Chen, F., Cheng, Y.J., Fan, G., et al. Bioassay-guided isolation and identification of antifungal components from propolis against *Penicillium italicum*. FOOD CHEMISTRY. 2011 doi 10.1016/j.foodchem.2010.12.011
- [349]. Yildirim Z, H.S., Kutlu NO, A.N., Kurkcuoglu M, I.M., R., D. Effect of water extract of Turkish propolis on tuberculosis infection in guinea-pigs. Pharmacol Res. 2004 doi 10.1016/j.phrs.2003.10.007
- [350]. Yonar, M.E., Yonar, S.M., Silici, S. Protective effect of propolis against oxidative stress and immunosuppression induced by oxytetracycline in rainbow trout (*Oncorhynchus mykiss*, W.). FISH & SHELLFISH IMMUNOLOGY. 2011 doi 10.1016/j.fsi.2011.05.019
- [351]. Yu-Xin, F., Yuan-Jun, X., Bin, C., Yan, L., Li-Ping, L. Analysis of Volatile Components from Inner Mongolia Propolis by Gas Chromatography-Mass Spectrometry. CHINESE JOURNAL OF ANALYTICAL CHEMISTRY. 2009
- [352]. Zeitoun R, N.F., Wehbi B, K.A., Fayyad-Kazan M, D.H.C., Faour WH, E.M.Y. Chemical Composition, Antioxidant and Anti-inflammatory Activity Evaluation of the Lebanese Propolis Extract. Curr Pharm Biotechnol. 2019 doi 10.2174/1389201020666190206201241
- [353]. Zhou J, L.Y., Zhao J, X.X., Wu L, C.F. Geographical traceability of propolis by high-performance liquid-chromatography fingerprints. Food Chem. 2008 doi 10.1016/j.foodchem.2007.11.009
- [354]. Zilius M, R.K., V., B. Release of propolis phenolic acids from semisolid formulations and their penetration into the human skin in vitro. Evid Based Complement Alternat Med. 2013 doi 10.1155/2013/958717
- [355]. da Silva Filho AA, S.J., Soares S, F.N., Andrade e Silva ML, C.W., Gregório LE, N.N., JK., B. Antimicrobial activity of the extract and isolated compounds from *Baccharis dracunculifolia* D. C. (Asteraceae). Z Naturforsch C J Biosci. 2008 doi 10.1515/znc-2008-1-208
- [356]. da Silva, M., Cito, A., Chaves, M., Lopes, J. Cycloartane triterpenoids of propolis from teresina. QUIMICA NOVA. 2005 doi 10.1590/S0100-40422005000500013
- [357]. da Silveira Regueira-Neto M, R.T.S., Pereira da Silva AR, S.C.M., de Moraes Oliveira-Tintino CD, A.B.A., Menezes IRA, Q.B.V., HD., M.C. Comparative Analysis of the Antibacterial Activity and HPLC Phytochemical Screening of the Brazilian Red Propolis and the Resin of *Dalbergia ecastaphyllum*. Chem Biodivers. 2019 doi 10.1002/cbdv.201900344
- [358]. de Albuquerque, I.L., Alves, L.A., Lemos, T.L.G., Dorneles, C.A., de Moraes, M.O. Constituents of the essential oil of Brazilian green propolis from Brazil. JOURNAL OF ESSENTIAL OIL RESEARCH. 2008 doi 10.1080/10412905.2008.9700044
- [359]. de Albuquerque, I.L., Alves, L.A., Lemos, T.L.G., Monte, F.J.Q., Braz Filho, R. Canaric acid (3,4-seco-lupane derivative) isolated from propolis of ceara. QUIMICA NOVA. 2007

- [360]. de Castro Ishida, V.F., Negri, G., Salatino, A., Bandeir, M.F.C.L. A new type of Brazilian propolis: Prenylated benzophenones in propolis from Amazon and effects against cariogenic bacteria. *FOOD CHEMISTRY*. 2011 doi 10.1016/j.foodchem.2010.09.089
- [361]. de Funari CS, O.F.V., MB., M. Analysis of propolis from *Baccharis dracunculifolia* DC. (Compositae) and its effects on mouse fibroblasts. *J Ethnopharmacol*. 2007 doi 10.1016/j.jep.2006.11.032
- [362]. de Sousa JP, B.P., Gregório LE, S.F.A., Furtado NA, S.M., JK., B. A reliable quantitative method for the analysis of phenolic compounds in Brazilian propolis by reverse phase high performance liquid chromatography. *J Sep Sci*. 2007 doi 10.1002/jssc.200700228
- [363]. dos Santos Pereira A, M.P.A., Trugo LC, A.N.F. Distribution of quinic acid derivatives and other phenolic compounds in Brazilian propolis. *Z Naturforsch C J Biosci*. 2003 doi 10.1515/znc-2003-7-824
- [364]. dos Santos TF, A.A., Dórea HS, N.S. MSPD procedure for determining buprofezin, tetradifon, vinclozolin, and bifenthrin residues in propolis by gas chromatography-mass spectrometry. *Anal Bioanal Chem*. 2008 doi 10.1007/s00216-007-1813-y