# Numerical methods for stochastic volatility models:

### Heston model and extensions

A Thesis Presented to The Division of Applied Mathematics Fundação Getulio Vargas

 $\label{eq:continuous} \begin{tabular}{ll} In Partial Fulfillment \\ of the Requirements for the Degree \\ M.Sc. of Mathematics \\ \end{tabular}$ 

Fernando O. Teixeira

May 2017

Approved for the Division (Mathematics)

Hugo de la Cruz

# Acknowledgements

I want to thank a few people.

# Table of Contents

Chapt	er 1: I	Delete line 6 if you only have one advisor	 1
Chapt	er 2: L	Literature Review	 3
2.1	Stocha	nastic Calculus	 3
	2.1.1		
	2.1.2		4
	2.1.3	TODO	4
	2.1.4		
2.2	Black-	z-Scholes Model	
	2.2.1		
2.3	Stocha	astic Volatility models	
2.0	2.3.1	•	-
	2.0.1	Other models	8
Chapt	er 3: N	Models	 9
Chapt	er 4: T	This chunk ensures that the thesisdown package is	 11
Chapt	er 5: C	Conclusion	 13
Chapt	er 6: <b>T</b>	The First Appendix	 15
Refere	ences .		17

# List of Tables

0.1	Box calculus.																															
∠.⊥	Dox calculus.	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•		٠

# List of Figures

### Abstract

The preface pretty much says it all. Second paragraph of abstract starts here.

### Dedication

You can have a dedication here if you wish.

Delete line 6 if you only have one advisor

### Literature Review

This chapter presents the concepts of stochastic calculus, from the historic conception of how it first arose through the basic principles and applications in finance. More precisely, we address the classical Black-Scholes model and its limitations and the Heston model. This model is also well known, it introduces the concept of stochastic volatility which brings us closer to reality.

#### 2.1 Stochastic Calculus

Stochastic calculus arises from stochastic processes and allows the creation of a theory of integration where both the integrand and integrator terms are stochastic processes. Stochastic calculus, also known as, Itô calculus due to the name of it's creator, the Japanese mathematician Kiyosi Itô in the 1940s and 1950s is used for modelling financial options and in another wide variety of fields [1]. In this chapter we present the historical contexts in which the tools and models used arise, but our focus is introducing the concepts and notations that will be further used in our work.

#### 2.1.1 Brownian Motion

The Brownian motion is the name given to the irregular motion observed in the motion of pollen particles suspended in fluid resulting from particle collision with atoms or molecules. It is named after Robert Brown, the first to have observed the movement in 1828. He noted two characteristic in the pollen movement [1]:

- the path of a given particle is very irregular, having a tangent at no point
- the motion of two distinct particles appear to be independent

The first quantitative works in brownian motion come from an interest in stock price fluctuation by Bachelier in 1900. Albert Einstein also leaned over the subject and in 1905 derived the transition density for Brownian motion from molecular-kinetic theory of heat [1,2].

In 1923, the Wiener process was coined in honor of Norbert Wiener mathematical proof of existence of the brownian motion and stating its properties as follows [3]:

- $W_0 = 0$
- The change in W, given by  $\Delta W = W_{t+1} W_t$ , is normally distributed with mean zero and standard deviation  $\sqrt{\Delta t}$ , meaning that  $\Delta W = \epsilon \sqrt{\Delta t}$ , where  $\epsilon$  is N(0,1).
- If the increment  $\Delta t_1$  does not overlap with the time increment  $\Delta t_2$ , then  $\Delta W_1$  and  $\Delta W_2$  are independent.
- The process is continuous, meaning that there are no jumps in the process.
- The process is a Markov process. This means that the conditional expectation of  $W_{t+1}$  given its entire history is equal to the conditional expectation of  $W_{t+1}$  given today's information. This can be written as:  $E[W_{t+1}|W_1,...,W_t] = E[W_{t+1}|W_t]$ .
- Consider the time interval [0,t] with n equally spaced intervals given by  $t_i = \frac{it}{n}$ . Then the paths of the Brownian motion have unbounded variation, this means that they are not differentiable and go towards infinity as n increases. The quadratic variation is given by  $\sum_{i=1}^{n} (Z_{t_i} Z_{t_{i-1}})^2 \to t$ , meaning that when n increases it stays constant at t.

#### 2.1.2 Correlated Brownian Motions

Two independent brownian motions that are correlated can describe a new process  $Z_t$ . Let  $W^1$  and  $W^2$  be these two *independent* Brownian motions and let  $-1 \le \rho \le 1$  be a given number. For  $0 \le t \le T$  define the new process  $Z_t$  as [1]:

$$Z_t = \rho W_t^1 sqrt(1 - \rho^2) W_t^2 \tag{2.1}$$

This equation is a linear combination of independent normals at each timestep t, so  $Z_t$  is normally distributed. It is proven that Z is a Brownian motion and that Z and  $W_t^1$  are correlated [1].

#### 2.1.3 TODO

Describe Arithmetic and Geometric brownian motions

#### 2.1.4 Itô's Lemma

Let  $X_t$  be a real-valued stochastic process that satisfies [4–6]:

$$S_{t} = S_{0} + \int_{0}^{t} \mu_{t} dt + \int_{0}^{t} \sigma_{t} dW_{t}$$
 (2.2)

for some  $\mu_t$ ,  $\sigma_t$  and  $t \in [0, T]$ . This equation is often rewritten in its differential stochastic form:

$$dS_t = \mu_t dt + \sigma_t dW_t \tag{2.3}$$

for  $0 \le t \le T$ .

#### Theorem

Assume that  $S_t$  has a stochastic differential given by:

$$dS_t = \mu_t dt + \sigma_t dW_t \tag{2.4}$$

for  $\mu_t$ ,  $\sigma_t$  and  $t \in [0,T]$ . Assume  $u : \mathbb{R} \times [0,T] \to \mathbb{R}$  is continuous and that  $\frac{\partial u}{\partial t}$ ,  $\frac{\partial u}{\partial x}$ ,  $\frac{\partial^2 u}{\partial x^2}$ exist and are continuous.

$$Y_t := u(S_t, t)$$

Then Y has the following stochastic differential:

$$dY_{t} = \frac{\partial u}{\partial t}dt + \frac{\partial u}{\partial x}dS_{t} + \frac{1}{2}\frac{\partial^{2} u}{\partial x^{2}}\sigma_{t}^{2}dt$$

$$= \left(\frac{\partial u}{\partial t} + \mu_{t}\frac{\partial u}{\partial x} + \frac{1}{2}\frac{\partial^{2} u}{\partial x^{2}}\sigma_{t}^{2}\right)dt + \sigma_{t}\frac{\partial u}{\partial x}dW_{t}$$
(2.5)

where the argument of u,  $\frac{\partial u}{\partial x}$  and  $\frac{\partial^2 u}{\partial x^2}$  above is  $(S_t, t)$ . Equation (2.5) is the stochastic equivalent to the chain rule, also known as Itô's formula or Itô's chain rule. The proof to this theorem is based on the Taylor expansion of the function  $f(S_t,t)$  [4,5]. For practical uses you should write out a second-order Taylor expansion for the function to be analyzed and apply the 2.1 multiplication table [1].

Table 2.1: Box calculus

	dt	$dW_t$
dt	0	0
$dW_t$	0	dt

#### 2.2 **Black-Scholes Model**

The Black-Scholes (B-S) model arises from the need to price european options in the derivative markets. Derivatives are financial instruments traded in the market, stock exchange or over-the-counter (OTC) market, whose values depend on the values of an underlying asset. [7–9]

- A call option is a derivative that gives its bearer the right, but not the obligation, to purchase a specific asset by a fixed price before or on a given date.
- A put option is a derivative that gives its bearer the right, but not the obligation, to sell a specific asset by a fixed price before or on a given date.

The trading price of the option is called the option *premium* and the asset from which the option derives is called the *underlying asset*. This asset may be the interest rate, exchange rates, stock exchanges rates, commodities or stocks. The fixed price in contract in which the underlying asset might to be bought or sold is the *strick price*. The option expiration date is called the *maturity*. [8,9]

There are two major different option types: the European and the American. The difference between these two is that the bearer of the first may exercise it only at the end of its life, at its maturity while the latter can be exercised at any given time until its maturity. [9,10]

#### 2.2.1 The model

The Black-Scholes model that provides analytical solution to the price of a European call at time t can be described as follows [3,7,9]:

$$C(S_t, t) = N(d_1)S_t - N(d_2)Ke^{-r(T-t)}$$
(2.6)

$$d_1 = \frac{1}{\sigma\sqrt{T-t}} \left[ \ln\left(\frac{S_t}{K}\right) + \left(r + \frac{\sigma^2}{2}\right) (T-t) \right]$$
 (2.7)

$$d_2 = d_1 - \sigma\sqrt{T - t} \tag{2.8}$$

Where:

- $S_t$  is the spot price of the underlying asset at time t
- r is the risk free rate (generally an annual rate)<sup>1</sup>
- $\sigma$  is the volatility of returns of the underlying asset <sup>2</sup>
- $N(\cdot)$  is the cumulative distribution function of the standard Gaussian distribution
- K is the strike price
- T-t is the time to maturity

Also, the stock price path is a Geometric Brownian Motion and is under the risk-neutral measure with the following dynamics [3,11]:

$$dS_t = (r - q)S_t d_t + \sigma S_t dW_t \tag{2.9}$$

<sup>&</sup>lt;sup>1</sup>Assumed to be constant.

<sup>&</sup>lt;sup>2</sup>See footnote 1.

Where  $dW_t$  is a Wiener process [9,11], r is the risk free rate and q is the dividend yield<sup>3</sup> and t denotes the current point in time.

Although the Black-Scholes is very popular and the *de facto* standard in the market there are implications to the B-S model assumptions that affect the results and that are unrealistic. The main assumption that does not hold up is the deterministic (constant) volatility, that can more accurately be described as a stochastic process since we observe that small moves usually are followed by small moves and large moves by large moves. [3,7]

Other assumptions that are critical to the B-S model and are not always observed in practice refer to the asset's continuity through time (no jumps), being allowed to perform continuous hedge without transactions costs and normal (Gaussian) returns.

Most models focus on the volatility problem because transaction costs often translate to rises in volatility and fat-tails (abnormal) returns can be simulated by stochastic volatility and market or volatility jumps.

### 2.3 Stochastic Volatility models

Introducing stochastic volatility to models brings complexity, but enables modeling some features observed in reality that are crucial like the randomic market volatility effects, skewness (market returns are more realistically modeled) and volatility smile. This kind of model is applied highly successfully in foreign exchange and credit markets.

**TODO:** Introduce CIR model

### Cox-Ingersoll-Ross (CIR) model

The Cox-Ingersoll-Ross (CIR) model is a well-known short-rate model that describes the interest rate movements driven by one source of market risk. The dynamics are described as follows:

$$dV_t = k(\theta - V_t)dt + \sigma\sqrt{V_t}dB_t \tag{2.10}$$

Where,  $V_t$  is the variance described by parameters k - the speed of mean reversion,  $\theta$  - the long-run mean variance and  $\sigma$  - the volatility of the variance process.

This model was widely used to describe the dynamics of the short rate interest because it has some fundamental features like intuitive parametrization, nonnegativity and pricing formulas. Furthermore, this equation and - this is the reason we present this model here - constitutes one of the two equations of the Heston model.

#### 2.3.1 Heston Model

Heston model solves the deterministic volatility problems introducing the following equations, which represents the dynamics of the stock price and the variance processes under the risk-neutral measure [12]:

 $<sup>^{3}</sup>r$  and q are assumed to be constant.

$$dS_t = \mu S_t dt + \rho \sqrt{V_t} dB_t + \sqrt{1 - \rho^2} \sqrt{V_t} S_t dW_t$$
  

$$dV_t = k(\theta - V_t) dt + \sigma \sqrt{V_t} dB_t$$
(2.11)

The second equation, as previously stated, is the CIR model equation. The first equation describes the asset price process.  $\mu$  is the asset's rate of return,  $dB_t$  and  $dW_t$  are two correlated whener processes with correlation coefficient of  $\rho$ .

$$dS_t = \mu S_t dt + \rho \sqrt{V_t} dB_t + \sqrt{1 - \rho^2} \sqrt{V_t} S_t dW_t$$
  

$$dV_t = k(\theta - V_t) dt + \sigma \sqrt{V_t} dB_t$$
(2.12)

[13]

#### 2.3.2 Other models

Models

This chunk ensures that the thesisdown package is

Chapter 5
Conclusion

Chapter 6
The First Appendix

### References

- [1] U.F. Wiersema, Brownian motion calculus, John Wiley & Sons, 2008.
- [2] I. Karatzas, S. Shreve, Brownian motion and stochastic calculus, Springer Science & Business Media, 2012.
  - [3] A.D. Helgadóttir, L. Ionescu, Option pricing within the heston model, (2016).
- [4] Z. Tong, Option pricing with long memory stochastic volatility models, PhD thesis, Université d'Ottawa/University of Ottawa, 2012.
- [5] L.C. Evans, An introduction to stochastic differential equations, American Mathematical Soc., 2012.
- [6] J.M. Steele, Stochastic calculus and financial applications, Springer Science & Business Media, 2012.
  - [7] Y. Yang, Valuing a european option with the heston model, (2013).
- [8] M. de F. Salomão, Precificação de opções financeiras: Um estudo sobre os modelos de black scholes e garch, (2011).
- [9] F. Black, M. Scholes, The pricing of options and corporate liabilities, Journal of Political Economy. 81 (1973) 637–654.
- [10] R.C. Merton, Theory of rational option pricing, The Bell Journal of Economics and Management Science. (1973) 141–183.
- [11] M. Gilli, D. Maringer, E. Schumann, Numerical methods and optimization in finance, Academic Press, Waltham, MA, USA, 2011. http://nmof.net.
- [12] M. Gilli, D. Maringer, E. Schumann, Numerical methods and optimization in finance, Academic Press, Waltham, MA, USA, 2011. http://nmof.net.
- [13] M. Broadie, Ö. Kaya, Exact simulation of stochastic volatility and other affine jump diffusion processes, Operations Research. 54 (2006) 217–231.