

# Numerical methods for the Heston model

---

School of Applied Mathematics

A Thesis

Presented to the

Getulio Vargas Foundation

---

In Partial Fulfillment

of the Requirements for the Degree

M.Sc. of Applied Mathematics

---

Fernando O. Teixeira

setembro 14, 2017



Approved for the Division  
(Mathematics)

---

Hugo Alexander de la Cruz Cancino



# Acknowledgements

Any one who considers arithmetical methods of producing random digits is, of course, in a state of sin. - John von Neumann

You get pseudo-order when you seek order; you only get a measure of order and control when you embrace randomness. — Nassim Nicholas Taleb

# Table of Contents

<b>Chapter 1: altadvisor: ‘Your Other Advisor’</b>	<b>1</b>
<b>Chapter 2: Literature Review</b>	<b>3</b>
2.1 Stochastic Calculus	3
2.1.1 The Stochastic differential equation - SDE	3
2.1.2 Brownian Motion	4
2.1.3 Itô’s Integral	6
2.2 Black-Scholes Model	7
2.2.1 Basics	7
2.2.2 The model	9
2.2.3 Limitations	10
2.3 Stochastic Volatility models	10
2.3.1 Cox-Ingersoll-Ross model	11
2.3.2 Heston Model	11
2.3.3 Other Models	12
2.4 Numerical Methods	12
2.4.1 Convergence	13
2.4.2 Discretization	14
2.4.3 Stability	16
<b>Chapter 3: The Heston Model Implementation</b>	<b>19</b>
<b>Chapter 4: This chunk ensures that the thesisdown package is</b>	<b>21</b>
<b>Chapter 5: Conclusion</b>	<b>23</b>
<b>Chapter 6: Black-Scholes formula</b>	<b>25</b>
<b>References</b>	<b>27</b>





# List of Tables

2.1	Box calculus . . . . .	7
-----	------------------------	---



# List of Figures

2.1	A Wiener process trajectory path example . . . . .	5
2.2	A GBM trajectory path example . . . . .	9
2.3	Volatility Smile . . . . .	11
2.4	Analytical x Euler solutions . . . . .	14
2.5	Euler's stability whith different timesteps . . . . .	17
2.6	Stability domains . . . . .	18



# Abstract

The preface pretty much says it all.

Second paragraph of abstract starts here.



# Dedication

You can have a dedication here if you wish.





# Chapter 1

**altadvisor: ‘Your Other Advisor’**



# Chapter 2

## Literature Review

This chapter presents the concepts of stochastic calculus, from the historic conception of how it first arose through the basic principles and applications in finance. We address with more care the classical Black-Scholes model and its limitations and the Heston model. This model is also well known, it brings the concept of stochastic volatility in it, which brings its results closer to reality.

### 2.1 Stochastic Calculus

Stochastic calculus arises from stochastic processes and allows the creation of a theory of integration where both the integrand and integrator terms are stochastic processes. Stochastic calculus was created by the Japanese mathematician Kiyosi Itô <sup>1</sup> in the 1940s and 1950s and is used for modeling financial options and in another wide variety of fields [27]. In this chapter we present the historical contexts in which the tools and models are used, but our focus is introducing the concepts and notations that will be further used in our work.

#### 2.1.1 The Stochastic differential equation - SDE

At first, before introducing stochastic differential equation, it is helpful to start with ordinary differential equation. Let  $x(t) = x_t$  denote a population at time  $t$  so that the change in the population at time  $t$  is given by the following deterministic differential equation:

$$\begin{aligned} dx_t &= f(t, x_t)dt \\ x(0) &= x_0 \end{aligned} \tag{2.1}$$

---

<sup>1</sup>There is another important stochastic integral, called the *Stratonovich Integral* that unlike the Itô's integral, respects the conventional calculus chain rule. Also, the integral is evaluated at the interval's midpoint, instead of its left extreme. A Stratonovich integral can be expressed as an Itô integral and vice versa.

We now add a “noise” to this equation:

$$\begin{aligned}
 dx_t &= \underbrace{f(t, x_t) dt}_{\text{drift}} + \underbrace{g(t, x_t) dW_t}_{\text{diffusion}} \\
 x(0) &= x_0
 \end{aligned} \tag{2.2}$$

This “noise”  $dW_t$  is a *random* Wiener process time derivative (which will be clarified below) and  $x_0$  is our initial value.

The  $g(t, x_t)$  part of the SDE is often referred as a *diffusion process*, these usually have a continuous paths. Before moving on, we must carefully define what the term *random* means and the best way to begin doing so is to precisely define a probability space:

**Definition 2.1.1.** (Probability Space) A triple  $(\Omega, \mathcal{U}, \mathcal{P})$  is called a *probability space* provided  $\Omega$  is any set,  $\mathcal{U}$  is a  $\sigma$ -algebra of subsets of  $\Omega$  and  $\mathcal{P}$  is a probability measure on  $\mathcal{U}$ .

## 2.1.2 Brownian Motion

The Brownian motion is the name given to the irregular motion observed in the motion of pollen particles suspended in fluid resulting from particle collision with atoms or molecules. It is named after Robert Brown, the first to have observed the movement in 1828. He noted two characteristic in the pollen movement [27]:

- the path of a given particle is very irregular, having a tangent at no point
- the motion of two distinct particles appear to be independent

The first quantitative works in Brownian motion come from an interest in stock price fluctuation by Bachelier in 1900. Albert Einstein also leaned over the subject and in 1905 derived the transition density for Brownian motion from molecular-kinetic theory of heat [16,27].

In 1923, the Wiener process was coined in honor of Norbert Wiener mathematical proof of existence of the Brownian motion and stating its properties.<sup>2</sup>

**Definition 2.1.2.** (Wiener Process) Given a probability space  $(\Omega, \mathcal{U}, \mathcal{P})$ , a stochastic process  $W_t$  defined in this space is a *\*Wiener process\** if it satisfies the following properties:

- $W_0 = 0$
- The change in  $W$ , given by  $\Delta W = W_{t+1} - W_t$ , is normally distributed with mean zero and standard deviation  $\sqrt{\Delta t}$ , meaning that  $\Delta W = \epsilon \sqrt{\Delta t}$ , where  $\epsilon$  is  $N(0, 1)$ .

---

<sup>2</sup>More can be found on [5,9,22].

- If the increment  $\Delta t_1$  does not overlap with the time increment  $\Delta t_2$ , then  $\Delta W_1$  and  $\Delta W_2$  are independent.
- The process is continuous, meaning that there are no jumps in the path.
- The process is Markovian. This means that the conditional expectation of  $W_{t+1}$  given its entire history is equal to the conditional expectation of  $W_{t+1}$  given today's information. This can be written as:  $E[W_{t+1}|W_1, \dots, W_t] = E[W_{t+1}|W_t]$ .
- Consider the time interval  $[0, t]$  with  $n$  equally spaced intervals given by  $t_i = \frac{it}{n}$ . Then the paths of the Brownian motion have unbounded variation, this means that they are not differentiable and go towards infinity as  $n$  increases. The quadratic variation is given by  $\sum_{i=1}^n (Z_{t_i} - Z_{t_{i-1}})^2 \rightarrow t$ , meaning that when  $n$  increases it stays constant at  $t$ .




Figure 2.1: A Wiener process trajectory path example

### Correlated Brownian Motions

Two independent Brownian motions that are correlated can describe a new process  $Z_t$ . Let  $W_1$  and  $W_2$  be these two *independent* Brownian motions and let  $-1 \leq \rho \leq 1$  be a given number. For  $0 \leq t \leq T$  define the new process  $Z_t$  as [27]:

$$Z_t = \rho W_{1,t} + \sqrt{1 - \rho^2} W_{2,t} \quad (2.3)$$

This equation is a linear combination of independent normals at each timestep  $t$ , so  $Z_t$  is normally distributed. It is proven that  $Z$  is a Brownian motion and that  $Z$  and  $W_{1,t}$  have correlation  $\rho$  [27].

### 2.1.3 Itô's Integral

Formally, the SDE presented in equation (2.2) only exists because we can rewrite it in the form [5,14,15,22,24,25]:

$$x_t = x_0 + \int_0^t f(s, x_s)ds + \int_0^t g(s, x_s)dW_s \quad (2.4)$$

for some  $f(s, x_s)$ ,  $g(s, x_s)$  and  $s \in [0, t]$ .

The Itô integral can, as the Riemann integral, be approximated by a finite sum. Also, it has a definition as a certain limit. Itô's lemma 2.1.1 plays the same role as the fundamental theorem of calculus in allowing to evaluate integrals. It is the formal definition and presents an extra term not encountered in the conventional calculus theorem that is due to the non-smoothness characteristics of Brownian motion paths. It is possible, though, to define the integral in a less rigorous way:

$$Y_{\Delta t}(t) \approx \sum_{t_k < t} g(t_k) \Delta W_k \quad (2.5)$$

with the usual notions  $t_k = k\Delta t$ , and  $\Delta W_k = W(t_{k+1}) - W(t_k)$ . And in a more rigorous form, if the limit exists, then the Ito integral is:

$$Y(t) = \lim_{\Delta t \rightarrow 0} Y_{\Delta t}(t) \quad (2.6)$$

It is essential that the *forward difference* is used rather than the backward difference, which would be wrong.

**Theorem 2.1.1** (Itô's Lemma). *Assume that  $S_t$  has a stochastic differential given by:*

$$dS_t = \mu_t dt + \sigma_t dW_t \quad (2.7)$$

*for  $\mu_t$ ,  $\sigma_t$  and  $t \in [0, T]$ . Assume  $u : \mathbb{R} \times [0, T] \rightarrow \mathbb{R}$  is continuous and that  $\frac{\partial u}{\partial t}$ ,  $\frac{\partial u}{\partial x}$ ,  $\frac{\partial^2 u}{\partial x^2}$  exist and are continuous.*

$$Y_t := u(S_t, t)$$

*Then  $Y$  has the following stochastic differential:*

$$\begin{aligned} dY_t &= \frac{\partial u}{\partial t} dt + \frac{\partial u}{\partial x} dS_t + \frac{1}{2} \frac{\partial^2 u}{\partial x^2} \sigma_t^2 dt \\ &= \left( \frac{\partial u}{\partial t} + \mu_t \frac{\partial u}{\partial x} + \frac{1}{2} \frac{\partial^2 u}{\partial x^2} \sigma_t^2 \right) dt + \sigma_t \frac{\partial u}{\partial x} dW_t \end{aligned} \quad (2.8)$$

*where the argument of  $u$ ,  $\frac{\partial u}{\partial x}$  and  $\frac{\partial^2 u}{\partial x^2}$  above is  $(S_t, t)$ .*

Equation (2.8) is the stochastic equivalent to the chain rule, also known as Itô's formula or Itô's chain rule. The proof to this theorem is based on the Taylor expansion of the function  $f(S_t, t)$  [5,25]. For practical use you should write out a second-order Taylor expansion for the function to be analyzed and apply the Table 2.1 multiplication table [27].

Table 2.1: Box calculus

	$dt$	$dW_t$
$dt$	0	0
$dW_t$	0	$dt$

### Itô's Integral Properties

Let  $f, g \in \mathcal{V}$  and let  $0 \leq t_0 < u < T$ . Then

$$(i) \int_{t_0}^T f dB_t = \int_{t_0}^u f dB_t + \int_u^T f dB_t$$

$$(ii) \int_{t_0}^T (\alpha f + \beta g) dB_t = \alpha \int_{t_0}^T f dB_t + \int_{t_0}^T \beta g dB_t$$

$$(iii) \mathbb{E} \left[ \int_{t_0}^T f dB_t \right] = 0$$

$$(iv) \mathbb{E} \left[ \left( \int_0^t H_s dB_s \right)^2 \right] = \mathbb{E} \left[ \int_0^t H_s^2 ds \right] \text{ (Isometry)}$$

$$(v) \mathbb{E} \left[ \int_{t_0}^T f dB_t \mid \mathcal{F}_s \right] = \int_{t_0}^s f dB_t, \quad \text{for } s < T. \text{ (Martingale}^3\text{)}$$

## 2.2 Black-Scholes Model

### 2.2.1 Basics

The Black-Scholes (B-S) model arises from the need to price european options in the derivative markets. Derivatives are financial instruments traded in the market, stock exchange or over-the-counter (OTC) market, whose values depend on the values of an underlying asset. [1,23,28]

- A call option is a derivative that gives its bearer the right, but not the obligation, to purchase a specific asset by a fixed price before or on a given date.
- A put option is a derivative that gives its bearer the right, but not the obligation, to sell a specific asset by a fixed price before or on a given date.

The trading price of the option is called the option *premium* and the asset from which the option derives is called the *underlying asset*. This asset may be interest rates, exchange rates, stock exchanges indices, commodities or stocks. The fixed price in

---

<sup>3</sup>A martingale is a stochastic process with certain characteristics. The main one is that the expected value in time  $t + 1$  for  $X$  is the  $X$  value in  $t$ . This means there are no winning strategies when we are dealing with martingales (unlike when we play poker, for example). A Wiener process is a martingale.

contract in which the underlying asset might to be bought or sold is the *strick price*. The option expiration date is called the *maturity*. [1,23]

There are two major different option types: European and American. The difference between these two is that the bearer of the first may exercise it only at the end of its life, at its maturity while the latter can be exercised at any given time until its maturity. [1,20]

**Definition 2.2.1.** (Intrinsic value) The intrinsic value of a call is the difference between the underlying asset price and the strike price. The put's intrinsic value operates the other way around, being the difference between the strike and the underling asset prices.

## Geometric Brownian Motion

A stochastic process  $S_t$  is a Geometric Brownian Motion<sup>4</sup> if it is described by the solution of the following stochastic differential equation [25–27].

$$dS_t = \mu S_t dt + \sigma S_t dW_t \quad (2.9)$$

for given constants  $\mu \in \mathbb{R}$  and  $\sigma > 0$ . Also, the assumed initial value is positive,  $S_0 > 0$ .

Figure 2.2 shows the GBM,<sup>5</sup> which is quite often applied to model the dynamics of some financial assets because of its properties [12]. Equation (2.10) shows the formula to generate a GBM and we provide proof of this solution in appendix 6<sup>6</sup>

$$S_t = S_0 \times \exp\left(\left(\mu - \frac{\sigma^2}{2}\right)t + \sigma W_t\right), \quad t > 0 \quad (2.10)$$

---

<sup>4</sup>There is an Arithmetic Brownian Motion:  $dS_t = \mu dt + \sigma dB_t$ . More information can be obtained at [27].

<sup>5</sup>Also known as exponential Brownian motion.

<sup>6</sup>An intuitive proof can be found at [18].






Figure 2.2: A GBM trajectory path example

### 2.2.2 The model

The Black-Scholes model provides analytical solution to the price of a European call at time  $t$  and can be described as follows [1,9,28]:

$$C(S_t, t) = N(d_1)S_t - N(d_2)Ke^{-r(T-t)} \quad (2.11)$$

$$d_1 = \frac{1}{\sigma\sqrt{T-t}} \left[ \ln\left(\frac{S_t}{K}\right) + \left(r + \frac{\sigma^2}{2}\right)(T-t) \right] \quad (2.12)$$

$$d_2 = d_1 - \sigma\sqrt{T-t} \quad (2.13)$$

Where:

- $S_t$  is the spot price of the underlying asset at time  $t$
- $r$  is the risk free rate (generally an annual rate)<sup>7</sup>
- $\sigma$  is the volatility of returns of the underlying asset<sup>8</sup>
- $N(\cdot)$  is the cumulative distribution function of the standard Gaussian distribution
- $K$  is the strike price
- $T - t$  is the time to maturity

Also, the stock price path is a Geometric Brownian Motion as previously stated, and is under the risk-neutral measure with the following dynamics [7,9]:

$$dS_t = (r - q)S_t dt + \sigma S_t dW_t \quad (2.14)$$

---

<sup>7</sup>Assumed to be constant.

<sup>8</sup>See footnote 1.

Where  $dW_t$  is a Wiener process [1,7],  $r$  is the risk free rate and  $q$  is the dividend yield<sup>9</sup> and  $t$  denotes the current point in time.

### 2.2.3 Limitations

Although the Black-Scholes is very popular and the *de facto* standard in the market there are implications to the B-S model assumptions that affect the results and that are unrealistic. The main assumption that does not hold up is the deterministic (constant) volatility, that can more accurately be described as a stochastic process since we observe that small moves usually are followed by small moves and large moves by large moves. [9,28]

Other assumptions that are critical to the B-S model and are not always observed in practice refer to the asset's continuity through time (no jumps), being allowed to perform continuous hedge without transactions costs and normal (Gaussian) returns.

Most models focus on the volatility problem because transaction costs often translate to rises in volatility and fat-tails (abnormal) returns can be simulated by stochastic volatility and market or volatility jumps.

## 2.3 Stochastic Volatility models

Introducing stochastic volatility to models brings complexity, but enables modeling some features observed in reality that are crucial, like the random market volatility effects, skewness (market returns are more realistically modeled) and volatility smile<sup>10</sup> (see Figure 2.3). This kind of model is applied highly successfully in foreign exchange and credit markets.

**Definition 2.3.1.** (Volatility Smile) Volatility smiles are implied volatility patterns that arise in pricing financial options. In particular for a given expiration, options whose strike price differs substantially from the underlying asset's price command higher prices (and thus implied volatilities) than what is suggested by standard option pricing models. These options are said to be either deep in-the-money or out-of-the-money.

Furthermore, stochastic volatility models use statistical methods as foundations to price and forecast options' behaviors and the underlying's security volatility is arbitrary. The Heston, the 3/2 and other models, like the GARCH<sup>11</sup> and SABR,<sup>12</sup> are considered standard smile models.

---

<sup>9</sup> $r$  and  $q$  are assumed to be constant.

<sup>10</sup>The name derives from the concave shape of the graph, which resembles a smile.

<sup>11</sup>generalized autoregressive conditional heteroscedasticity.

<sup>12</sup>stochastic alpha, beta, rho.




Figure 2.3: Volatility Smile

### 2.3.1 Cox-Ingersoll-Ross model

The Cox-Ingersoll-Ross (CIR) model is a well-known short-rate model that describes the interest rate movements driven by one source of market risk. The dynamics are described as follows[4,10]:

$$dr_t = k(\theta - r_t)dt + \sigma\sqrt{r_t}dB_t \quad (2.15)$$

Where,  $r_t$  is the short rate interest described by parameters  $\kappa$  the speed of mean reversion,  $\theta$  the long-run mean variance and  $\sigma$  the volatility of the variance process.

This model has been widely used to describe the dynamics of the short rate interest because it has some fundamental features like intuitive parametrization, nonnegativity and pricing formulas. Besides, it takes account of anticipations, risk aversion, investment alternatives and preferences about consumption timing and allows for detailed predictions about how changes in a wide range of underlying variables affect the term structure[4]. Furthermore, this equation constitutes one of the two Heston model equations with the volatility taking the short rate interest place.

### 2.3.2 Heston Model

Heston model was introduced in 1993 by Steven Heston to solve the deterministic volatility problems. It was designed to analyze bond and currency options and it introduced the following equations, which represent the dynamics of the stock price

and the variance processes under the risk-neutral measure [8,10]:

$$dS_t = \mu S_t dt + \sqrt{V_t} S_t dW_t^* \quad (2.16)$$

$$dV_t = \kappa(\theta - V_t)dt + \sigma\sqrt{V_t}dB_t \quad (2.17)$$

The second equation, as described in Section 2.3.1, is the CIR model equation. The first equation states the asset price process.  $\mu$  is the asset's rate of return,  $dW_{t,1}$  and  $dW_{t,2}$  are two correlated wiener processes with correlation coefficient of  $\rho$ .

### 2.3.3 Other Models

#### Ornstein-Uhlenbeck

The Ornstein-Uhlenbeck is the earliest recorded SDE. Named after Leonard Ornstein and George Eugene Uhlenbeck, it is a stochastic process that describes the acceleration of a pollen particle in a liquid subject to bombardments by molecules [27]. As we can observe in equation (2.18),  $x_t$  represents the one dimension velocity of the particle, thus  $dx_t$  is the *change* in velocity, in other words, its acceleration. The  $-\theta x_t$  component slows down the acceleration and is to be understood as frictional force. Besides, we add a noise  $W_t$  with intensity  $\sigma$  that models the random bombardment by the molecules.

$$dx_t = -\theta x_t dt + \sigma dW_t \quad (2.18)$$

With  $\theta$  and  $\sigma$  being positive constants. Expressing in terms of  $x_t$  we get:

$$x_t = e^{-\theta t} \times \left[ x_0 + \sigma \int_{t=0}^T e^{\theta t} dW_s \right]. \quad (2.19)$$

#### Langevin

The Langevin equation describes a system that consists of the molecular bombardment of a speck of dust on a water surface. We know that the intensity of the bombardement does not depend on the state variables [6,17].

$$m \frac{dv}{dt} = -\zeta v + \delta F(t) \quad (2.20)$$

$m$  is the mass of the particle,  $v$  it's velocity,  $-\zeta v$  is the frictional force, which is proportional to the velocity, and  $\delta F(t)$  is a *fluctuating* force (random) to the frictional force.

## 2.4 Numerical Methods

Numerical methods are tools that are often applied to solve stochastic differential equations because most of these do not have explicit solution. This means that we are not able to solve these equations using symbolic computation. Although we are

unable to find an analytical solution, when facing real problems, the approximation given by a numerical method is often sufficient. Alongside the analytical issue, the need to calculate the SDE's trajectory through time is the main reason why studying numerical methods is so important. An implementation of a numerical method is called a numerical algorithm.

We will simulate sample paths of time discrete approximations implemented in the R programming language [21] that we base on a finite discretization of a time interval  $[t_0, T]$ . We shall generate approximate values of the sample path for each step contained in the discretized interval [17].

In the fixed step methods, the distance between two contiguous points is the distance  $d_i = t_i - t_{i-1} = \frac{T-t_0}{N} \quad \forall i \mid 1 \leq i \leq N \in \mathbb{N}$ .  $N$  being the time interval partition number.

According to Kloeden [17], in the stochastic realm, simulated sample paths can be statistically analysed to find how good an approximation is compared to an exact solution. Moreover, the computational costs such as time and memory increases polynomially with the problem's dimension, which is good, and it is possible to apply variance reduction methods that allow a considerable decrease in the required sample size.

### 2.4.1 Convergence

As soon as we talk about numerical methods we are required to approach the topic of approximations and how to handle them. Methods efficiency receive the name of *convergence order*. In the SDE domain there are two main methods of convergency, that are classified according to their criteria. Firstly, we present the *strong order of convergence*. A method is said to have strong convergence  $\delta$  to  $Y$  if a time discretized  $Y_\delta$  of a continuous-time process  $Y$ , with  $\delta$  being the maximum time increment of the discretization, and for any fixed time horizon  $T$  holds true that [12]:

$$\mathbb{E} \mid Y_\delta(T) - Y(T) \mid \leq C\delta^\gamma, \quad \forall \delta < \delta_0$$

with  $\delta_0 > 0$  and  $C$  a constant not depending on  $\delta$ . Strong convergence addresses the problem of solutions' trajectories. For specific conditions, the Euler method has strong convergence order  $\gamma = \frac{1}{2}$ . Furthermore, there is the *weak order of convergence*. The weak convergence

$$\mid \mathbb{E}p(Y_n) - \mathbb{E}p(Y(\tau)) \mid \leq C\Delta t^\gamma$$

Strong and weak convergence are not mutually exclusive [12]. That means that a method with a given strong order of convergence might have a higher weak order of convergence too. This is the case for the Euler scheme, with a strong order of convergence of  $1/2$  and a weak order of  $1$  (under some conditions). For a more detailed and rigorous explanation of convergence we recommend consulting [11].

It is worth noting that, although schemes have a given convergency order, it is not unusual that they behave better than their order for some SDEs specifications.

### 2.4.2 Discretization

We know that convergence is an important feature to a numerical method and studies have found not all time discrete possible approximations of an SDE converge in a useful sense to the solution process as the step size adopted tends toward zero [2,3]. Moreover, particularly for SDEs, some of the more rapidly convergent methods available for ordinary differential equations (ODE) do not work, such as higher order Runge-Kutta methods.<sup>13</sup>

One of the methods that do work for ODEs and SDEs is the Euler method, named after the Swiss mathematician Leonhard Euler. Figure 2.4 shows an example of an implementation for the Newton's cooling law with timestep of 2 seconds compared to its analytical solution. This method (*a.k.a.* forward Euler method) is a first-order numerical procedure. It is the most basic explicit method<sup>14</sup> for numerical integration.




Figure 2.4: Analytical x Euler solutions

The method is first-order, as stated above, this means that the error in each step is a proportion of the square of the step size. Also, the global error at a given time is a function of the step size. We proceed to apply the Euler method to SDEs. Consider the equation:

$$dS_t = \mu(S_t, t)dt + \sigma(S_t, t)dW_t \quad (2.21)$$

$dW_t$  is the Brownian motion,  $\mu$  and  $\sigma$  are functions depending on  $S_t$  and  $t$ , over an

<sup>13</sup>The euler method is the simplest Runge-Kutta method.

<sup>14</sup>Explicit methods calculate the state of a system at a later time from the state of the system at the current time. Mathematically we have something like  $Y(t + \Delta t) = F(Y(t))$ .

interval  $[0, T]$ , and we want to discretize it as  $0 = t_1 < t_2 < \dots < t_m = T$  with increments equally spaced  $d_t$ .

Integrating it from  $t$  to  $t+dt$  we have the starting point for our (and any) discretization scheme:

$$S_{t+dt} = S_t + \int_t^{t+dt} \mu(S_u, u) du + \int_t^{t+dt} \sigma(S_u, u) dW_u \quad (2.22)$$

To use the Euler discretization is the equivalent of approximating integrals using the left-point rule, we then have:

$$\begin{aligned} \int_t^{t+dt} \mu(S_u, u) dW_u &\approx \mu(S_t, t) \int_t^{t+dt} dW_u \\ &= \mu(S_t, t)(W_{t+dt} - W_t) \\ \int_t^{t+dt} \sigma(S_u, u) dW_u &\approx \sigma(S_t, t) \int_t^{t+dt} dW_u \\ &= \sigma(S_t, t)(W_{t+dt} - W_t) \\ &= \sigma(S_t, t)\sqrt{dt}Z \end{aligned}$$

$W_{t+dt} - W_t$  and  $\sqrt{dt}Z$  have identical distribution,  $Z$  being a standard gaussian variable. The Euler discretization of equation (2.22) is then:

$$S_{t+dt} = S_t + \mu(S_t, t)dt + \sigma(S_t, t)\sqrt{dt}Z \quad (2.23)$$

### Euler method - Heston model

We now proceed to apply the method to our model of interest. We retake the equations (2.16) and (2.17). We begin showing how to discretize the latter [11,12]:

$$V_{t+dt} = V_t + \int_t^{t+dt} \kappa(\theta - V_u) du + \int_t^{t+dt} \sigma\sqrt{V_u} dB_u \quad (2.24)$$

Which discretized turns out as:

$$\begin{aligned} \int_t^{t+dt} \kappa(\theta - V_u) du &\approx \kappa(\theta - V_t)dt \\ \int_t^{t+dt} \sigma\sqrt{V_u} dB_u &\approx \sigma\sqrt{V_t}(W_{t+dt} - W_t) \\ &= \sigma\sqrt{V_t dt}Z_v \end{aligned}$$

And leaves us with:

$$V_{t+dt} = V_t + \kappa(\theta - V_t)dt + \sigma\sqrt{V_t dt}Z_v \quad (2.25)$$

$Z_v$  is a standard normal variable. To avoid problems with negative values in  $\sqrt{V_t}$  we apply the *full truncation* scheme, which substitutes  $V_t$  with  $V_t^+ = \max(0, V_t)$ .<sup>15</sup>

---

<sup>15</sup>Another possible scheme (not used in this work) is the *reflection* scheme where we replace  $V_t$  with  $|V_t|$

For the  $S_t$  SDE we proceed similarly:

$$S_{t+dt} = S_t + \mu \int_t^{t+dt} S_u du + \int_t^{t+dt} \sqrt{V_u} S_u dW_u \quad (2.26)$$

Discretizing we have:

$$\begin{aligned} \int_t^{t+dt} S_u du &\approx S_t dt \\ \int_t^{t+dt} \sqrt{V_u} S_u dW_u &\approx \sqrt{V_t} S_t (W_{t+dt} - W_t) \\ &= \sqrt{V_t dt} S_t Z_s \end{aligned}$$

$Z_s$  is a standard normal variable with correlation  $\rho$  with  $Z_v$ . We have:

$$S_{t+dt} = S_t + \mu S_t dt + \sqrt{V_t dt} S_t Z_s \quad (2.27)$$

### 2.4.3 Stability

Most differential equations, deterministic or stochastic, cannot be solved explicitly [17]. Hence, stability studies begin with computers and is associated with numerical methods and approximations. Convergent methods were resulting in bigger errors than what was expected that could not be only due to discretization error. Eventually, scientists discovered that this unexpected problem was caused by accumulation of successive truncation errors. Figure 2.5 retakes the cooling example previously approached to show instability due to an increase in size of the timestep and extending to 600 seconds.






Figure 2.5: Euler's stability with different timesteps

We know that binary machines like computers are not able to represent all the real numbers, but only a subset of them. Thus, solving these errors is not straightforward since it's not possible to eliminate *all* truncation error when using a computer and dealing with numerical solutions. When faced to an incorrect (not acceptable) solution, we have to evaluate and distinguish between two distinct situations:

- i Rounding errors are considerably amplified by the algorithm. This situation is called numerical instability.
- ii Small perturbations of data generate large changes in the solution. This is termed an ill-conditioned (or sensitive) problem.

Examples of these two classes of problem can be found in [8].

Stiff equations appear very often in mathematical problems and refer to differential equations for which a numerical methods might be unstable for not small enough stepsizes [13]. A differential equation of the form  $y' = f(t, y)$ , if its exact solution  $y(t)$  includes a term that decays exponentially to zero as  $t$  increases, but whose derivatives are greater in magnitude than the term itself. In other words, if it requires a significant

depression of the stepsize to avoid stability lost. This is a loose definition but, since we are dealing with numerical methods a proper mathematical definition isn't required. An example of the stiff behavior is the following differential equation [19]:

$$y' = -100y, \quad t > 0, \quad y_0 = 1$$

Whose exact solution is  $y_t = e^{-100t}$  and goes to zero as  $t$  increases. Applying Euler's method to this equation with  $h = 0.1$  we stumble in the following equation

$$y_{n+1} = y_n - 100hy_n = -9y_n$$

which is wrong, since it yields an exponentially growing solution  $y_n = (-9)^n$ . On the other hand, if our timestep is smaller  $h = 10^{-3}$ , our solution using Euler's method becomes  $y_n = (0.9)^n$ . This solution leads to an accurate behavior regarding the exact solution, it rapidly decays to zero.

Sometimes, it is interesting to rank differential equations that are more or less stiff. Thus, people compute the quotient of the largest and the smallest eigenvalues of a linear system. They call it the equations' *stiffness ratio* and, usually the bigger the stiffness ratio, the more likely is to be stiff [13].




Figure 2.6: Stability domains

Thereby, we claim that stability method study is an important topic, since it enables achieving solutions that are good to stiff equations without having to overly reduce our timesteps which can be very computationally costly.

## **Chapter 3**

# **The Heston Model Implementation**



## Chapter 4

This chunk ensures that the  
thesisdown package is



## Chapter 5

## Conclusion





## Chapter 6

### Black-Scholes formula



# References

- [1] F. Black, M. Scholes, The pricing of options and corporate liabilities, *Journal of Political Economy*. 81 (1973) 637–654.
- [2] J.M. Clark, R. Cameron, The maximum rate of convergence of discrete approximations for stochastic differential equations, in: *Stochastic Differential Systems Filtering and Control*, Springer, 1980: pp. 162–171.
- [3] D. Clements, B. Anderson, Well-behaved itô equations with simulations that always misbehave, *IEEE Transactions on Automatic Control*. 18 (1973) 676–677.
- [4] J.C. Cox, J.E. Ingersoll Jr, S.A. Ross, A theory of the term structure of interest rates, *Econometrica: Journal of the Econometric Society*. (1985) 385–407.
- [5] L.C. Evans, *An introduction to stochastic differential equations*, American Mathematical Soc., 2012.
- [6] T.C. Gard, *Introduction to stochastic differential equations*, M. Dekker, 1988.
- [7] M. Gilli, D. Maringer, E. Schumann, *Numerical methods and optimization in finance*, Academic Press, Waltham, MA, USA, 2011.
- [8] M. Gilli, D. Maringer, E. Schumann, *Numerical methods and optimization in finance*, Academic Press, 2011.
- [9] A.D. Helgadóttir, L. Ionescu, Option pricing within the heston model, (2016).
- [10] S.L. Heston, A closed-form solution for options with stochastic volatility with applications to bond and currency options, *Review of Financial Studies*. 6 (1993) 327–343.
- [11] D.J. Higham, An algorithmic introduction to numerical simulation of stochastic differential equations, *SIAM Review*. 43 (2001) 525–546.
- [12] S.M. Iacus, *Simulation and inference for stochastic differential equations: With r examples*, Springer Science & Business Media, 2009.
- [13] A. Iserles, *A first course in the numerical analysis of differential equations*, Cambridge university press, 2009.
- [14] K. Itô, The brownian motion and tensor fields on riemannian manifold, *Proc. Int. Congr. Math.*, Stockholm. 2 (1962).
- [15] K. Itô, On stochastic differential equations, *American Mathematical Soc.*, 1951.
- [16] I. Karatzas, S. Shreve, *Brownian motion and stochastic calculus*, Springer Science & Business Media, 2012.
- [17] P.E. Kloeden, E. Platen, *Numerical solution of stochastic differential equations* springer-verlag, 1992.
- [18] A. Krouglov, Intuitive proof of black-scholes formula based on arbitrage and

properties of lognormal distribution, arXiv Preprint Physics/0612022. (2006).

[19] J.V. Lambers, A.C. Sumner, Explorations in numerical analysis, University O California at Irvine. (2016).

[20] R.C. Merton, Theory of rational option pricing, The Bell Journal of Economics and Management Science. (1973) 141–183.

[21] R Core Team, R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria, 2017.

[22] J.S. Rosenthal, A first look at rigorous probability theory, World Scientific Publishing Co Inc, 2006.

[23] M. de F. Salomão, Precificação de opções financeiras: Um estudo sobre os modelos de black scholes e garch, (2011).

[24] J.M. Steele, Stochastic calculus and financial applications, Springer Science & Business Media, 2012.

[25] Z. Tong, Option pricing with long memory stochastic volatility models, PhD thesis, Université d'Ottawa/University of Ottawa, 2012.

[26] R.S. Tsay, Analysis of financial time series, John Wiley & Sons, 2005.

[27] U.F. Wiersema, Brownian motion calculus, John Wiley & Sons, 2008.

[28] Y. Yang, Valuing a european option with the heston model, (2013).