

# Joint analysis of geological map units and topography to support soil survey - lessons from a case study in South Tyrol OR From geological to soil parent material maps - a random forest-supported analysis of geological map units and topography in South Tyrol

Fabian E. Gruber<sup>a,\*</sup>, Jasmin Baruck<sup>a</sup>, Clemens Geitner<sup>a</sup>

<sup>a</sup>*Institute of Geography, University of Innsbruck, Innrain 52f, 6020 Innsbruck, Austria*

---

## Abstract

*Keywords:*

---

## 1. Introduction

*general introduction.* Geologic maps have always been an important aid in soil survey as parent material is a decisive factor in soil formation (Jenny, 1941). The importance of this relationship is highlighted by the fact that, vice versa, soil maps have themselves been applied to support and improve geologic mapping (Brevik and Miller, 2015).

Geologic maps have always been an important aid in soil survey as parent material is a decisive factor in soil formation (Jenny, 1941). The importance of this relationship is highlighted by the fact that, vice versa, soil maps have themselves been applied to support and improve geologic mapping (Brevik and Miller, 2015). Providing both the physical structure and the chemical composition of the mineral constituents, parent material plays a fundamental role regarding the direction as well as speed of soil evolution. This is particularly the case in young soils (e.g. Schaetzl et al. (2000)) such as those predominantly found in the Alps (Geitner et al., 2017). Thus, in order to understand the spatial pattern of soils in the Alps, it is essential to identify

---

\*Corresponding author

*Email address:* `Fabian.Grubert@uibk.ac.at` (Fabian E. Gruber)

the types and origins of parent materials, which are, at least in the lower and medium elevations of the Alpine environment, dominated by quaternary unconsolidated sediments. These deposits vary considerably in thickness; they are often multi-layered and exposed to recent morphodynamics, all of which control soil horizon development and properties (Phillips and Lorz, 2008). In this context, it is indicated to include characteristics of the subsolum as often as possible, mainly in order to make soil information more suitable for a wide range of environmental issues, as discussed in detail by Juilleret et al. (2016). Consequently, geological maps at various scales have been used as an environmental variable in digital soil mapping (DSM), representing the soil forming factor parent material, or simply 'p'. In their study which presents the 'scorpan' framework of inferring soil information, McBratney et al. (2003) present a table of studies applying DSM, which also indicates in which of these studies the parent material was involved as an independent variable. How this important variable is classified, however, will vary greatly depending on the the available data, the soil classification system used, the specific mapping guidelines applied, and most importantly the particular geologic and geomorphological setting of the investigated area. In its guidelines for soil description, the Food and Agriculture Organization of the United Nations promotes a hierarchical system for describing lithologies that constitute the soil parent material, based on the major classes igneous rock, metamorphic rock, consolidated and unconsolidated sedimentary rock (FAO, 2006). KA5? While the lithologies regarding bedrock as parent material are similar to the types in the classification system used by the surveyors employed by the Forestry service, the latter system is closely adapted to the Alpine environment. Specifically, the major class of unconsolidated sedimentary rocks has a far greater number of types in order to satisfy the demands posed by the diversity of the glacial, but also the more recent deposits, driven mainly by the high relief present in Alpine regions.

While such an adaptation of the classes and types of soil parent material to the given circumstances is certainly necessary, communication between soil scientists regarding soil parent materials and comparability is hindered by the multitude of classifications. Juilleret et al. (2016), who stress the importance of describing the subsolum in soil survey, propose a morphogenetic procedure for characterising and classifying subsolum material applying a structure similar to that of the WRB.

Herbst zitat fr bedeutung der Schrfte der Geologischen Karte fr die genauigkeit von DSM??

55 A number of studies have compared the information from soil surveys  
with geologic maps. HERE SOME MORE LITERATURE, like Miller and  
Lee Burras (2015), Juilleret(2012), Brevik and Miller (2015). While most of  
the previously mentioned studies analyse the possibility of using soil survey  
information for mapping surficial geology, the aim of the presented study is  
60 to highlight those geologic units of the study area where the soil parent ma-  
terial cannot be simply derived from the detailed geologic map. McBratney  
et al. (2003) list some examples of DSM studies which use geologic maps as  
environmental variables.

Situation in the Alps bezüglich boden und geologie wäre auch noch in-  
65 teressant

A second, immensely important soil forming factor is topography or relief.  
It is considered in traditional soil survey, for instance by mapping landscape  
position and local slope and curvature (FAO, 2006), and also DSM, where  
the representing variables implemented in a given model can be chosen from  
70 a wide set of available parameters. Examples of such terrain parameters can  
be found, amongst others, in Böhner and Antonić (2009), Gallant and Wilson  
(2000) and Olaya (2009) . Regarding the geomorphometric characterisation  
of geologic or soil parent material units, a number of considerations have to  
be taken into account when choosing which parameter groups to investigate.  
75 While regional parameters well describe the, hydrologically relevant, relative  
position in the landscape, they, as well as absolute and relative height-related  
parameters, are strongly correlated to the underlying geological structure of  
a given region. Local parameters such as slope and curvature are often used  
to infer soil properties and give insight into local dynamics, but may also  
80 vary strongly within a map unit. To characterise parent material units,  
especially with regard to topographic, and as a result, soil, variability, an in-  
termediate terrain parameter describing a unit's land surface is of particular  
interest. Researchers have long investigated ways to quantify the roughness  
or ruggedness of terrain, from the analysis of field data and topographic maps  
85 to computing roughness indices on raster grids. Geology, geomorphology as  
well as habitat modelling and wildlife management have been the main sci-  
entific research areas in which such investigations were performed on land  
surfaces. Hobson (1972) presented three different roughness values and ap-  
plied them to field measurements, correlating them to rock type. In another  
90 early study aimed at quantifying roughness, Beasom et al. (1983) presented  
the land surface ruggedness index, which is based on the total length of con-  
tour lines per area. Similarly analysing topographic maps, Nellemann and

Thomsen (1994) describe the calculation of a terrain ruggedness index based on the variability of contour lines along transects, which they correlate with caribou forage availability. Regarding field methods, they calculate micro-topographic diversity by analysing the horizontal distance of a chain laid on the ground in their study plots. Riley et al. (1999) proposed the topographic ruggedness index (TRI), which compares the elevation of a central pixel to the elevations of cells within a given search window. In an attempt to decorrelate roughness from slope, Sappington et al. (2007) expanded on the work of Hobson (1972) to introduce vector ruggedness measure (VRM), which is calculated based on the orientation of vectors normal to the surface in a given area. (Grohmann et al., 2010) analysed several roughness measures at different resolutions and window sizes with regard to their ability to depict terrain features. They highlight the ability of VRM to detect fine-scale roughness features and attribute low roughness values to steep but smooth slopes, but also acknowledge its inability to delimit slope breaks and identify regional relief. The Melton ruggedness index, which relates the elevation difference of a basin to the drainage area, was applied by Marchi and Dalla Fontana (2005) to investigate sediment transport, however compared to VRM it is more of a measure of general relief than roughness. Similar to VRM, roughness measures based on eigenvalue ratios of an orientation matrix have been used in geology to describe land surfaces, especially bedrock fabric. Coblenz et al. (2014) combined such a roughness measures with parameters representing the drainage network of the investigated geologic units to create terrain characterisation types to distinguish various lithologies, with emphasis on discriminating soft and hard rock areas.

*Intention and aims.* The objective of this study is to evaluate how to make best use of available geologic and topographic information (with emphasis on terrain roughness measures) for soils survey. By applying random forest classification and feature selection, we investigate which terrain parameters, with emphasis on roughness measures, are best suited to produce a parent material map based on an available geological maps as well as topography. Additionally, the same method is applied to distinguish terrain parameters that, for each soil parent material class, best separate those profile site points that are correctly classified in the geological map from those of the same class that are misclassified. Based on this analysis and a similar investigation into characteristic terrain parameters of the geological map units, each of these is characterised with regard to topography and soil and we highlight those

130 units were there is often dissent between soil parent material as mapped by  
the soil surveyor and the geologic units mapped by geologists. The main aim  
of the random forest classification is not necessarily to improve the geological  
map with regard to its application as a parent material map, but foremost to  
135 identify the topographic characteristics of the parent material units in order  
to facilitate future, more detailed surveys.

## 2. Study area and data

### 2.1. General description

The study area includes the wide vale of Eppan-Kaltern, the Überetsch,  
located just south-west of Bozen in the Autonomous Province of Bolzano  
140 - South Tyrol, and extends in the north to the debris fan of Andrian in  
the Etsch Valley and the adjacent hill slope on the orographic right of the  
Etsch River. The western border of the study area is the steep slope of the  
Mendola-Roèn-Ridge, whereas the eastern border of the Überetsch as well as  
the study area is represented by the the Mitterberg, a ridge of Permian  
145 canites from which steep slopes descend to the Etsch Valley (approx. 200 m  
a.s.l.). The Kalterer Lake represents the southern limits of the investigated  
area. The land use of the paleovalley and its debris cones as well as the  
Etsch valley is dominantly apple orchards and vineyards, whereas the slope  
of the Mendola-Roèn-Ridge and the hilly outcrops of Vulcanites are covered  
150 by forests. Pastures are located mainly on till covering the flat areas of the  
Mendola-Roèn-Ridge.

### 2.2. Surficial Geology

A detailed description of the geologic situation can be found in the com-  
mentary to the new geologic map of Eppan (Avanzini et al., 2006). The  
155 paleovalley of Überetsch is described by Scholz et al. (2005) as a complex  
system of gravelly lateral moraines and large kame terraces, the result of the  
'Kaltern lobe', a Pleniglacial tongue of the Etsch valley glacier. Additionally,  
eroded remainders of debris flows that were deposited against the recessing  
glacier can be found along the slopes of the Mendola-Roèn-Ridge, as well as  
160 recent debris flow deposits, often composed of mainly limestone and dolomite  
fragment. The vale bottom itself is filled with Pleistocene sediments and con-  
tains a number of valleys carved into the gravels by fossil melt-water. At the  
eastern and western borders of the Pleistocene sediments, outcrops of Per-  
mian igneous Rhyolite and Lapilli-Tuff are responsible for a hilly relief, most

165 prominently at the eastern border of the study area where the Überetsch is  
separated from the Etsch valley by a steep slope down from the Mitterberg  
with an elevation difference of approximately 400 m. The steep slopes of the  
Mendola-Roèn-Ridge are dominated by various Dolomite units, with inter-  
mittent layers of sand and siltstones. Except for the very steep Dolomite  
170 walls of the ridge, the rarely occurring outcrops of these formations are sur-  
rounded and mostly covered by Pleistocene and Holocene slope debris, and  
in locally flatter areas by till.

The study area comprises two map sheets of the new geologic map of Italy,  
sheet Eppan, which covers the northern and major part of the area, as well as  
175 sheet Mezzo-Lombardo in the southern part. The sheets were published at a  
scale of 1 : 50,000 in 2007 and 2012, respectively. Mapping was performed at a  
scale of 1 : 10,000, this information was kindly provided by the Department of  
Geology and Building Material Tests of the Autonomous Province Bolzano,  
South Tyrol, in shapefile format and used for the analysis performed in this  
180 study.

As means for simplification of the analysis and data harmonisation, the  
geologic map units were generalised to the 16 SGUs described in Table 1, that  
allow for comparison with the parent material units described and identified  
by the soil surveyors in the field.

SGU	Abbrev.	short description	% area
alluvial deposits	AD	Holocene and Pleistocene deposits of silt, sand and gravels	14.9
coarse blocky debris	CBD	Holocene and Pleistocene blocky deposits of mass movements	1.8
colluvial deposits	CD	footslope deposits	2.4
calcareous sedimentary rock	CSR	limestones and dolomites	8.4
debris cones	DC	Holocene conic deposits from debris flows and torrents	12.7
glacio- and lacustrine deposits	GLD	(fine) sand deposits (with dropstones)	2.5
ice-marginal sediments	IMS	clast-supported gravels	0.2
intermediate sedimentary rock	ISR	silt- and sandstones	0.2
landslide deposits	LD	large landslide deposits	1.2
mire deposits	MrD	Holocene and Pleistocene silt and peat deposits	3.3
mixed deposits	MxD	Pleistocene deposits from debris flows, torrents and avalanches	2.1
siliceous bedrock	SB	rhyolite and rhyodazite tuffs and ignimbrites	13.0
slope debris	SD	Holocene and Pleistocene debris on slopes	10.3
siliceous sedimentary rock	SSR	sandstones and siltstones	1.1
till in general	TG	undifferentiated glacial sediment	25.9

Table 1: Table of the generalised parent material units with abbreviations and short description. Additionally, the proportion of the study area covered by each unit is given. Anthropogenic deposits and water bodies are not included in the analysis.

### 185 2.3. Soils

Overview of soils of the study area.

### 2.3.1. soil classification

The soil classification scheme applied in this study is the Austrian system (Nestroy et al., 2011), as most of the soil profile descriptions available for  
190 this study apply this system and it is generally recommendable to use local systems for large-scale mapping. Additionally, not all available soil profile data, especially those from points investigated only with augering, included sufficient information for deriving the reference soil group according to the World Reference Base for soil resources (IUSS Working Group WRB, 2015).  
195 Table 2 gives an overview of which reference soil groups are correlated to the relevant soil types in the Austrian classification. Classification of the soil profiles was performed at the subtype level, a basic overview of the Austrian soil classification system can be found in Baruck et al. (2016).

*data base 1: soil survey of agricultural areas in the Überetsch/Oltradige re-*  
200 *gion.* From 1993-1995 a soil survey of the farmlands in the region Überetsch was conducted (Thalheimer, 2006). Soil types were classified according to Soil Taxonomy, resulting in a soil map with 18 different soil series. 58 detailed soil pit descriptions were incorporated into the presented study, all located either in vineyards or apple orchards. Using the horizon descriptions, chem-  
205 ical properties as well as photographs of the pit face, theses soil profiles were reclassified applying the Austrian System.

*data base 2: soil survey 'ReBo - Terrain Classification of ALS Data to sup-*  
*port Digital Soil Mapping'.* During this project which was funded by the Autonomous Province Bolzano - South Tyrol and had the aim to investi-  
210 gate optimal cooperation between soil survey and terrain classification, 55 soil pit profiles were described in the presented studies area of interest. Soil classification was performed following Kilian (2015).

*data base 3: data base of the Forestry Service of the Autonomous Province Bolzano - South Tyrol.* 42 pit descriptions from the Forestry Service data set

215 *data base 4.* 227 auger observations (WLM)

### 2.4. Digital elevation data

## 3. Methods

### 3.1. General workflow

In a first step we analyse how well the geologic units of the high reso-  
220 lution geologic map correspond to the parent material identified by the soil

soil type	possible WRB group	short description	soil class
Grobmaterial-Rohboden	Leptosol, Regosol, Histosol	same as Feinmaterial-Rohboden but with more than 40 V.-% coarse fraction	Terrestrische Rohböden
Feinmaterial-Rohboden	Leptosol, Regosol, Histosol, Arenosol	only initial soil formation (Ai horizon) on parent material with less than 40 V.-% coarse fraction.	Terrestrische Rohböden
Rendzina	Leptosols, Histosols	with organic horizon on calcareous bedrock.	Terrestrische Humusböden
Kalklehm-Rendzina	Leptosol	soils with a loamy organic horizon on calcareous bedrock.	Terrestrische Humusböden
Pararendzina	Leptosol, Regosol, Umbrisol, Histosol	with organic horizon on carbonatic siliceous bedrock.	Terrestrische Humusböden
Ranker	Leptosol, Regosol, Umbrisol	with organic horizon on siliceous bedrock.	Terrestrische Humusböden
Braunerde	Cambisol, Fluvisol, Luvisol, Umbrisol, Regosol	with brown B-horizon owing to weathering and re-formation of clay minerals.	Braunerden
Parabraunerde	Luvisol, Albiluvisol, Cambisol	with eluvial horizon over clay-enriched B-horizon.	Braunerden
Semipodsol	Podzol, Regosol	characterized by moderate podzolization.	Podsole
Kalkbraunlehm	Cambisol, Luvisol	with a yellow- to redbrown cohesive B-horizon on calcareous bedrock, often fossil soils.	Kalklehme
Farb-Substratboden	Regosol, Alisol, Ferralsol, Luvisol, Nitisol, Arenosol	strong influence of color of parent material, overprinting horizon differentiation.	Substratböden
Textur-Substratboden	Regosol, Arenosol, Vertisol	strong influence of texture of parent material, overprinting horizon differentiation.	Substratböden
Kolluvisol	Anthrosol	developed from fine soil material relocated by (often human-induced) erosion.	Umgelagerte Böden
Rigolboden	Anthrosol	influenced by deep, homogenizing human cultivation.	Umgelagerte Böden
Haftnsse-Pseudogley	Stagnosol, Planosol	influenced by shallow, capillary stagnation phases.	Pseudogleye

Table 2: Table relating the Austrian soil types to WRB reference groups along with a simplified description, based on Kilian (2015).

surveyor, thus evaluating the performance and reliability of geologic maps to support soil survey in South Tyrol. This requires generalisation of the geologic units into surficial geology units (SGUs) that can be compared to the parent material units used in the soil (or forestry) surveys. The result is a confusion matrix that shows to which extent geologic units are in accordance with the parent material mapped by the surveyor. We highlight those units that are often confused or show overlap, and which should consequently be surveyed with greater detail and in consideration of relevant topographic information.

The next step is to perform a morphometric characterisation of the geological units. To better understand the topographic characteristics of a geologic unit, a data mining approach using random forest classification is performed. By applying a forward stepwise feature selection as well as the analysis of the parameter 'mean decrease accuracy', which quantifies the importance of a variable in the prediction procedure, we then identify which terrain parameters best separate parent materials units as well as geological map units and discuss how they can be related to and interpreted with regard to soil formation and the distribution of soil units. This data mining



procedure was applied to several groups of terrain parameters. One group  
240 included all computed terrain parameters, while other groups focus either on  
local or regional terrain parameters, or parameters related to surface rough-  
ness. An emphasis is set on the roughness measures, as they are deemed best  
suited to enhance the geologic map, as the geological units already implicitly  
contain information with regard to regional and local terrain parameters. In  
245 this study, this analysis is presented only for the area-wise most relevant ge-  
ologic units, and the focus is on separating those units which share common  
borders.

The connection between the two important soil forming factors, parent  
material and topography, on the one hand, and soil as the result of theses  
250 factors on the other, is then investigated by analysing the diversity and dis-  
tribution of soils for each geologic unit. This is performed from two points of  
view: the soil type distribution is done for profile sites per geologic unit, but  
also per parent material unit as attributed by the soil surveyor. This gives  
insight into how the surveyors' soil landscape model relates specific parent  
255 material units to specific soil types, especially when applying a morphologic-  
genetic classification such as the Austrian soil classification (Nestroy et al.,  
2011). The synthesis of this information then leads to a geologic-topographic  
characterisation (GTC) that describes each geologic unit.

### *3.2. Terrain parameters with emphasis on roughness measures*

#### *260 3.2.1. Local and regional terrain parameters*

#### *3.2.2. surface roughness-related terrain parameters*

ws=window size

*Topographic roughness index (TRI).* (Riley et al., 1999)

*Terrain texture.*

#### *265 Vector-based roughness measures (VRM and vectorruggedness).*

*Landform diversity.*

### *3.3. Random Forest classification*

Random Forest classifications was performed for several applications in  
the course of this study:

- 270 • To expand the geological map to a parent material map by model parent material based on the profile site descriptions, the topography of their location and the information provided by the new geologic map with its detailed information with regard to surficial geology
- 275 • To investigate which terrain parameters are best suited to discriminate for each SGU those points that were correctly classified to a parent material class by the geological map from those where this information differed to the parent material identified by the field surveyors.
- 280 • To analyse which terrain parameters best distinguish between adjacent soil parent material classes as identified by the soil surveyors during field survey
- To identify the terrain parameters that best separate and consequently characterise the SGUs as described by the geological map

285 Considering the confusion matrices and the accuracy measures calculated, this was performed as if the field observations described by the surveyors had been planned to validate the use of the geologic map as a map of soil parent material.

290 We can use the information on the confusion between certain classes implicitly contained in the sample point data, as well as the relative height information contained in the geologic map units. Consequently, roughness measure are sufficient to improve the parent material map.

## 4. Results and discussion

### 4.1. Comparison of soil parent material at soil profile sites with geologic map units

A comparison of the soil parent material class identified in the field survey and the SGU from the geological map for the same position shows that while these match for 186 of the profile sites, leading to an overall accuracy, or correct classification rate, of 49%, there is a high number of misclassifications. A first view at the confusion matrix (Table 3) indicates that a considerable amount of these misclassifications involve the classes till and slope deposits.

	AD	CBD	CD	CSR	DC	GLD	IMS	ISR	LD	MrD	MxD	SB	SD	SSR	TG
AD	7	0	0	0	0	1	0	0	0	0	0	0	1	0	3
CBD	0	4	0	0	0	0	0	0	0	0	0	0	8	0	2
CD	0	0	3	0	0	0	0	1	0	0	0	0	1	0	0
CSR	0	0	1	2	0	0	0	7	0	0	1	0	13	0	12
DC	4	0	3	0	5	0	0	2	0	0	0	0	20	0	2
GLD	0	0	0	0	0	5	0	0	0	0	0	0	0	0	2
IMS	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
ISR	0	0	0	0	0	0	0	0	0	0	0	0	3	0	0
LD	0	1	0	0	0	0	0	0	0	0	0	2	0	0	0
MrD	0	0	0	0	0	2	0	0	0	0	0	0	0	0	1
MxD	0	0	1	0	0	0	0	0	0	0	0	0	6	0	1
SB	0	1	1	0	0	2	0	1	0	0	2	14	3	1	24
SD	0	3	0	2	1	0	0	4	0	0	0	4	55	0	15
SSR	0	0	0	0	0	0	0	2	0	0	1	0	3	3	1
TG	1	0	0	1	0	1	0	3	0	1	5	0	9	0	88

Table 3: Tabular comparison of parent material geounits as observed by soil surveyor (columns) and in the geologic map (rows).

### 4.2. Geologic-topographic characterisation of the predominant SGUs

Through synthesis of results of the comparison of the soil point data and the geologic map, the random forest analysis of terrain parameters as well as the distribution of soil types in the study area, a geologic-topographic characterisation of the SGUs is performed. Only those units with substantial areal extent and sufficient soil profile points are described in detail.

#### 4.2.1. Alluvial deposits

The SGU alluvial deposits occupies 16.3 km, amounting to 14.9 % of the study area. It incorporates alluvial deposits in the paleovalley but also  
310 in the Etsch valley, with to soil profile points limited to the former. The alluvial deposits share very long borders with the SGUs till in general, debris cones, mire deposits and colluvial deposits, due to its bifurcated vertical transection of the study area. Long borders can therefore also be found with the units glaciolacustrine deposits, mixed deposits, slope debris and till in  
315 general. There is agreement between the geological map and the soil profile description in 7 (54%) of the 13 profiles for which the soil surveyors identified this SGU as the soil parent material. Some confusion with the SGUs debris cones and till in general can be observed, the former from the viewpoint of producer's reliability and the latter as an error of commission.

An investigation of the soil profile sites that were identified by surveyors as having alluvial deposits as parent material but are located on a different unit on the geological map, shows that these are characterized by consistently higher fischer k values, whereas slope shows less clear thresholds. Addressing the confusion of alluvial deposits with till, the random forest investigation  
320 shows that besides obvious channel-level related terrain parameters, which well separate the units, convexity contributes strongly to distinguishing between alluvial deposits and till on the geologic map, with the former characterised by lower values concentrating around 0.42. It must however be taken into account that the SGU in the study area also incorporates parts of the  
330 wide Etsch valley floor, whereas the soil profile points are mostly situated in the paleovalley of the Überetsch. The same investigation, but performed from the viewpoint of the surveyor (i.e. using the field data points and their parent material information), identifies a very local (small window size) version of the TRI as best suited to distinguish between the two parent material groups, with AD interestingly characterised by higher values indicating  
335 higher ruggedness. The separation of the profile sites of the parent material classes AD and debris cones, the second most common confusion for AD, is less clear than for AD and TG. High resolution roughness values are less important than in the case of till, with texture computed with 50 m grid cell  
340 size being the most decisive parameter. A comparison of the geological map units' topography also highlights the importance of this roughness parameter, which is surpassed only by the regional terrain parameter catchment slope, computed at the same low DTM resolution. The choice of these pa-

rameters seem closely linked to the transitional landscape position of debris  
345 cones, situated between the steep slopes of the Mendola-Roèn-Ridge and the  
flatter valley bottoms.

Of the 11 soil profile or auger points located on areas covered by alluvial  
deposits according to the reclassified geological map, the soil type Braunerde  
is predominant, with some occasional anthrosols. These brown soils are to be  
350 expected as the more or less pronounced stability of the flat alluvial deposits  
have allowed a certain degree of pedogenetic processes to occur. The Rigosols  
on the other hand are typical for the vineyards and orchards commonly found  
on alluvial deposits in the region, where landscape as well as soil have seen  
strong anthropogenic influence. The distribution of the soil types of the  
355 profile points identified to have alluvial deposits as parent material by the  
surveyors is comparable regarding the dominance of brown soils. Addition-  
ally, the soil types Kalklehm-Rendzina and Pararendzina were encountered,  
both characterised by organic horizons on more or less unweathered parent  
material, indicating that alluvial deposits were also identified at places lack-  
360 ing the stable conditions necessary for the development of a B-horizon as  
presented by the alluvial deposits unit of the geological map. So while the  
alluvial deposits unit is characterised on the geological map by the lowest  
mean slope aside from mire deposits, the soil data as well as the topographic  
analysis, indicate it is nevertheless necessary for future surveys to also inves-  
365 tigate the less typical, rougher and sloping areas at the border or in proximity  
of alluvial deposit units.

#### 4.2.2. *Siliceous bedrock*

The SGU siliceous bedrock in the study area is characterised by outcrops  
of rhyolitic ignimbrite in the Überetsch paleovalley, for instance forming the  
370 Mitterberg which separates the paleovalley from the current Etsch valley. It  
represents 13 % of the study area, and shares a long border with the SGU till  
in general, but also the units coarse blocky debris and debris cones. Shorter  
borders exist with almost all other SGU, especially mixed deposits and slope  
deposits. Regarding the comparison of the unit on the geological map with  
375 the parent material as reported by the surveyors, there is an interesting  
discrepancy between user's accuracy and producer's reliability when at-  
tempting the use of the geological map as a parent material map. While the  
parent material of only 14 of the 49 profile sites located on siliceous bedrock  
according to the geological map was identified as siliceous bedrock by the  
380 surveyors, 70 % of all soil pits with this parent material were actually on the

correct unit of the geological map. This means that when investigating soils on siliceous bedrock, the probability that they are encountered in the unit SB is higher than for other units, however it is similarly probable to encounter other parent materials in this unit. The greatest confusion occurred with the unit till in general, which was found to be the parent material of almost every second soil profile located on the SGU siliceous bedrock. Further parent materials identified on the SB unit include mixed and slope deposits, as well as glaciolacustrine deposits (Table 3). Confusion with slope debris or mixed deposits is understandable due to the often fuzzy transition from weathered bedrock to slope debris, or the fact that mixed deposits may very well contain almost only siliceous material from bedrock units at higher elevations in the catchment or underneath the debris cover layer. Additionally, the soils resulting from such parent material must not necessarily be very different to that on siliceous bedrock. Till, as a parent material, is a different case, as the material may be derived from catchments with a very different geology and the grain distribution is not comparable to that of slope debris or siliceous bedrock. When evaluating the misclassification of the parent materials till and siliceous bedrock, it is essential to consider different mapping procedures. Furthermore, the parent material layer must not necessarily be very thick. So, whereas the soil surveyor is particularly interested in the parent material, i.e. the material which through pedogenic processes and weathering slowly becomes the solum, no matter the thickness of this specific layer, the geologist's main concern is the underlying material, and may consider cover layers of till only once their thickness reaches a certain threshold, for instance 1 m as conveyed by the surveyors of the geologic survey of South Tyrol.

The random forest-based analysis of how to topographically separate the profile sites with siliceous bedrock as parent material that are situated on the SB unit of the geological map from those with the same parent material but on different SGUs showed that the latter group is situated in areas with lower convexity (computed with the 10 m DTM). Regarding roughness parameters, the vector ruggedness measure based on the 2.5 m grid and a large search window of approx. 130 m performed best. Soil profiles that evolved from siliceous bedrock and are also located on the SGU SB are characterised by higher vector ruggedness values. Given the strong confusion with the unit till in general, as well as the fact that these two units share a long border, it is of interest to examine the topographical differences between points that are attributed to one of these two parent material units. The TRI at a window size of approx. 100 m leads the terrain parameters with a 5-fold

cross-validated classification error of approx. 15 %. The minimum curvature  
420 based on the 10 m grid performs best from the group of local terrain parameters. In general, soil profile sites with SB as parent material are consequently characterised by rougher topography (increased TRI values) and slightly positive minimal curvature values when compared with data points where till was identified as parent material. This is well in line with increased distribution  
425 of this parent material on convex, ridge-like structures in the SGU siliceous bedrock. As Table 3 shows, confusion with slope deposits is present from both the user's and the producer's point of view. The random forest-based data mining shows that, regarding the parent material as indicated by the surveyors, the terrain parameters that are most characteristic of the difference between the two units are based on landform diversity. In the case of  
430 SB and SD, the patch density of the geomorphon-based landform map computed with a lookout distance of 7.5 m helped separate the two parent units due to the latter groups lower values, implying that areas of slope deposits have a slightly more homogeneous landform distribution, in the specific case  
435 large polygons of the landform slope, which seem to be a good indicator for slope deposits. Furthermore, the same TRI mentioned when separating SB and TG can also be a very useful addition to the parameter set when distinguishing SB from SD.

An investigation into the soil types classes of the soil profile sites located  
440 on the SGU siliceous bedrock shows that brown soils are the dominating soil type, represented by 37 of the 49 locations (75%). The class *Terrestrische Rohböden*, represented by the soil type Ranker, is second most common on this SGU, on which single examples of the soil types Semipodsol, Kalkbraunlehm, Farb-Substratböden and Kolluvisol were also performed. Taking into  
445 account only the soil profile sites for which the surveyors identified SB as parent material, whether or not actually located on this SGU, the soil type brown soils similarly dominates the soil type Ranker, with 13 and 6 profile sites, respectively. All of the other soil types found on the SGU on the geological map were not identified by the surveyors as having SB parent material,  
450 however one soil profile site with this parent material was attributed the soil type Grobmaterial-Rohboden. This soil type however does fit well for soils on siliceous bedrock outcrops, where only initial soil formation is possible, whereas the other soil types located on the SGU of the geological map are less plausible on SB, especially Kalkbraunlehm. In addition, the vast majority of the soil profile sites located on this SGU but identified to have a  
455 different parent material class (mostly till) by the surveyors, also feature the

soil type Braunerde, thereby decreasing the severity of misclassifying SB and TG, at least from a soil type perspective. From this perspective, the soil type Ranker is more characteristic of the SB unit, as it is much less common on  
460 TG with its characteristically more developed soils and the increased influence of calcareous materials. Furthermore, an examination of the points were parent material classes from observation and SGU map coincide, shows the dominance of brown soils compared to Rankers decrease, with proportions of 57 and 35 % for both soil types, respectively. In conclusion, the soil types  
465 Braunerde and Ranker are characteristic of this unit, with the former representing locations where weathering and clay-reformation has led to horizon differentiation not present in the less developed Ranker soils which are found on siliceous bedrock outcrops.

#### 4.2.3. *Till in general*

470 The unit till in general comprises lodgement and subglacial till, as well as other, undifferentiated till materials. It is found in the paleovalley as well as on flat terraces of the the Mendola-Roèn-Ridge. Covering more than 25 % of the study area, it is the most common SGU on the geological map, sharing borders with every single SGU, the longest, each with a length of at least  
475 30 km, being those with the SGUs alluvial deposits, siliceous bedrock and slope deposits. According to the geologic map, 109 of the soil data points are located on the SGU till in general, accounting for 29% of the points. In 88 of these locations, the surveyors agreed with regard to the parent material being till, leading to a user's accuracy of 80%. The majority of  
480 the other soil data points on this SGU were attributed the parent materials slope deposits or mixed deposits, with some intermediary sedimentary rocks also identified as parent material. While the user's accuracy for till is the best amongst the SGUs, the producer's reliability is not comparable, as the surveyors also identified till as the parent material at 63 further locations on  
485 different SGUs. The relative majority of these locations are found on the unit siliceous bedrock, already discussed above, but a large number of soil profiles with this parent material were also located on the units calcareous sedimentary rock and slope debris. Furthermore, till was identified as the parent material of soils on the units SSR, MxD, MrD, GLD, DC, CBD and  
490 AD. An important takeaway point from these results is that, at least in this study area, while the SGU unit till in general is a good indicator for where to reliably encounter soils evolved from till, it is of great importance to expect this parent material also on various other SGUs.



A topographic comparison of those points with till as parent material  
 located on the SGU till with points with the same parent material but on  
 495 different SGUs (Table x), shows that the misclassified, latter group is char-  
 acterized by a higher roughness (mean TRI of 25 compared to 13 for the  
 correctly classified group) and steeper slope (mean value of 21 compared to  
 12°). Regarding confusion with the soil parent material unit calcareous sedi-  
 500 mentary rock, data mining the terrain parameters of the relevant soil profile  
 sites again highlights the role of the parameter TRI, in this case dominantly  
 at a window size of 50 m, in topographically separating the unit till in general  
 from other parent material units. CSR profile sites are characterised by TRI  
 values in the upper quartile of the values characteristically displayed by TG  
 505 profile sites. Of the local terrain parameters, slope, together with minimal  
 curvature, performs well in separating the points of both groups, but not as  
 clearly as it separates both units on the geological map, as TG profile points  
 in the study area can be found at slope values higher than indicated in the  
 geological map. To characterise the difference of the TG sites with slope de-  
 510 posits, the TRI with a window size in the range of 100 m performs best of all  
 local, regional and roughness-related terrain parameters in the forward step-  
 wise feature selection. Similar to the situation with CSR, slope is the best  
 performing local terrain parameter, but as before, the separation is clearer  
 on the geological map, signifying that till can be found at steeper angles than  
 515 expected from the geological map. Till as the parent material of soil profiles  
 on the surficial geological map units CBD, DC and GLD is not as common  
 as for the above discussed units and additionally these units do not have  
 as many overall members, nevertheless some interesting observations can be  
 derived from the random forest-based investigation. For instance when com-  
 520 pared with till, debris cones, which have a considerably uniform topography  
 at the analysed grid cell size of 2.5 m, are characterised by lower local rough-  
 ness (vectorruggedness with  $ws = 50$  m or TRI with  $ws=7.5$  m) and a low  
 landform diversity as represented by a high mean patch size of landforms  
 calculated at a lookup distance of 7.5 m and a flatness threshold of 10°. On  
 525 the contrary, the profile site topography of the unit coarse blocky debris is  
 distinguishable from till by a higher TRI ( $ws=125m$ ) as well as steeper slope.

Other aspects that may account for the high number of till parent material  
 sites on units other than till in general have been discussed in the siliceous  
 bedrock section.

530 4.2.4. *Slope debris*

Slope debris, as a SGU of the geologic map, occupies 10% of the study area. Its by far longest border is shared with the unit siliceous bedrock, other important borders are with the units till in general, calcareous sedimentary rock and debris cones, the first three units greatly influencing the distribution of components in the slope debris units. When comparing the parent material of soil profile sites with the SGUs, the slope debris unit has slightly better user's accuracy than producer's reliability, as surveyors established slope debris as the parent material of 55 of the 84 profile points on this SGU, but also for 69 soil data points on other SGUs. Similar to the SGU siliceous bedrock, the most confusion regarding parent material on the SGU slope debris occurred with the unit till in general, which was identified at 15 soil profiles. As is the case with the SB unit, some of this confusion may be attributed to thin layers or punctual deposits of till. Considering slope debris, another important aspect is that the debris in question may very well be composed of till material that has been transported gravitationally. The same explanations may hold for other parent materials which were identified on the slope debris unit, especially for the bedrock units SB, ISR and CSR. While some isolated outcrops are possible, the most likely cause is that the constituents of the slope debris are so dominated by transported material of one of these bedrock classes, that the surveyors determine this unit as the parent material of the examined soil profile. On the other hand, misclassification between the units coarse blocky debris and slope deposits can be attributed to the fuzzy border between these units, ultimately linked to the grain size distribution, and the subjective interpretation thereof, especially during field survey. Contrary to most other SGUs (with the exception of CBD, LD and MxD), this unit itself does not provide information regarding the mineralogy of its component, which can only partially be derived through interpretation of its location in the catchment and the uphill geologic situation. Consequently, this unit is much better described by its topography than its material, as the latter may be highly diverse.

Those profile sites for which the geological map correctly proposes slope debris as the soil parent material differ from the slope deposit points on other SGUs mainly by higher slope heights and also higher slope angles, that is the geological map tends to under-represent slope debris situations in lower regions of the catchments and at less steep positions. 20 of the 122 soil profiles with slope debris as parent material are located on debris cones ac-

cording to the geological map, which are good examples of the mentioned situation. This is presumably closely related to the furrowed and disturbed character of debris cones in the southern half of the study area, where the  
570 situation may not present itself clearly during field survey. Additionally, the material composition as well as its origin is basically the same for slope debris and debris cones in this area. Topographically, roughness measures do well in separating the slope debris points from the debris cones points of the field survey data set. Slope debris is characterised by higher vector-  
575 ruggedness ( $ws=52.5$  m) values, which are very low for debris cones. Slope angles also differ considerably, implying that the surveyors tend to annotate slope debris if the topography does not show the low roughness and slope values characteristic of debris cones. This difference of roughness and slope with regard to the conceptual difference of slope debris and debris cones is  
580 also apparent when comparing the topography of both units as delineated by the geological map, however the difference is more distinct in this case. Similarly, calcareous sedimentary rock as parent material seems connected to roughness values slightly higher than those present for the average slope debris site, a relationship which is again more pronounced in the geological  
585 map. Consequently, it is in these transitional zones where the misclassification occurs. This topographical proximity of the units is of course reinforced by the fuzziness of distinguishing between the two parent material as such. As is the case with siliceous bedrock and slope debris, the border between the source material, in this case the calcareous bedrock, and its derivative, the  
590 slope debris, is often not clear when considering the parent material of a soil profile. The degree of weathering of the source material certainly adds to the confusion. The issues are comparable for the parent material units intermediate and siliceous sedimentary rock. While the gradual transition from source material, in these cases bedrock of differing mineralogical com-  
595 position, to slope debris poses various issues that cannot be explained only by topography, this is even more the case when considering the possibility of confusion between slope debris and coarse blocky debris, the distinction between which raises a number of questions with regard to the objectivity of estimating block or grain size of soil profile parent material. Misclassification  
600 between the units slope debris and mixed deposits are also predominantly a question of definition and the explanatory power of topography is very limited.

#### 4.2.5. *Calcareous sedimentary rock*

The calcareous sedimentary rocks in the study area are predominantly responsible for the steep walls of the upper elevations of the Mendola-Roèn-Ridge (Dolomite and limestone), but also represent some thin layers in lower parts of that slope, interchanging with intermediate and siliceous sedimentary rock layers. Despite covering only 8.4% of the area, it has long borders due to its layers that span from north to south of the study area. Slope debris units often occupy locations downslope of the CSR units, accounting for the long border length of 40 km with this unit. The fuzzy transition from weathered bedrock to slope debris is a major issue also for this SGU. The confusion matrix representing the comparison of the parent material as indicated by the geologic map and the parent material identified in field survey (Table 3) shows that, consequently, this is the unit responsible for the most misclassification, contributing to the very low user's accuracy of this SGU. Additionally, due to its intermittent layering with ISR, it is not surprising that this unit was found to be the parent material for 7 investigated soil profiles in the calcareous sedimentary rock unit. The producer's reliability is slightly better, however, all in all only five of the soil profiles were attributed the parent material unit calcareous sedimentary rock, of which two were in fact situated on the SGU slope debris. This again accentuates the problem with differentiating these two units when adjacent, as one is often the result of weathering, and gravitational transport, of the other unit. Similarly to the other bedrock units in the study area, till was reported also in this unit as the soil parent material at 12 sites, accounting for a third of the soil profile locations on this SGU and highlighting the necessity to expect thin layers of till in areas mapped as bedrock. Due to the small number of soil profile points actually identified as having CSR as parent material, the random forest classification approach of identifying important terrain parameter differences between the correctly classified points and those on other units is not very meaningful. Topographical separation from intermediate sedimentary rock, a major cause of misclassification, is also not very adequate, as the most important parameter in this case is slope height. This is to be expected, since the main difference is the vertical position in the overall geological structure of the study area, a difference which is however diminished by the above mentioned intermittent layering and gravitational movement down-slope, in addition to the above discussed problem regarding the definition of slope debris versus weathered bedrock.

640 4.2.6. *Debris cones*

The unit debris cones are located west of the center of the paleovalley, between the slope debris of the Mendola-Roèn-Ridge slope in the west, which are often the source area of these deposits, and the till deposited in the paleovalley in the east. Other units with which long borders exist are mixed, 645 colluvial, and alluvial deposits. The number of profile sites is comparably small with 36 profile sites, considering that the unit occupies almost 13% of the study area. A reason is that a large part of the debris cones are covered by settlements. Although the soil surveyors noted debris cones as parent material for only 5 soil profile sites, it must be considered that 20 of 650 the misclassified parent materials on the debris cones unit were identified as slope debris. Given the long mutual border and the fact that the source of the debris cones material is mainly the slope debris and the calcareous bedrock units which themselves are the origin of the slope debris, this misclassification may seem acceptable. In fact, it rather points out a difference in the point 655 of view of the soil and the geologic surveyors, where the first are interested in the material and the latter more in the landform presented by the debris cones unit. The remaining misclassifications are all with units that border this SGU, or, in the case of intermediate sedimentary rock, are located in the source region of the debris cones.

660 The topographic issues regarding the confusion with other parent materials such as AD, SD, TG have been discussed in the unit-specific subsections, highlighting this units landscape position as the main distinguishing characteristic.

4.2.7. *Intermediate sedimentary rock and siliceous sedimentary rock*

665 The units intermediate and siliceous sedimentary rock are situated in the slope of the Mendola-Roèn-Ridge and are characterised by more or less thin layer layers, often intermittent with layers of calcareous sedimentary rock, and only limited outcrops. Another common characteristic is that none of the soil profile points for which the surveyors identified one of these units 670 as the soil's parent material is actually situated on the geological map unit. The parent material of all three soil profile sites on ISR was classified as parent material, which is understandable considering the issues discussed in the previous sections regarding the slope debris and weathered bedrock. As further evidence of this issue, ISR was detected as the parent material of 4 675 soil profiles on the map unit slope debris. Additionally, intermediate sedimentary rock was identified as the parent material of 7 soils on the CSR map

unit and two on the SSR unit. There is of course no question that the layers are correctly located on the geological map with regard to the underlying geological structure of the study area, these soil profile sites rather highlight  
680 the possibility of finding any or all of the materials of these intermittent layers on these units as well as any potential down-slope debris units due to the prevailing geomorphodynamics in the study area. Gravitational transportation leads to these materials serving as soil parent material not where the layer is indicated on the geological map, but in fact down-slope of these  
685 layers. Additionally, these materials are seldom found as homogeneous units due to mixing and multilayering caused by gravitational transport. For these and other reasons discussed in the CSR section, topographical separation is limited for these units.

#### *4.2.8. Colluvial, mixed, and landslide deposits*

690 All three of these SGUs are characterised as material that has been transported down-slope by various processes. Regarding colluvial deposits, the surveyors indeed identified this parent material unit for three of the five profile sites on this map unit. The parent material of the remain two sites was mapped as ISR and SD. While the latter can be attributed to the similarity  
695 of the material and the uncertainty of the definition of the difference between slope deposits and colluvial deposits, the former can also indicate that the soil surveyor may value the information with regard to the mineralogy of greater importance to explaining a soil profile than the information regarding the morphodynamic history of the material. The definition of mixed  
700 deposits as deposits from debris flows, torrents and avalanches is similarly dominated by the means of transportation rather than the material components. Interestingly, none of the soil profiles located on this map unit were linked to this parent material by the surveyors, but 5 of the profiles on till were. With regards to soil parent material, an important issue is whether  
705 this material is in fact constituted of the same material as till and has been simply transported by one the mentioned processes. The parent material of the majority of the soil profiles on the MxD unit of the geological map were identified as slope deposits, highlighting the definitional proximity of these units, at least concerning the view of surveyors focussing on soil. Compared  
710 to these two units, the issue regarding landslide depositions is different one. This unit was never used by surveyors to describe the parent material of a soil profile site. Of the three data points on this map unit, the surveyors explicitly chose the parent material siliceous bedrock of two locations, which

is in fact the material of the landslide body in the study area. Coarse blocky  
715 debris was also used at one soil profile sites, which well describes the struc-  
ture of the material involved. This example provides insight into how the  
different emphasis and focus of soil and geologic surveyors influences their  
classification, each highlighting a different aspect of the same unit.

Topographically, these three units are not easy to differentiate, especially  
720 with regard to the unit slope debris, which is very close with regard to both  
definition and morphometry and discussed in the slope debris subsection.

#### 4.2.9. *Glacio-lacustrine deposits, ice-marginal sediments and mire deposits*

Even when taking into account that only 7 soil profile sites are located on  
this map unit, the geological map has relatively good user's accuracy for the  
725 soil parent material unit glacio-lacustrine deposits, comprising bother glacio-  
lacustrine and lacustrine deposits which are concentrated in the northern bor-  
der of the paleovalley. The producer's reliability is below 0.5, as surveyors  
indicated this parent material also for soil profile sites on the SGUs alluvial  
deposits, mire deposits, siliceous bedrock and till. Confusion with the first  
730 two SGUs seems acceptable due to the common fluviatile or hydromorphic  
history, although the other misclassifications do indicate the possibility of  
GLD as the parent material on these units. The topographical analysis of  
those points with GLD as parent material on other SGUS hints the these are  
located in regions where landform diversity is partly dominated by a single  
735 class at meso scale (ws=50 m), indicating a smoother surface than for the  
GLD map unit, as well as being characterised by a more pronounced negative  
minimal curvature (concave topography). The random forest based investi-  
gation into the topographical difference between soil profile points with SB  
and with GLD as parent material indicate that the latter class is character-  
740 ized by lower roughness values (TRI and vectorruggedness at window sizes  
between 100-130 m), indicating similar regions on SB as possible GLD areas.

Ice-marginal sediments are closely associated with till units and are mostly  
located at the border of these units, with whom they consequently share a  
long border. Given their often thin, elongated form and the fact that they  
745 cover only 0.2% of the study area, only one soil profile site is located on this  
SGU on the geologic map, of which the parent material however is classi-  
fied as alluvial deposits. This situation again highlights the problem fuzzy  
definitions that may overlap between closely related parent material units.  
Similarly, it may not be as important to the soil surveyors that the gravelly  
750 parent material is of ice-marginal origin, compared to alluvial, as it is to the

geologic surveyor specialized on the quarternary. Similarly, on the SGU mire deposits the surveyors identified the soil parent material glacio-lacustrine deposits twice and till once, with the only soil profile with mire deposits as parent material located on the SGU till. The influence of stagnant water on both the GLD and MrD can help understand the confusion between both units, and local mire deposits due to reduced permeability on clay-rich sub-glacial till is also rather likely. Given these local phenomena as well as the small number of sample profile sites, the topographical separation of the units IMS and MrD provides little relevant information.

#### 4.3. Random forest classification of parent material based on geologic map data and topography

A five-fold cross-validated forward step-wise feature selection was performed with random forest classification to produce a parent material map of the study area. Feature selection was performed to identify a small number of parameters capable of turning the geologic map into a parent material map. In contrast to using all possible terrain parameters, this procedure that produces only a limited number of explanatory variables, has the advantage of being easier to interpret, as the main aim is also to identify, highlight, and understand situations where the most confusion between parent material units as documented by surveyors and the geologic map occur. Additionally to a large number of terrain parameters, the geologic map, reclassified to the SGUs in Table 1, was added to the set of explanatory variables from which the random forest classifier could add features to the model. The whole feature selection procedure was also performed in 5-fold crossvalidation, and for 4 parameter sets: 1) all terrain parameters, 2) local terrain parameters, 3) regional terrain parameters, 4) surface roughness-related parameters. Without exception the SGU information was chosen first in every single feature selection run. Contrary to when investigating which terrain parameters best highlight the differences between two specific parent material units, local and regional terrain parameters, including relative elevations, were not identified as important predictor variables. Instead, the roughness-related measures dominated the selection process, even with the parameter set including all available terrain parameters, resulting in the choice of two of these measures and the SGU-information. The addition of further explanatory variables was not seen as an improvement with regard to both the model performance as well as its interpretability after evaluating some possible expanded models both with visual inspection of the results as well as the 1-standard error rule



(James et al., 2013). The reason for the decreased importance of regional, but also local, terrain parameters is that this information is already implicitly accounted for by the geologic map, as the various SGU classes are closely linked to certain relative elevations, especially the vertical distance to channel base level, due to overall geologic structure of the study area. Additionally, at least to a certain degree, some units like till, slope debris or alluvial deposits also provide implicit information regarding slope angles. Furthermore, a certain amount of slope information is also included in the TRI, which increases with slope as well with increasing roughness. The vector ruggedness measure well compliments the TRI and the SGU as it provides information on surface roughness independent of the slope gradient. Figure 1 shows the original SGU map as well as the parent material map based on SGUs, TRI and vector ruggedness.




Figure 1: Original reclassified SGUs (left) and random forest-based parent material map modified based on vector ruggedness and TRI (middle) and parent material map with only till modeled (right)

#### 4.3.1. *Discussion of accuracy*

As Olofsson et al. (2013) point out: uncertainty in remote sensing applications, but also in general, can be related to geolocation and interpreter variability. In this study...

- 805     -multiple interpreters, but not on same profiles, what about powell 2004 and their 5 interpretators
- Wickham 2013: constant communication in order to discuss and document difficult cases is important
- in future: response protocols!

#### 810   4.4. *Differences between geologic survey and parent material from profile site descriptions*

#### 4.5. *General difficulties*

- confusion at categorical transitions
- analyst biases (Foody 2010)
- 815   • Olofsson et al. (2013) state that with regard to reference labeling, in this case the parent material, difficult case need be noted for future reference and consensus development

#### 4.5.1. *Differences with regard to mapping purpose*

Between the two different frameworks of mapping, geology on the one  
820 hand and soil on the other, it is important to acknowledge the main focus of attention of each branch of research. There may exist a difference with regard to how pronounced a certain feature or characteristic must be in order to considered for mapping. Miller and Lee Burras (2015) note that the resulting maps of the two sciences try to communicate different aspects.  
825 While geology refers to geologic materials and general landform regions, soil science is concerned more with soil properties with regard to land use and management decisions.

A typical example is the unit landslide debris, which seems only of interest to the mappers with focus on geology, whereas the soil surveyor...

830   Regarding glacial deposits, the soil or forestry surveyors seldom differentiated between lodgement till and other glacial deposits, only when an decisively higher clay content was determined. On the other hand, the chemical properties, especially with regard to acidic properties of the tillic material are of much more interest to the soil surveyors who further differentiate till

835 with regard to this aspect. The geologic map on the other hand also contains information with regard to the age and the geologic system(synthem) to which certain moraines belong, whereas this is only of interest to the soil surveyor if it features additional information with regard to the mineralogic content of the different constituents of the moraine.

840 Geologist only map regolith cover layers thicker than approximately 1 m. What is more important for soil mapping, the transportation history of material or information regarding its composition?

#### *4.5.2. Nomenclatural differences and overlapping classes*

Congalton

845 *4.6. Pedologic interpretation of terrain parameters that best separate the geological units*

Do the points surveyed as moraine have different terrain parameters from those mapped as moraine on the geologic map.

*4.7. Distribution of soil types with regard to geologic unit as well as parent material unit*

850 *4.8. Influence of the Alpine environment on interpretability of geologic units as parent material units*

Heung et al. (2014) note that while traditional geologic maps focusing on bed rock are a valuable input for DSM when the residual materials form the soil parent material, but less so in areas distinguished by glaciation and high geomorphodynamics. Similarly, in their comparison of surficial geology maps derived from Soil Survey maps on the one hand and the Geologic Survey on the other, Miller and Lee Burras (2015) point out that the level of agreement was lower for areas with complicated geologic histories.

860 *4.8.1. High relief areas and multilayering*

Are there thresholds regarding terrain parameters?

*4.8.2. thin cover layers of till - an essential new parent material unit?*

Why is important to differentiate till and slope debris? While there may exist situations where the slope debris is composed of till material, the general difference is that soils from slope debris can be understood with knowledge of local geology and that they may not be so different from soils higher up in the catchment that evolved from bedrock, whereas till can consist of

chemically very different components. Till can be found at steeper slopes than expected, is this due to lateral consolidation of tillic material by glaciers in these U-shaped valleys?

*4.8.3. Is the morphodynamic background of deposits a necessary distinctive attribute from a pedological point of view?*

In the study area, mixed deposits from mass movement and torrents have the same components as till or hillside debris, which themselves are often the same...

*4.9. Terrain parameters - issues and results*

- the landform diversities are all heavily correlated and have the same meaning in the end.
- beware of the different class sizes when using random forest and box-plots to investigate differences
- In general local terrain parameters and regional terrain parameters are only important if the parent material map does not apply the information already implicitly provided in the new detailed geological map.

## 5. Conclusion

We propose that future surveys focus increasingly on these units with greater uncertainty with regard to soil parent material to strengthen understanding of the pedologic relevance of these units. By performing a GTC prior to future detailed field soil surveys, the surveyor can make best use of available information and concentrate the time and money consuming task of field work, involving soil pits and auguring, on units identified as highly variable and uncertain regarding soils. This information can be additionally helpful for devising future sampling procedures and also for consideration when attempting to regionalise point information.

## Acknowledgements

This research was performed within the project 'Terrain Classification of ALS Data to support Digital Soil Mapping', funded by the Autonomous Province Bolzano – South Tyrol (15/40.3).

	CBD	CD	CSR	DC	GLD	IMS	ISR	LD	MrD	MxD	SB	SD	SSR	TG
AD	0.2	21.1	0.0	23.6	14.8	2.5	0.0	1.2	22.2	10.9	5.7	12.7	0.2	45.2
CBD		0.5	6.1	8.5	0.0	0.0	1.1	0.9	0.5	1.5	10.1	9.7	0.9	9.1
CD			1.3	17.6	10.7	2.8	0.0	0.6	0.7	10.0	5.8	4.0	0.0	31.1
CSR				11.8	0.0	0.0	5.5	0.8	0.0	1.3	0.6	40.7	5.0	23.1
DC					2.5	0.6	0.7	2.4	5.9	17.4	9.3	25.6	1.6	40.6
GLD						2.2	0.0	0.2	0.8	3.4	0.4	0.5	0.0	7
IMS							0.0	0.0	0.0	0.6	0.7	0.4	0.0	1.1
ISR								0.2	0.0	0.0	0.0	4.3	0.0	0.4
LD									0.0	0.1	1.4	3.7	0.5	4.2
MrD										0.2	1.2	0.6	0.0	1.9
MxD											2.7	2.7	0.1	19.5
SB												109.3	4.4	67.4
SD													8.3	50.4
SSR														5.5

Table 4: Length in kilometers of the borders of adjacent SGUs

## additional tables

## References

- 900 Avanzini, M., Bargossi, G., Borsato, A., Castiglioni, G., Cucato, M., Morelli, C., Prosser, G., Sapelza, A., 2006. Erläuterungen zur geologischen Karten von Italien im Maßstab 1:50000 Blatt 026 Eppan. Technical Report. Agenzia per la protezione dell’ambiente e per i servizi tecnici.
- Baruck, J., Nestroy, O., Sartori, G., Baize, D., Traidl, R., Vraj, B., Brm, E., Gruber, F.E., Heinrich, K., Geitner, C., 2016. Soil classification and mapping in the alps: The current state and future challenges. *Geoderma* 264, 312 – 331. URL: <http://www.sciencedirect.com/science/article/pii/S0016706115300343>, doi:<http://dx.doi.org/10.1016/j.geoderma.2015.08.005>. soil mapping, classification, and modelling: history and future directions.
- 910 Beasom, S.L., Wiggers, E.P., Giardino, J.R., 1983. A technique for assessing land surface ruggedness. *The Journal of Wildlife Management* 47, 1163–1166. doi:10.2307/3808184.
- Böhner, J., AntoniĆ, O., 2009. Chapter 8 land-surface parameters specific to topo-climatology, in: Hengl, T., Reuter, H.I.
- 915

- (Eds.), Geomorphometry Concepts, Software, Applications. Elsevier. volume 33 of *Developments in Soil Science*, pp. 195 – 226. URL: <http://www.sciencedirect.com/science/article/pii/S0166248108000081>, doi:[http://dx.doi.org/10.1016/S0166-2481\(08\)00008-1](http://dx.doi.org/10.1016/S0166-2481(08)00008-1).  
920
- Brevik, E.C., Miller, B.A., 2015. The use of soil surveys to aid in geologic mapping with an emphasis on the eastern and midwestern united states. *Soil Horizons* 56. URL: <http://dx.doi.org/10.2136/sh15-01-0001>, doi:10.2136/sh15-01-0001.
- 925 Coblentz, D., Pabian, F., Prasad, L., 2014. Quantitative geomorphometrics for terrain characterization. *International Journal of Geosciences* 5, 247–266. URL: <http://dx.doi.org/10.4236/ijg.2014.53026>.
- FAO, 2006. Guidelines for soil description. Food and Agricultural Organisation of the United Nations. Rome.
- 930 Gallant, J.C., Wilson, J.P., 2000. Terrain Analysis - Principles and Applications. John Wiley & Sons, Inc.. chapter Primary Topographic Attributes. pp. 51–85.
- Geitner, C., Baruck, J., Freppaz, M., Godone, D., Grashey-Jansen, S., Gruber, F.E., Heinrich, K., Papritz, A., Simon, A., Stanchi, S., Traidl, R., von Albertini, N., Vraj, B., 2017. Chapter 8 - soil and land use in the alps challenges and examples of soil-survey and soil-data use to support sustainable development, in: Pereira, P., Brevik, E.C., Muoz-Rojas, M., Miller, B.A. (Eds.), *Soil Mapping and Process Modeling for Sustainable Land Use Management*. Elsevier, pp. 221 – 292. URL: <http://www.sciencedirect.com/science/article/pii/B9780128052006000086>,  
940 doi:<http://doi.org/10.1016/B978-0-12-805200-6.00008-6>.
- Grohmann, C.H., Smith, M.J., Riccomini, C., 2010. Multiscale analysis of topographic surface roughness in the midland valley, scotland. *IEEE Transactions on Geoscience and Remote Sensing* PP, 1 – 14. doi:10.1109/TGRS.2010.2053546.  
945
- Heung, B., Bulmer, C.E., Schmidt, M.G., 2014. Predictive soil parent material mapping at a regional-scale: A random forest approach. *Geoderma* 214215, 141 – 154. URL: <http://www.sciencedirect.com/science/>

- 950 article/pii/S0016706113003443, doi:<http://dx.doi.org/10.1016/j.geoderma.2013.09.016>.
- Hobson, R.D., 1972. Spatial analysis in geomorphology. Harper and Row, New York. chapter Surface roughness in topography: quantitative approach. pp. 221–245.
- IUSS Working Group WRB, 2015. World Reference Base for Soil Resources 955 2014, update 2015 International soil classification system for naming soils and creating legends for soil maps. World Soil Resources Reports 106. FAO, Rome. Rome.
- James, G., Witten, D., Hastie, T., Tibshirani, R., 2013. An Introduction to Statistical Learning. Springer-Verlag New York. URL: <http://www-bcf.usc.edu/~garth/ISL/index.html>, doi:10.1007/978-1-4614-7138-7. 960
- Jenny, H., 1941. Factors of Soil Formation - A System of Quantitative Pedology. Dover Publications, Inc. New York.
- Juilleret, J., Dondeyne, S., Vancampenhout, K., Deckers, J., Hissler, C., 2016. Mind the gap: A classification system for integrating the subsolum into soil surveys. Geoderma 264, Part B, 332 – 339. URL: <http://www.sciencedirect.com/science/article/pii/S0016706115300604>, doi:<http://dx.doi.org/10.1016/j.geoderma.2015.08.031>. soil mapping, classification, and modelling: history and future directions. 965
- 970 Kilian, W., 2015. Schlüssel zur Bestimmung der Böden Österreichs. volume 81. 2 ed., Österr. Bodenkundl. Ges.
- Marchi, L., Dalla Fontana, G., 2005. Gis morphometric indicators for the analysis of sediment dynamics in mountain basins. Environmental Geology 48, 218–228. URL: <http://dx.doi.org/10.1007/s00254-005-1292-4>, doi:10.1007/s00254-005-1292-4. 975
- McBratney, A.B., Mendonca Santos, M., Minasny, B., 2003. On digital soil mapping. Geoderma 117, 3 – 52.
- Miller, B.A., Lee Burras, C., 2015. Comparison of surficial geology maps based on soil survey and in depth geological survey. Soil Horizons

- 980 56. URL: <http://dx.doi.org/10.2136/sh14-05-0005>, doi:10.2136/sh14-05-0005.
- Nellemann, C., Thomsen, M.G., 1994. Terrain ruggedness and caribou forage availability during snowmelt on the arctic coastal plain, alaska. *Arctic* 47, 361–367. URL: <http://www.jstor.org/stable/40511597>.
- 985 Nestroy, O., Aust, G., Blum, W., Englisch, M., Hager, H., Herzberger, E., Kilian, W., Nelhiebel, P., G. Ortner and, E.P., und J. Wagner, A.P.W.S., 2011. Systematische gliederung der bden sterreichs. sterreichische boden-systematik 2000 in der revidierten fassung von 2011. *Mitt. sterr. Bodenkdl. Ges.* 79.
- 990 Olaya, V., 2009. Chapter 6 basic land-surface parameters, in: Hengl, T., Reuter, H.I. (Eds.), *Geomorphometry Concepts, Software, Applications*. Elsevier. volume 33 of *Developments in Soil Science*, pp. 141 – 169. URL: <http://www.sciencedirect.com/science/article/pii/S0166248108000068>, doi:[http://dx.doi.org/10.1016/S0166-2481\(08\)](http://dx.doi.org/10.1016/S0166-2481(08)00006-8)  
995 00006-8.
- Olofsson, P., Foody, G.M., Stehman, S.V., Woodcock, C.E., 2013. Making better use of accuracy data in land change studies: Estimating accuracy and area and quantifying uncertainty using stratified estimation. *Remote Sensing of Environment* 129, 122 – 131. URL: <http://www.sciencedirect.com/science/article/pii/S0034425712004191>, doi:<https://doi.org/10.1016/j.rse.2012.10.031>.  
1000
- Phillips, J.D., Lorz, C., 2008. Origins and implications of soil layering. *Earth-Science Reviews* 89, 144 – 155. URL: <http://www.sciencedirect.com/science/article/pii/S001282520800055X>, doi:<https://doi.org/10.1016/j.earscirev.2008.04.003>.  
1005
- Riley, S.J., DeGloria, S.D., Elliot, R., 1999. A terrain ruggedness index that quantifies topographic heterogeneity. *Intermountain Journal of Sciences* 5, 23 – 27.
- 1010 Sappington, J.M., Longshore, K.M., Thompson, D.B., 2007. Quantifying landscape ruggedness for animal habitat analysis: A case study using



bighorn sheep in the mojave desert. *Journal of Wildlife Management* 75.  
URL: <http://dx.doi.org/10.2193/2005-723>, doi:10.2193/2005-723.

1015 Schaetzl, R.J., Krist Jr., F.J., Rindfleisch, P.R., Liebens, J., Williams,  
T.E., 2000. Postglacial landscape evolution of northeastern lower michi-  
gan, interpreted from soils and sediments. *Annals of the Association of  
American Geographers* 90, 443–466. URL: <http://dx.doi.org/10.1111/0004-5608.00204>, doi:10.1111/0004-5608.00204.

1020 Scholz, H., Bestle, K.H., Willerich, S., 2005. Quartärgeologische untersuchun-  
gen im überetsch. *Geo.Alp* 2, 1–23.

Thalheimer, M., 2006. Kartierung der landwirtschaftlich genutzten  
bden des Überetsch in südtirol (italien). *Laimburg Journal* 3, 135–  
177. URL: [http://www.laimburg.it/de/projekte-publikationen/  
blickpunkte.asp?news\\_action=300&news\\_image\\_id=828290](http://www.laimburg.it/de/projekte-publikationen/blickpunkte.asp?news_action=300&news_image_id=828290).