

# Joint analysis of geological map units and topography to support soil survey - a case study from South Tyrol

Fabian E. Gruber<sup>a,\*</sup>, Jasmin Baruck<sup>a</sup>, Clemens Geitner<sup>a</sup>

<sup>a</sup>*Institute of Geography, University of Innsbruck, Innrain 52f, 6020 Innsbruck, Austria*

---

## Abstract

*Keywords:*

---

## 1. Introduction

*general introduction.* Geologic maps have always been an important aid in soil survey as parent material is an important factor in soil formation (Jenny, 1941). The importance of this relationship is highlighted by the fact that, 5 vice versa, soil maps have themselves been applied to support and improve geologic mapping (Brevik and Miller, 2015).

Hier vielleicht Juilleret und so für die Wichtigkeit des Ausgangsmaterials.  
Situation in the Alps

*overview of literature with regard to the use of geologic maps in classic field 10 soil survey and digital soil mapping in literature- what parent material classes are there in the classifications.*

*literature overview with regard to terrain parameters for characterisation and soil survey, allgemeine geomorphometry kurz zitieren.* Researchers have long investigated ways to quantify the roughness or ruggedness of terrain, from 15 the analysis of field data and topographic maps to computing roughness indices on raster grids. Geology, geomorphology as well as habitat modelling and wildlife management have been the main research areas in which such investigations were performed on land surfaces. Hobson (1972) presented three

---

\*Corresponding author

Email address: `Fabian.Grubert@uibk.ac.at` (Fabian E. Gruber)

different roughness values and applied them to field measurements, correlat-  
 20 ing them to rock type. In another early study aimed at quantifying roughness,  
 Beasom et al. (1983) presented the land surface ruggedness index, which is  
 based on the total length of contour lines per area. Similarly analysing topo-  
 graphic maps, Nellemann and Thomsen (1994) describe the calculation of a  
 terrain ruggedness index based on the variability of contour lines along tran-  
 25 sects, which they correlate with caribou forage availability. Regarding field  
 methods, they calculate microtopographic diversity by analysing the hori-  
 zontal distance of a chain laid on the ground in their study plots. Riley et al.  
 (1999) proposed the topographic ruggedness index (TRI), which compares  
 the elevation of a central pixel to the elevations of cells within a given search  
 30 window. In an attempt to decorrelate roughness from slope, Sappington et al.  
 (2007) expanded on the work of Hobson (1972) to introduce vector rugged-  
 ness measure (VRM), which is calculated based on the orientation of vectors  
 normal to the surface in a given area. (Grohmann et al., 2010) analysed  
 several roughness measures at different resolutions and window sizes with  
 35 regard to their ability to depict terrain features. They highlight the ability  
 of VRM to detect fine-scale roughness features and attribute low roughness  
 values to steep but smooth slopes, but also acknowledge its inability to de-  
 limit slope breaks and identify regional relief. The Melton ruggedness index,  
 which relates the elevation difference of a basin to the drainage area, was  
 40 applied by Marchi and Dalla Fontana (2005) to investigate sediment trans-  
 port, however compared to VRM it is more of a measure of general relief  
 than roughness. Similar to VRM, roughness measures based on eigenvalue  
 ratios of an orientation matrix have long been used in geology to describe  
 land surfaces, especially of bedrock. Coblenz et al. (2014) investigated the  
 45 use of such measures ...

*overview of intention and aims.* In a first step we analyse how well the ge-  
 ologic units of the high resolution geologic map correspond to the parent  
 material identified by the soil surveyor, thus evaluating the performance and  
 reliability of geologic maps to support soil survey in South Tyrol. This re-  
 50 quires generalisation of the geologic units into surficial geology units (SGUs)  
 that can be compared to the parent material units used in the soil (or forestry)  
 surveys, called The result is a confusion matrix that shows to which extent  
 geologic units are in accordance with the parent material mapped by the  
 surveyor. We highlight those units that are often confused or show over-  
 55 lap, and which should consequently be surveyed with greater detail and in

consideration of relevant topographic information.

The next step is to perform a morphometric characterisation of the geological units. Applying a data mining approach based on a forward stepwise feature selection with a SVM classifier, we then identify which terrain parameters best separate geologic units and discuss how they can be related to and interpreted with regard to soil formation and the distribution of soil units.

The connection between the two important soil forming factors, parent material and topography, on the one hand, and soil as the result of these factors on the other, is then investigated by analysing the diversity and distribution of soils for each geologic unit. This is performed from two points of view: the soil type distribution is done for profile sites per geologic unit, but also per parent material unit as attributed by the soil surveyor. This gives insight into how the surveyors' soil landscape model relates specific parent material units to specific soil types, especially when applying a morphologic-genetic classification such as the Austrian soil classification (Nestroy et al., 2011). The synthesis of this information then leads to a geologic-topographic characterisation (GTC) that describes each geologic unit.

The aim of this study is to evaluate how to make best use of available geologic and topographic information for soils survey. Hence each geologic unit is characterised with regard to topography and soil and we highlight those units where there is often dissent between soil parent material as mapped by the soil surveyor and the geologic units mapped by geologists.

## 2. Material and Methods

### 2.1. Study area and data

*General description.* The study area includes the wide vale of Eppan-Kaltein, the Überetsch, located just south-west of Bozen in the Autonomous Province of Bolzano - South Tyrol, and extends in the north to the debris fan of Andrian in the Etsch Valley and the adjacent hillslope on the orographic right of the Etsch River. The western border of the study area is the steep slope of the Mendola-Roè-Ridge, whereas the eastern border of the Überetsch as well as the study area is represented by the Mitterberg, a ridge of Permian Vulkanites from which steep slopes descend to the Etsch Valley (approx. 200 m a.s.l.). The Kalterer Lake represents the southern limits of the investigated area. The land use of the paleovalley and its debris cones as well as the Etsch valley is dominantly apple orchards and vineyards, whereas the slope

of the Mendola-Roèn-Ridge and the hilly outcrops of Vulcanites are covered by forests. Pastures are located mainly on till covering the flat areas of the Mendola-Roèn-Ridge.

95 *2.1.1. Surficial Geology*

A detailed description of the geologic situation can be found in the commentary to the new geologic map of Eppan (Avanzini et al., 2006). The paleovalley of Überetsch is described by Scholz et al. (2005) as a complex system of gravelly lateral moraines and large kame terraces, the result if the  
100 'Kaltern lobe', a Pleniglacial tongue of the Etsch valley glacier. Additionally, eroded remainders of debris flows that were deposited against the recessing glacier can be found along the slopes of the Mendola-Roèn-Ridge, as well as recent debris flow deposits, often composed of mainly limestone and dolomite fragment. The vale bottom itself is filled with Pleistocene sediments and contains a number of valleys carved into the gravels by fossil meltwater. At the  
105 eastern and western borders of the Pleistocene sediments, outcrops of Permian igneous Rhyolite and Lapilli-Tuff are responsible for a hilly relief, most prominently at the eastern border of the study area where the Überetsch is separated from the Etsch valley by a steep slope down from the Mitterberg with an elevation difference of approximately 400 m. The steep slopes of the  
110 Mendola-Roèn-Ridge are dominated by various Dolomite units, with intermittent layers of sand and siltstones. Except for the very steep Dolomite walls of the ridge, the rarely occurring outcrops of these formations are surrounded and mostly covered by Pleistocene and Holocene slope debris, and  
115 in locally flatter areas by till.

The study area comprises two map sheets of the new geologic map of Italy, sheet Eppan, which covers the northern and major part of the area, as well as sheet Mezzo-Lombardo in the southern part. The sheets were published at a scale of 1 : 50,000 in 2007 and 2012, respectively. Mapping was performed at  
120 a scale of 1 : 10,000, this information was kindly provided by the Department of Geology and Building material tests of the Autonomous Province Bolzano, South Tyrol, in shapefile format and used for the analysis performed in this study.

As means for simplification of the analysis and data harmonisation, the  
125 geologic map units were generalised to the 15 SGUs described in Table 1, that allow for comparison with the parent material units described and identified by the soil surveyors in the field.

### *2.1.2. Soils*

*Overview of soils in study area.*

130 *Soil profile data.*

.

### *2.1.3. Digital elevation data*

## *2.2. Methods*

### *2.2.1. Terrain parameters with emphasis on roughness measures*

135 (Riley et al., 1999)

### *2.2.2. general methodology*

SGU	Abbrev.	short description	% area
alluvial deposits	AD	Holocene and Pleistocene deposits of silt, sand and gravels	10.6
coarse blocky debris	CBD	Holocene and Pleistocene blocky deposits of mass movements	2.1
colluvial deposits	CD	footslope deposits	2.8
calcareous sedimentary rock	CSR	limestones and dolomites	7.4
debris cones	DC	Holocene conic deposits from debris flows and torrents	14.2
glaciolacustrine deposits	GD	(fine) sand deposits with dropstones	2.8
ice-marginal sediments	IMS	clast-supported gravels	0.3
intermediate sedimentary rock	ISR	silt- and sandstones	2.8
mire deposits	MrD	Holocene and Pleistocene silt and peat deposits	3.9
mixed deposits	MxD	Pleistocene deposits from debris flows, torrents and avalanches	3.2
siliceous bedrock	SB	rhyolite and rhyodazite tuffs and ignimbrites	7.4
slope debris	SD	Holocene and Pleistocene debris on slopes	11.4
sub-glacial till	SGT	compacted sub-glacial sediment	18.7
siliceous sedimentary rock	SSR	sandstones and siltstones	1.1
till in general	TG	undifferentiated glacial sediment	11.3

Table 1: Table of the generalised parent material geounits with abbreviations and short description. Additionally, the proportion of the study area covered by each geounit is given.

soil type	possible WRB group	short description
Braunerde	Cambisol, Fluvisol, Luvisol, Umbrisol, Regosol	with brown B-horizon owing to weathering and re-formation of clay minerals.
Farb-Substratboden	Regosol, Alisol, Ferralsol, Luvisol, Nitisol, Arenosol	strong influence of color of parent material, overprinting horizon differentiation.
Feinmaterial-Rohboden	Leptosol, Regosol, Histosol, Arenosol	only initial soil formation (Ai horizon) on parent material with less than 40 V.-% coarse fraction.
Grobmaterial-Rohboden	Leptosol, Regosol, Histosol	same as Feinmaterial-Rohboden but with more than 40 V.-% coarse fraction
Haftnsse-Pseudogley	Stagnosol, Planosol	influenced by shallow, capillary stagnation phases.
Kalkbraunlehm	Cambisol, Luvisol	with a yellow- to redbrown cohesive B-horizon on calcareous bedrock, often fossil soils.
Kalklehm-Rendzina	Leptosol	soils with a loamy organic horizon on calcareous bedrock.
Kolluvisol	Anthrosol	developed from fine soil material relocated by (often human-induced) erosion.
Parabraunerde	Luvisol, Albiluvisol, Cambisol	with eluvial horizon over clay-enriched B-horizon.
Pararendzina	Leptosol, Regosol, Umbrisol, Histosol	with organic horizon on carbonatic siliceous bedrock.
Ranker	Leptosol, Umbrisol, Regosol	with organic horizon on siliceous bedrock.
Rendzina	Leptosols, Histosols	with organic horizon on calcareous bedrock.
Rigolboden	Anthrosol	influenced by deep, homogenizing human cultivation.
Semipodsol	Podzol, Regosol	characterized by moderate podzolization.
Textur-Substratboden	Regosol, Arenosol, Vertisol	strong influence of texture of parent material, overprinting horizon differentiation.

Table 2: Table relating the Austrian soil types to WRB reference groups along with a simplified description, based on Kilian (2015).

### 3. Results

	AD	CD	CSR	DC	GD	IMS	ISR	MrD	MxD	SB	SC	SD	SGT	SSR	TG
AD		1.96	0.00	2.19	1.37	0.23	0.00	2.29	1.01	0.53	0.01	1.18	31.49	0.02	1.05
CD			0.12	1.63	0.89	0.26	0.00	0.06	0.93	0.54	0.05	0.37	0.59	0.00	1.18
CSR				1.10	0.00	0.00	0.51	0.00	0.12	0.06	0.57	3.89	0.07	0.47	2.06
DC					0.23	0.05	0.07	0.58	1.62	0.88	0.79	2.41	0.72	0.15	3.05
GD						0.20	0.00	0.00	0.31	0.03	0.00	0.03	0.27	0.00	0.24
IMS							0.00	0.00	0.05	0.06	0.00	0.03	0.09	0.00	0.09
ISR								0.00	0.00	0.00	0.10	0.40	0.00	0.00	0.04
MrD									0.02	0.11	0.05	0.05	0.17	0.00	0.01
MxD										0.25	0.14	0.25	0.93	0.01	0.82
SB											0.94	10.18	2.03	0.41	4.14
SC												0.90	0.12	0.09	0.73
SD													0.79	0.77	3.80
SGT														0.12	0.70
SSR															0.20
TG															

Table 3: Length in kilometers of the borders of adjacent SGUs

#### 3.1. Geomorphometric analysis of geologic map units

#### 3.2. Distribution of soils with regard to geologic units

### 4. Discussion

#### 4.1. Differences between geologic survey and parent material from profile site descriptions

##### 4.1.1. Differences with regard to mapping purpose

Between the two different frameworks of mapping, geology on the one hand and soil on the other, it is important to acknowledge the main focus of attention of each branch of research. There may exist a difference with regard to how pronounced a certain feature or characteristic must be in order to considered for mapping.

A typical example is...



- 150 4.1.2. *Nomenclatural differences and overlapping classes*
- 4.2. *Pedologic interpretation of terrain parameters that best separate the geological units*
- 4.3. *Distribution of soil types with regard to geologic unit as well as parent material unit*
- 155 4.4. *Influence of the Alpine environment on interpretability of geologic units as parent material units*
- 4.4.1. *High relief areas and multilayering*
- Are there thresholds regarding terrain parameters?
- 4.4.2. *thin cover layers of till - an essential new parentmaterial unit?*
- 160 4.4.3. *Is the morphodynamic background of deposits a necessary distinctive attribute from a pedological point of view?*

In the study are, mixed deposits from mass movement and torrents have the same components as till or hillside debris, which themselves are often the same...

## 165 5. Conclusion

We propose that future surveys focus increasingly on these units with greater uncertainty with regard to soil parent material to strengthen understanding of the pedologic relevance of these units. By performing a GTC prior to future detailed field soil surveys, the surveyor can make best use of

170 available information and concentrate the time and money consuming task of field work, involving soil pits and auguring, on units identified as highly variable and uncertain regarding soils. This information can be additionally helpful for devising future sampling procedures and also for consideration when attempting to regionalise point information

## 175 References

- Avanzini, M., Bargossi, G., Borsato, A., Castiglioni, G., Cucato, M., Morelli, C., Prosser, G., Sapelza, A., 2006. Erläuterungen zur geologischen Karten von Italien im Maßstab 1:50000 Blatt 026 Eppan. Technical Report. Agenzia per la protezione dell'ambiente e per i servizi tecnici.
- 180 Beasom, S.L., Wiggers, E.P., Giardino, J.R., 1983. A technique for assessing land surface ruggedness. The Journal of Wildlife Management 47, 1163–1166. doi:10.2307/3808184.

- Brevik, E.C., Miller, B.A., 2015. The use of soil surveys to aid in geologic mapping with an emphasis on the eastern and midwestern united states. Soil Horizons 56. URL: <http://dx.doi.org/10.2136/sh15-01-0001>, doi:10.2136/sh15-01-0001.
- Coblentz, D., Pabian, F., Prasad, L., 2014. Quantitative geomorphometrics for terrain characterization. International Journal of Geosciences 5, 247–266. URL: <http://dx.doi.org/10.4236/ijg.2014.53026>.
- Grohmann, C.H., Smith, M.J., Riccomini, C., 2010. Multiscale analysis of topographic surface roughness in the midland valley, scotland. IEEE Transactions on Geoscience and Remote Sensing PP, 1 – 14. doi:10.1109/TGRS.2010.2053546.
- Hobson, R.D., 1972. Spatial analysis in geomorphology. Harper and Row, New York. chapter Surface roughness in topography: quantitative approach. pp. 221–245.
- Jenny, H., 1941. Factors of Soil Formation - A System of Quantitative Pedology. Dover Publications, Inc. New York.
- Kilian, W., 2015. schlüssel zur Bestimmung der Böden Österreichs. volume 81. 2 ed., Österr. Bodenkundl. Ges.
- Marchi, L., Dalla Fontana, G., 2005. Gis morphometric indicators for the analysis of sediment dynamics in mountain basins. Environmental Geology 48, 218–228. URL: <http://dx.doi.org/10.1007/s00254-005-1292-4>, doi:10.1007/s00254-005-1292-4.
- Nellemann, C., Thomsen, M.G., 1994. Terrain ruggedness and caribou forage availability during snowmelt on the arctic coastal plain, alaska. Arctic 47, 361–367. URL: <http://www.jstor.org/stable/40511597>.
- Nestroy, O., Aust, G., Blum, W., Englisch, M., Hager, H., Herzberger, E., Kilian, W., Nelhiebel, P., G. Ortner and, E.P., und J. Wagner, A.P.W.S., 2011. Systematische gliederung der bden sterreichs. sterreichische boden-systematik 2000 in der revidierten fassung von 2011. Mitt. sterr. Bodenkdl. Ges. 79.

- 215 Riley, S.J., DeGloria, S.D., Elliot, R., 1999. A terrain ruggedness index that  
quantifies topographic heterogeneity. *Intermountain Journal of Sciences* 5,  
23 – 27.
- Sappington, J.M., Longshore, K.M., Thompson, D.B., 2007. Quantifying  
landscape ruggedness for animal habitat analysis: A case study using  
bighorn sheep in the mojave desert. *Journal of Wildlife Management* 75.  
URL: <http://dx.doi.org/10.2193/2005-723>, doi:10.2193/2005-723.
- 220 Scholz, H., Bestle, K.H., Willerich, S., 2005. Quartärgeologische untersuchun-  
gen im überetsch. *Geo.Alp* 2, 1–23.