

# Finding Partite Hypergraphs Efficiently

Ferran Espuña Bertomeu

Supervisor: Richard Lang

June 2025

- 1 Hypergraphs
- 2 Turán-Type Problems
- 3 Algorithms
- 4 Future Work

# $k$ -Graphs

## Definition

A  $k$ -graph is a pair  $G = (V, E)$  where  $V$  is a finite set of *vertices* and  $E \subseteq \binom{V}{k}$  is a set of *edges*.




Figure: A complete 3-graph on 4 vertices:  $K_4^{(3)}$ .

# Partite $k$ -Graphs

## Definition

A  $k$ -graph  $G = (V, E)$  is  $r$ -partite if there exists a partition  $V = V_1 \cup \dots \cup V_r$  such that every edge of  $G$  intersects every part  $V_i$  in at most one vertex. We write  $G = (V_1, \dots, V_r; E)$ .




Figure: A complete 3-partite 2-graph:  $K^{(2)}(2, 2, 2)$ .

# Partite $k$ -Graphs

## Remark

We may identify  $E$  as a subset of  $\mathcal{C} = \bigcup_{\{i_1, \dots, i_k\} \in \binom{[r]}{k}} V_{i_1} \times \dots \times V_{i_k}$ . If  $E = \mathcal{C}$ , we say that  $G$  is a *complete  $r$ -partite  $k$ -graph*.




Figure: A complete 3-partite 3-graph:  $K^{(3)}(2, 2, 2)$ .

# Turán-Type Problems

## Definition

Let  $G = (V, E)$  be a  $k$ -graph and  $n \geq |V|$  an integer.

The *Turán number*  $\text{ex}(n, G)$  is the maximum number of edges in a  $k$ -graph on  $n$  vertices that does not contain a copy of  $G$  as a subgraph.

Determining  $\text{ex}(n, G)$  or estimating it as  $n \rightarrow \infty$  is known as the *Turán problem* for  $G$ .

## Theorem

For all  $k$ -graphs  $G$  there exists a constant  $\alpha(G) \in [0, 1)$  such that

$$\text{ex}(n, G) = (\alpha(G) + o(1)) \cdot \binom{n}{k} \quad \text{as } n \rightarrow \infty.$$

Furthermore,  $\alpha(G) = 0$  if and only if  $G$  is  $k$ -partite.

# The Kővari–Sós–Turán Theorem

The bound  $\text{ex}(n, G) = o(n^k)$  can be improved by a lot.

## Definition

The *Zarankiewicz number*  $z(v_1, \dots, v_k; t_1, \dots, t_k)$  is the largest integer  $z$  for which there is a  $k$ -partite  $k$ -graph  $H = (V_1, \dots, V_k, F)$  with

- Part sizes  $|V_i| = v_i$
- $|F| = z$  edges
- No complete subgraph  $K(W_1, \dots, W_k)$  with  $W_i \subset V_i$  and  $|W_i| = t_i$ .


## Theorem (Kővari–Sós–Turán)

$$z(u, w; s, t) \leq (s-1)^{1/t}(w-t+1)u^{1-1/t} + (t-1)u$$

By a probabilistic argument, this implies that  $\text{ex}(n, K(s, t)) = \mathcal{O}(n^{2-1/t})$ .

# Kővari–Sós–Turán: Proof Sketch. $(u, s) = (5, 3); (w, t) = (4, 2)$

This graph has the maximum number of edges ( $|E| = 13$ ) to be  $K_{3,2}$ -free.




- **Hypothesis:**  $H = (U, W; E)$  is a  $K(s, t)$ -free bipartite  $k$ -graph with  $z = z(u, w; s, t)$  edges, where  $|U| = u$  and  $|W| = w$ .



# Kővari–Sós–Turán: Proof Sketch. $(u, s) = (5, 3); (w, t) = (4, 2)$

This graph has the maximum number of edges ( $|E| = 13$ ) to be  $K_{3,2}$ -free.




For example, adding the edge  $\{U_4, W_3\}$  creates a  $K_{3,2}$  on vertices  $\{U_1, U_2, U_4\}$  and  $\{W_2, W_3\}$ .

- Hypothesis:**  $H = (U, W; E)$  is a  $K(s, t)$ -free bipartite  $k$ -graph with  $z = z(u, w; s, t)$  edges, where  $|U| = u$  and  $|W| = w$ .

# Kővari–Sós–Turán: Proof Sketch. $(u, s) = (5, 3); (w, t) = (4, 2)$

This graph has the maximum number of edges ( $|E| = 13$ ) to be  $K_{3,2}$ -free.




For  $x = U_1$ , we count its  $\binom{4}{2} = 6$  stars.

- **Hypothesis:**  $H = (U, W; E)$  is a  $K(s, t)$ -free bipartite  $k$ -graph with  $z = z(u, w; s, t)$  edges, where  $|U| = u$  and  $|W| = w$ .
- **Counting Stars:** For each  $x \in U$ , there are  $\binom{d_H(x)}{t}$  sets  $T \subset W$  of  $t$  neighbors of  $x$ .

# Kővari–Sós–Turán: Proof Sketch. $(u, s) = (5, 3); (w, t) = (4, 2)$

This graph has the maximum number of edges ( $|E| = 13$ ) to be  $K_{3,2}$ -free.




For  $x = U_1$ , we count  
its  $\binom{4}{2} = 6$  stars.

- **Hypothesis:**  $H = (U, W; E)$  is a  $K(s, t)$ -free bipartite  $k$ -graph with  $z = z(u, w; s, t)$  edges, where  $|U| = u$  and  $|W| = w$ .
- **Counting Stars:** For each  $x \in U$ , there are  $\binom{d_H(x)}{t}$  sets  $T \subset W$  of  $t$  neighbors of  $x$ .

# Kővari–Sós–Turán: Proof Sketch. $(u, s) = (5, 3); (w, t) = (4, 2)$

This graph has the maximum number of edges ( $|E| = 13$ ) to be  $K_{3,2}$ -free.




For  $x = U_1$ , we count  
its  $\binom{4}{2} = 6$  stars.

- **Hypothesis:**  $H = (U, W; E)$  is a  $K(s, t)$ -free bipartite  $k$ -graph with  $z = z(u, w; s, t)$  edges, where  $|U| = u$  and  $|W| = w$ .
- **Counting Stars:** For each  $x \in U$ , there are  $\binom{d_H(x)}{t}$  sets  $T \subset W$  of  $t$  neighbors of  $x$ .

# Kővari–Sós–Turán: Proof Sketch. $(u, s) = (5, 3); (w, t) = (4, 2)$

This graph has the maximum number of edges ( $|E| = 13$ ) to be  $K_{3,2}$ -free.




For  $x = U_1$ , we count  
its  $\binom{4}{2} = 6$  stars.

- **Hypothesis:**  $H = (U, W; E)$  is a  $K(s, t)$ -free bipartite  $k$ -graph with  $z = z(u, w; s, t)$  edges, where  $|U| = u$  and  $|W| = w$ .
- **Counting Stars:** For each  $x \in U$ , there are  $\binom{d_H(x)}{t}$  sets  $T \subset W$  of  $t$  neighbors of  $x$ .

# Kővari–Sós–Turán: Proof Sketch. $(u, s) = (5, 3); (w, t) = (4, 2)$

This graph has the maximum number of edges ( $|E| = 13$ ) to be  $K_{3,2}$ -free.




For  $x = U_1$ , we count its  $\binom{4}{2} = 6$  stars.

- **Hypothesis:**  $H = (U, W; E)$  is a  $K(s, t)$ -free bipartite  $k$ -graph with  $z = z(u, w; s, t)$  edges, where  $|U| = u$  and  $|W| = w$ .
- **Counting Stars:** For each  $x \in U$ , there are  $\binom{d_H(x)}{t}$  sets  $T \subset W$  of  $t$  neighbors of  $x$ .

# Kővari–Sós–Turán: Proof Sketch. $(u, s) = (5, 3); (w, t) = (4, 2)$

This graph has the maximum number of edges ( $|E| = 13$ ) to be  $K_{3,2}$ -free.




For  $x = U_1$ , we count  
its  $\binom{4}{2} = 6$  stars.

- **Hypothesis:**  $H = (U, W; E)$  is a  $K(s, t)$ -free bipartite  $k$ -graph with  $z = z(u, w; s, t)$  edges, where  $|U| = u$  and  $|W| = w$ .
- **Counting Stars:** For each  $x \in U$ , there are  $\binom{d_H(x)}{t}$  sets  $T \subset W$  of  $t$  neighbors of  $x$ .

# Kővari–Sós–Turán: Proof Sketch. $(u, s) = (5, 3); (w, t) = (4, 2)$

This graph has the maximum number of edges ( $|E| = 13$ ) to be  $K_{3,2}$ -free.




In the example, there are at least  $5 \binom{13/5}{2} = 10.4$  stars (there are actually 12)

- **Hypothesis:**  $H = (U, W; E)$  is a  $K(s, t)$ -free bipartite  $k$ -graph with  $z = z(u, w; s, t)$  edges, where  $|U| = u$  and  $|W| = w$ .
- **Counting Stars:** For each  $x \in U$ , there are  $\binom{d_H(x)}{t}$  sets  $T \subset W$  of  $t$  neighbors of  $x$ .
- **Averaging:** By a convexity argument, the number of stars is at least  $u \binom{z/u}{t}$ .



# Kővari–Sós–Turán: Proof Sketch. $(u, s) = (5, 3); (w, t) = (4, 2)$

This graph has the maximum number of edges ( $|E| = 13$ ) to be  $K_{3,2}$ -free.




Each set  $T \subset W$  (in this case,  $T = \{W_1, W_2\}$ ) is in at most  $s - 1 = 3 - 1 = 2$  stars. In total, at most  $2\binom{4}{2} = 12$  stars.

- **Hypothesis:**  $H = (U, W; E)$  is a  $K(s, t)$ -free bipartite  $k$ -graph with  $z = z(u, w; s, t)$  edges, where  $|U| = u$  and  $|W| = w$ .
- **Counting Stars:** For each  $x \in U$ , there are  $\binom{d_H(x)}{t}$  sets  $T \subset W$  of  $t$  neighbors of  $x$ .
- **Averaging:** By a convexity argument, the number of stars is at least  $u\binom{z/u}{t}$ .
- **Bounding:** Because  $H$  is  $K(s, t)$ -free, each set  $T \subset W$  is the right component of at most  $(s - 1)$  stars.

# Kővari–Sós–Turán: Proof Sketch. $(u, s) = (5, 3); (w, t) = (4, 2)$

This graph has the maximum number of edges ( $|E| = 13$ ) to be  $K_{3,2}$ -free.




Each set  $T \subset W$  (in this case,  $T = \{W_1, W_2\}$ ) is in at most  $s - 1 = 3 - 1 = 2$  stars. In total, at most  $2\binom{4}{2} = 12$  stars.

- **Hypothesis:**  $H = (U, W; E)$  is a  $K(s, t)$ -free bipartite  $k$ -graph with  $z = z(u, w; s, t)$  edges, where  $|U| = u$  and  $|W| = w$ .
- **Counting Stars:** For each  $x \in U$ , there are  $\binom{d_H(x)}{t}$  sets  $T \subset W$  of  $t$  neighbors of  $x$ .
- **Averaging:** By a convexity argument, the number of stars is at least  $u\binom{z/t}{t}$ .
- **Bounding:** Because  $H$  is  $K(s, t)$ -free, each set  $T \subset W$  is the right component of at most  $(s - 1)$  stars.

# Kővari–Sós–Turán: Proof Sketch. $(u, s) = (5, 3); (w, t) = (4, 2)$

This graph has the maximum number of edges ( $|E| = 13$ ) to be  $K_{3,2}$ -free.




Each set  $T \subset W$  (in this case,  $T = \{w_1, w_2\}$ ) is in at most  $s - 1 = 3 - 1 = 2$  stars. In total, at most  $2\binom{4}{2} = 12$  stars.

- **Hypothesis:**  $H = (U, W; E)$  is a  $K(s, t)$ -free bipartite  $k$ -graph with  $z = z(u, w; s, t)$  edges, where  $|U| = u$  and  $|W| = w$ .
- **Counting Stars:** For each  $x \in U$ , there are  $\binom{d_H(x)}{t}$  sets  $T \subset W$  of  $t$  neighbors of  $x$ .
- **Averaging:** By a convexity argument, the number of stars is at least  $u\binom{z/t}{t}$ .
- **Bounding:** Because  $H$  is  $K(s, t)$ -free, each set  $T \subset W$  is the right component of at most  $(s - 1)$  stars.

# Kővari–Sós–Turán: Proof Sketch. $(u, s) = (5, 3); (w, t) = (4, 2)$

This graph has the maximum number of edges ( $|E| = 13$ ) to be  $K_{3,2}$ -free.



In the example, we conclude that  $10.4 \leq 12$ , which is true. For bigger values of  $z$  this would fail, leading to contradiction and therefore upper bounding  $z$ .

- **Hypothesis:**  $H = (U, W; E)$  is a  $K(s, t)$ -free bipartite  $k$ -graph with  $z = z(u, w; s, t)$  edges, where  $|U| = u$  and  $|W| = w$ .
- **Counting Stars:** For each  $x \in U$ , there are  $\binom{d_H(x)}{t}$  sets  $T \subset W$  of  $t$  neighbors of  $x$ .
- **Averaging:** By a convexity argument, the number of stars is at least  $u \binom{z/u}{t}$ .
- **Bounding:** Because  $H$  is  $K(s, t)$ -free, each set  $T \subset W$  is the right component of at most  $(s - 1)$  stars.
- **Conclusion:**  $u \binom{z/u}{t} \leq (s - 1) \binom{w}{t}$ , from which the theorem follows.

# Erdős's Bound for Hypergraphs (1964)

## Theorem (Erdős '64)

For integers  $k \geq 2, t \geq 2$ ,  $ex(n, K(t, \overset{k}{\cdot}, t)) = \mathcal{O}\left(n^{k - \frac{1}{t^{k-1}}}\right)$ .

This generalizes the Kővari–Sós–Turán theorem to  $k$ -graphs.

It follows from a similar bound on the corresponding generalized Zarankiewicz number, obtained by induction.

Suppose that  $H = (V_1, \dots, V_k; F)$  is a  $k$ -graph with  $|W_i| = w$ . Let  $H$  have  $z$  edges and no copy of  $K(t, \overset{k}{\cdot}, t)$ .



We set up a bipartite  $k$ -graph  $H' = (U, W; F')$  with

$$U = W_1 \times \dots \times W_{k-1}$$



$$W = W_k$$

$$F' = \{(X, y) \in U \times W \mid X \cup \{y\} \in F\}.$$



# Erdős's Bound: Proof Sketch ( $k = 3, t = 2$ )



# Erdős's Bound: Proof Sketch ( $k = 3, t = 2$ )





# Erdős's Bound: Proof Sketch ( $k = 3, t = 2$ )









# Erdős's Bound: Proof Sketch ( $k = 3, t = 2$ )





# Erdős's Bound: Proof Sketch ( $k = 3, t = 2$ )





# Erdős's Bound: Proof Sketch ( $k = 3, t = 2$ )





# Erdős's Bound: Proof Sketch ( $k = 3, t = 2$ )





# Erdős's Bound: Proof Sketch ( $k = 3, t = 2$ )





# Erdős's Bound: Proof Sketch ( $k = 3, t = 2$ )



# Erdős's Bound: Proof Sketch ( $k = 3, t = 2$ )



# Erdős's Bound: Proof Sketch ( $k = 3, t = 2$ )





# Implications of Erdős's Bound

Doing the calculations more carefully, we can show:

## Theorem

*Let  $k \geq 2$  and  $d > 0$ . Then there exists a constant  $\delta = \delta(k, d) > 0$  such that every  $k$ -graph  $G$  with  $n$  vertices and  $dn^k$  edges contains  $K(t, \dots, t)$  with*

$$t \geq \delta \cdot (\log n)^{1/(k-1)}.$$

# Implications of Erdős's Bound

Doing the calculations more carefully, we can show:

## Theorem

*Let  $k \geq 2$  and  $d > 0$ . Then there exists a constant  $\delta = \delta(k, d) > 0$  such that every  $k$ -graph  $G$  with  $n$  vertices and  $dn^k$  edges contains  $K(t, \dots, t)$  with*

$$t \geq \delta \cdot (\log n)^{1/(k-1)}.$$

That is,  $k$ -graphs of constant density contain complete balanced  $k$ -partite  $k$ -subgraphs of **growing** part size.

# Implications of Erdős's Bound

Doing the calculations more carefully, we can show:

## Theorem

*Let  $k \geq 2$  and  $d > 0$ . Then there exists a constant  $\delta = \delta(k, d) > 0$  such that every  $k$ -graph  $G$  with  $n$  vertices and  $dn^k$  edges contains  $K(t, \dots, t)$  with*

$$t \geq \delta \cdot (\log n)^{1/(k-1)}.$$

That is,  $k$ -graphs of constant density contain complete balanced  $k$ -partite  $k$ -subgraphs of **growing** part size.

A probabilistic argument shows that this is tight up to the constant  $\delta(k, d)$ .

# The Algorithmic problem

The proofs shown are **non-constructive**.

# The Algorithmic problem

The proofs shown are **non-constructive**.

We want to **find**  $K(t, \dots, t)$  in  $H = (V, E)$  of constant density  $d = m/n^k$ .

# The Algorithmic problem

The proofs shown are **non-constructive**.

We want to **find**  $K(t, \dots, t)$  in  $H = (V, E)$  of constant density  $d = m/n^k$ .

Naïve approach: enumerate all  $k$ -partite  $k$ -subgraphs of  $H$ .

# The Algorithmic problem

The proofs shown are **non-constructive**.

We want to **find**  $K(t, \dots, t)$  in  $H = (V, E)$  of constant density  $d = m/n^k$ .

Naïve approach: enumerate all  $k$ -partite  $k$ -subgraphs of  $H$ .

**Problem:** There are on the order of

$$\binom{n}{t} \geq \binom{n}{\delta(\log n)^{1/(k-1)}}$$

such sets, which is **not** polynomial in  $n$ .

# The Algorithmic problem

The proofs shown are **non-constructive**.

We want to **find**  $K(t, \dots, t)$  in  $H = (V, E)$  of constant density  $d = m/n^k$ .

Naïve approach: enumerate all  $k$ -partite  $k$ -subgraphs of  $H$ .

**Problem:** There are on the order of

$$\binom{n}{t} \geq \binom{n}{\delta(\log n)^{1/(k-1)}}$$

such sets, which is **not** polynomial in  $n$ .

Can we do it in polynomial time?



# An Algorithm for $k = 2$ (D. Mubayi and G. Turán)

**Idea:** Mimick the proof the Kővari–Sós–Turán theorem:

# An Algorithm for $k = 2$ (D. Mubayi and G. Turán)

**Idea:** Mimick the proof the Kővari–Sós–Turán theorem:

- Find a bipartite subgraph  $(U, W; E)$  of  $H$  with many edges.

# An Algorithm for $k = 2$ (D. Mubayi and G. Turán)

**Idea:** Mimick the proof the Kővari–Sós–Turán theorem:

- Find a bipartite subgraph  $(U, W; E)$  of  $H$  with many edges.
- For each subset  $T \in \binom{W}{t}$ , calculate the common neighborhood. For some  $T$  it has size  $s = t$ .

# An Algorithm for $k = 2$ (D. Mubayi and G. Turán)

**Idea:** Mimick the proof the Kővari–Sós–Turán theorem:

- Find a bipartite subgraph  $(U, W; E)$  of  $H$  with many edges.
- For each subset  $T \in \binom{W}{t}$ , calculate the common neighborhood. For some  $T$  it has size  $s = t$ .

**Problems:**

# An Algorithm for $k = 2$ (D. Mubayi and G. Turán)

**Idea:** Mimick the proof the Kővari–Sós–Turán theorem:

- Find a bipartite subgraph  $(U, W; E)$  of  $H$  with many edges.
- For each subset  $T \in \binom{W}{t}$ , calculate the common neighborhood. For some  $T$  it has size  $s = t$ .

**Problems:**

- Finding a dense bipartite subgraph.

# An Algorithm for $k = 2$ (D. Mubayi and G. Turán)

**Idea:** Mimick the proof the Kővari–Sós–Turán theorem:

- Find a bipartite subgraph  $(U, W; E)$  of  $H$  with many edges.
- For each subset  $T \in \binom{W}{t}$ , calculate the common neighborhood. For some  $T$  it has size  $s = t$ .

**Problems:**

- Finding a dense bipartite subgraph.
- If we are not careful, the search space  $\binom{W}{t}$  might still be too large ...

# An Algorithm for $k = 2$ (D. Mubayi and G. Turán)

**Idea:** Mimick the proof the Kővari–Sós–Turán theorem:

- Find a bipartite subgraph  $(U, W; E)$  of  $H$  with many edges.
- For each subset  $T \in \binom{W}{t}$ , calculate the common neighborhood. For some  $T$  it has size  $s = t$ .

**Problems:**

- Finding a dense bipartite subgraph.
- If we are not careful, the search space  $\binom{W}{t}$  might still be too large ...

**Solution:**  $W$  are the  $\mathbf{w} = \lfloor \mathbf{t}/\mathbf{d} \rfloor$  vertices of **maximum degree**.  $U = V \setminus W$ .

# An Algorithm for $k = 2$ (D. Mubayi and G. Turán)

**Idea:** Mimick the proof the Kővari–Sós–Turán theorem:

- Find a bipartite subgraph  $(U, W; E)$  of  $H$  with many edges.
- For each subset  $T \in \binom{W}{t}$ , calculate the common neighborhood. For some  $T$  it has size  $s = t$ .

**Problems:**

- Finding a dense bipartite subgraph.
- If we are not careful, the search space  $\binom{W}{t}$  might still be too large ...

**Solution:**  $W$  are the  $\mathbf{w} = \lfloor \mathbf{t}/\mathbf{d} \rfloor$  vertices of **maximum degree**.  $U = V \setminus W$ .

- Max degree  $\implies$  The bipartite subgraph has many edges.



# An Algorithm for $k = 2$ (D. Mubayi and G. Turán)

**Idea:** Mimick the proof the Kővari–Sós–Turán theorem:

- Find a bipartite subgraph  $(U, W; E)$  of  $H$  with many edges.
- For each subset  $T \in \binom{W}{t}$ , calculate the common neighborhood. For some  $T$  it has size  $s = t$ .

**Problems:**

- Finding a dense bipartite subgraph.
- If we are not careful, the search space  $\binom{W}{t}$  might still be too large ...

**Solution:**  $W$  are the  $\mathbf{w} = \lfloor \mathbf{t}/\mathbf{d} \rfloor$  vertices of **maximum degree**.  $U = V \setminus W$ .

- Max degree  $\implies$  The bipartite subgraph has many edges.
- $w$  is “just right”. If too small, the extra term in the  $KST$  bound is too large; if too big, the search space  $\binom{W}{t}$  is too large.

## Our Contribution: Extending to $k$ -Graphs

We present a polynomial algorithm that finds a  $K(t, \dots, t)$  in a  $k$ -graph  $H = (V, E)$  with  $dn^k$  edges, where

$$t = \left\lfloor \left( \frac{\log(n/2^{(k-1)})}{\log(3/d)} \right)^{\frac{1}{k-1}} \right\rfloor$$

- **First (failed) attempt:** Try to find a  $k$ -partite subgraph of  $H$  with logarithmically sized parts except one large part.

## Our Contribution: Extending to $k$ -Graphs

We present a polynomial algorithm that finds a  $K(t, \dots, t)$  in a  $k$ -graph  $H = (V, E)$  with  $dn^k$  edges, where

$$t = \left\lfloor \left( \frac{\log(n/2^{(k-1)})}{\log(3/d)} \right)^{\frac{1}{k-1}} \right\rfloor$$

- **First (failed) attempt:** Try to find a  $k$ -partite subgraph of  $H$  with logarithmically sized parts except one large part.
- **Actual solution:** Use a **recursive** approach from the beginning.

## Our Contribution: Extending to $k$ -Graphs

We present a polynomial algorithm that finds a  $K(t, \dots, t)$  in a  $k$ -graph  $H = (V, E)$  with  $dn^k$  edges, where

$$t = \left\lfloor \left( \frac{\log(n/2^{(k-1)})}{\log(3/d)} \right)^{\frac{1}{k-1}} \right\rfloor$$

- **First (failed) attempt:** Try to find a  $k$ -partite subgraph of  $H$  with logarithmically sized parts except one large part.
- **Actual solution:** Use a **recursive** approach from the beginning.
  - Find a set  $W$  of  $\mathbf{w} = \lfloor 2\mathbf{t}/\mathbf{d} \rfloor$  vertices of maximum degree.

## Our Contribution: Extending to $k$ -Graphs

We present a polynomial algorithm that finds a  $K(t, \dots, t)$  in a  $k$ -graph  $H = (V, E)$  with  $dn^k$  edges, where

$$t = \left\lfloor \left( \frac{\log(n/2^{(k-1)})}{\log(3/d)} \right)^{\frac{1}{k-1}} \right\rfloor$$

- **First (failed) attempt:** Try to find a  $k$ -partite subgraph of  $H$  with logarithmically sized parts except one large part.
- **Actual solution:** Use a **recursive** approach from the beginning.
  - Find a set  $W$  of  $\mathbf{w} = \lfloor 2\mathbf{t}/\mathbf{d} \rfloor$  vertices of maximum degree.
  - There are many edges with exactly 1 vertex in  $W$  (by averaging).

## Our Contribution: Extending to $k$ -Graphs

We present a polynomial algorithm that finds a  $K(t, \dots, t)$  in a  $k$ -graph  $H = (V, E)$  with  $dn^k$  edges, where

$$t = \left\lfloor \left( \frac{\log(n/2^{(k-1)})}{\log(3/d)} \right)^{\frac{1}{k-1}} \right\rfloor$$

- **First (failed) attempt:** Try to find a  $k$ -partite subgraph of  $H$  with logarithmically sized parts except one large part.
- **Actual solution:** Use a **recursive** approach from the beginning.
  - Find a set  $W$  of  $\mathbf{w} = \lfloor 2\mathbf{t}/\mathbf{d} \rfloor$  vertices of maximum degree.
  - There are many edges with exactly 1 vertex in  $W$  (by averaging).
  - By  $KST$ , for some  $T \subset W$  of size  $t$ , the set  $L_T$  of  $(k-1)$ -sets of  $V \setminus W$  forming an edge with all vertices in  $T$  has size at least  $\mathbf{s} = \mathbf{d}^t \mathbf{n}^{k-1}$ .

# Our Contribution: Extending to $k$ -Graphs

We present a polynomial algorithm that finds a  $K(t, \dots, t)$  in a  $k$ -graph  $H = (V, E)$  with  $dn^k$  edges, where

$$t = \left\lfloor \left( \frac{\log(n/2^{(k-1)})}{\log(3/d)} \right)^{\frac{1}{k-1}} \right\rfloor$$

- **First (failed) attempt:** Try to find a  $k$ -partite subgraph of  $H$  with logarithmically sized parts except one large part.
- **Actual solution:** Use a **recursive** approach from the beginning.
  - Find a set  $W$  of  $\mathbf{w} = \lfloor 2\mathbf{t}/\mathbf{d} \rfloor$  vertices of maximum degree.
  - There are many edges with exactly 1 vertex in  $W$  (by averaging).
  - By  $KST$ , for some  $T \subset W$  of size  $t$ , the set  $L_T$  of  $(k-1)$ -sets of  $V \setminus W$  forming an edge with all vertices in  $T$  has size at least  $\mathbf{s} = \mathbf{d}^t \mathbf{n}^{k-1}$ .
  - Applying the algorithm to  $H' = (V \setminus W, S)$  with  $\mathbf{n}' = \mathbf{n} - \mathbf{w}$ ,  $\mathbf{d}' = \mathbf{d}^t$  we get  $\mathbf{t}' \geq \mathbf{t}$ .

## Our Contribution: Extending to $k$ -Graphs

We present a polynomial algorithm that finds a  $K(t, \dots, t)$  in a  $k$ -graph  $H = (V, E)$  with  $dn^k$  edges, where

$$t = \left\lfloor \left( \frac{\log(n/2^{(k-1)})}{\log(3/d)} \right)^{\frac{1}{k-1}} \right\rfloor$$

- **First (failed) attempt:** Try to find a  $k$ -partite subgraph of  $H$  with logarithmically sized parts except one large part.
- **Actual solution:** Use a **recursive** approach from the beginning.
  - Find a set  $W$  of  $\mathbf{w} = \lfloor 2\mathbf{t}/\mathbf{d} \rfloor$  vertices of maximum degree.
  - There are many edges with exactly 1 vertex in  $W$  (by averaging).
  - By *KST*, for some  $T \subset W$  of size  $t$ , the set  $L_T$  of  $(k-1)$ -sets of  $V \setminus W$  forming an edge with all vertices in  $T$  has size at least  $\mathbf{s} = \mathbf{d}^t \mathbf{n}^{k-1}$ .
  - Applying the algorithm to  $H' = (V \setminus W, S)$  with  $\mathbf{n}' = \mathbf{n} - \mathbf{w}$ ,  $\mathbf{d}' = \mathbf{d}^t$  we get  $\mathbf{t}' \geq \mathbf{t}$ .
  - It finds  $\mathbf{T}_1, \dots, \mathbf{T}_{k-1}$  complete in  $H' \implies \mathbf{T}_1, \dots, \mathbf{T}_{k-1}, \mathbf{T}$  complete in  $H$ .



# Future Work

# Future Work

- **Refine the algorithm:** The current analysis is not tight. The algorithm is not yet practical.

# Future Work

- **Refine the algorithm:** The current analysis is not tight. The algorithm is not yet practical.
- **Implement and test:** Evaluate the algorithm on synthetic and real-world hypergraphs.

# Future Work

- **Refine the algorithm:** The current analysis is not tight. The algorithm is not yet practical.
- **Implement and test:** Evaluate the algorithm on synthetic and real-world hypergraphs.
- **Generalize the structures found:** The algorithm finds complete balanced  $k$ -partite  $k$ -graphs. Extending it to unbalanced seems feasible.

# Future Work

- **Refine the algorithm:** The current analysis is not tight. The algorithm is not yet practical.
- **Implement and test:** Evaluate the algorithm on synthetic and real-world hypergraphs.
- **Generalize the structures found:** The algorithm finds complete balanced  $k$ -partite  $k$ -graphs. Extending it to unbalanced seems feasible.
- **Generalize the blow-ups:** The algorithm finds  $t_n$ -blowups of a single edge. Obvious way to extend it does not yield optimal  $t$ .

# Future Work

- **Refine the algorithm:** The current analysis is not tight. The algorithm is not yet practical.
- **Implement and test:** Evaluate the algorithm on synthetic and real-world hypergraphs.
- **Generalize the structures found:** The algorithm finds complete balanced  $k$ -partite  $k$ -graphs. Extending it to unbalanced seems feasible.
- **Generalize the blow-ups:** The algorithm finds  $t_n$ -blowups of a single edge. Obvious way to extend it does not yield optimal  $t$ .
- **Extremely dense hypergraphs:** The algorithm does not yield the best order of  $t$  for increasing  $d \rightarrow 1/k!$ .

# Thank You

# Questions?

---

Ferran Espuña Bertomeu  
ferranespuna@gmail.com