Problem 4: Let $T_{d,k}$ be a (d-1)-ary rooted tree of height k. Show that

- (a) $\lambda_1(T_{d,k}) \le 2\sqrt{d-1}$
- (b) $\lim_{k\to\infty} \lambda_1(T_{d,k}) = 2\sqrt{d-1}$

Solution (by Ferran Espuña): Let A be the adjacency matrix of $T_{d,k}$. For example, if d=3 and k=2,

$$A = \begin{pmatrix} 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \end{pmatrix}$$

where the first row corresponds to the root of the tree, the second and third rows correspond to its children and the other four rows correspond to the leaves. Now we can rephrase (a) as the following:

Proposition. $||A||_2 \leq 2\sqrt{d-1}$.

Proof. We will use this famous inequality from linear algebra: $||B||_2 \le \sqrt{||B||_1 \cdot ||B||_{\infty}}$. In this case, we will apply this to powers of A, which is non-negative and symmetric. Therefore, for $B = A^p$

$$||B||_1 = ||B||_{\infty} = \max_i \sum_j B_{ij}$$

so in particular

(1)
$$||A||_2^p = ||A^p||_2 \le \sqrt{||A^p||_1 \cdot ||A^p||_{\infty}} = \max_i \sum_i A_{ij}^p$$

Which is just a row sum of A^p . This corresponds to the number $P_{i,p}$ of paths of length p from a given vertex v_i . Let's try to approximate this number.

At each step of the path, the height (distance from the root) either increases by 1 (we say we go down) or decreases by 1 (we say we go up). Because both at the start and end of the path the height must be in the range [0, k], the difference betwen the number of up steps s_u and the number of down steps s_d must satisfy

$$|s_u - s_d| \le k$$

and obviously

$$s_u + s_d = p$$

All together,

$$\frac{p}{2} - k \le s_d \le \frac{p}{2} + k$$

If we go up, there is only one choice (the parent of the current node). If we go down, there are at most d-1 choices (the children of the current node). Furtheremore, for each choice of s_d there are $\binom{p}{s_d}$ ways to choose

the down steps. Note that not all the ways of choosing are valid, as we must also ensure that the height stays within the range [0, k] during the whole path. All in all, for all i,

$$P_{i,p} \le \sum_{s_d = \frac{p}{2} - k}^{\frac{p}{2} + k} \binom{p}{s_d} (d - 1)^{s_d} \le 2 \cdot k \cdot 2^p \cdot (d - 1)^{\frac{p}{2}}$$

Putting this together with (1), we get

$$||A||_2^p \le 2 \cdot k \cdot 2^p \cdot (d-1)^{\frac{p}{2}} \implies ||A||_2 \le \lim_{p \to \infty} \left(2 \cdot k \cdot 2^p \cdot (d-1)^{\frac{p}{2}} \right)^{\frac{1}{p}} = 2\sqrt{d-1}$$

Similarly, we can rephrase (b) as the following:

Proposition. $\lim_{k\to\infty} ||A(k)|| = 2\sqrt{d-1}$.

Proof. We would now like to bound below the 2-norm of A(k), so will use the inequality $||B||_2 \ge \frac{1}{\sqrt{n}} ||B||_{\infty}$, where n is the size of the matrix. We repeat the same calculation for the number of paths of length p from a given vertex v_i , but now we need to bound it below. Because of the maximum in the formula in the ∞ -norm, we're free to choose the startring vertex v_i . We will choose the root. We're also free to choose p = 2k. If we restrict ourselves to paths that come back to the root, we get $s_d = s_u = k$. The number of valid combinations of up and down steps is the corresponding $Catalan\ number\ C_k = \frac{1}{k+1} {2k \choose k}$. All in all,

$$||A(k)||_2^{2k} = ||A(k)||_2^{2k} = \frac{1}{\sqrt{n}} P_{1,2k} \ge \frac{1}{\sqrt{n}} C_k (d-1)^k \ge \frac{1}{(k+1)\sqrt{n}} {2k \choose k} (d-1)^k \ge \frac{4^k}{(k+1)2\sqrt{kn}} (d-1)^k$$

Which implies

$$||A(k)||_2 \ge \lim_{k \to \infty} \left(\frac{4^k}{(k+1)2\sqrt{kn}} (d-1)^k \right)^{\frac{1}{2k}} = 2\sqrt{d-1}$$