

Maximum length of a 3-sum-free set consisting of 2 intervals

Ferran Espuña

June 22, 2026

Theorem 1. *Let A be a 3-sum-free subset of $\mathbb{T}$ consisting of 2 disjoint closed intervals. with measure at least $\frac{1}{5}$. Then, $A + A - 3 \cdot A = \mathbb{T}$.*

Proof. Note that if the statement is true for a set A , then it is also true for any translate of A and for the set $-A$. Without loss of generality, we can therefore assume $A = [0, \alpha] \cup [x, x + \beta]$ satisfying the following conditions:

$$0 < \alpha < x \leq x + \beta < 1, \quad (1)$$

$$\alpha \geq \beta, \quad (2)$$

$$x - \alpha \leq 1 - (x + \beta), \quad \text{i.e.} \quad x \leq \frac{1 + \alpha - \beta}{2}. \quad (3)$$

Furthermore, by monotonicity, we can assume that $\alpha + \beta = \frac{1}{5}$, and that therefore by (2), $\alpha \geq \frac{1}{10}$ and $\beta \leq \frac{1}{10}$. Let us denote $I = [0, \alpha]$ and $J = [x, x + \beta]$. We will prove that $\mathbb{T}$ is covered by the following 5 intervals:

$$R_1 = I + I - 3 \cdot I = [-3\alpha, 2\alpha],$$

$$R_2 = I + J - 3 \cdot I = [-3\alpha + x, \alpha + x + \beta],$$

$$R_3 = J + J - 3 \cdot I = [2x - 3\alpha, 2x + 2\beta],$$

$$R_4 = I + I - 3 \cdot J = [-3x - 3\beta, -3x + 2\alpha],$$

$$R_5 = I + J - 3 \cdot J = [-2x - 3\beta, \alpha + \beta - 2x].$$

Note that all of these intervals as written are of the form $[a, b]$ where $0 < b - a < 1$, (we say that the intervals are in *proper* form) and therefore checking if a point $y \in \mathbb{T}$ is contained in one of these intervals is equivalent to checking if for some integer k , $y + k \in [a, b]$.

We denote $U = R_1 \cup R_2 \cup R_3 \cup R_4 \cup R_5$ and $V = \mathbb{T} \setminus U$. We assume for the sake of contradiction that V is non-empty. Note that $V \subset \mathbb{T} \setminus R_1 = (2\alpha, 1 - 3\alpha)$, which again is in proper form. The rest of the proof consists in iteratively shrinking this interval through a series of claims on the intervals R_i until we show that V is empty. The first claim is the following:

Claim 1. $[2\alpha, \alpha + x + \beta] \subset R_2$.

Proof. We just need to show that $2\alpha \in R_2$. In fact, we prove that

$$-3\alpha + x \leq 2\alpha \leq \alpha + x + \beta. \quad (4)$$

The first inequality is equivalent to $x \leq 5\alpha$. By assumption (3), we have

$$x \leq \frac{1 + \alpha - \beta}{2} \leq \frac{1 + \alpha - (1/5 - \alpha)}{2} = \alpha + \frac{2}{5}.$$

and since $\alpha \geq \frac{1}{10}$, we have $\alpha + \frac{2}{5} \leq 5\alpha$. The second inequality is equivalent to $\alpha \leq x + \beta$, which is true by assumption (1). $\square$

We can then conclude that

$$V \subset (\alpha + x + \beta, 1 - 3\alpha). \quad (5)$$

Furthermore, since we assuming V is non-empty, we get that $1 - 3\alpha \notin R_2$, and therefore by inequalities (4), together with the fact that $1 - 3\alpha > x - 3\alpha$, we have that

$$1 - 3\alpha > \alpha + x + \beta \quad \text{i.e.} \quad x < 1 - 4\alpha - \beta. \quad (6)$$

In particular, the interval in (5) is in proper form. We also note that from inequality (6) we also get the weaker but more convenient

$$x < 1 - (\alpha + \beta) - 3\alpha \leq 1 - 1/5 - 3/10 = 1/2. \quad (7)$$

Claim 2. $[\alpha + \beta + x, 2x + 2\beta] \subset R_3$.

Proof. Similarly to the previous claim, it suffices to show that

$$2x - 3\alpha \leq \alpha + \beta + x \leq 2x + 2\beta. \quad (8)$$

The first inequality is equivalent to $x \leq 4\alpha + \beta$. From condition (3) and the fact that $\alpha + \beta = \frac{1}{5}$, we have

$$x \leq \frac{1 + \alpha - \beta}{2} = \frac{\alpha - \beta + 5(\alpha + \beta)}{2} = 3\alpha + 2\beta. \quad (9)$$

But since $\alpha \geq \beta$, we have $3\alpha + 2\beta \leq 4\alpha + \beta$, and we are done.

The second inequality is equivalent to $\alpha \leq x + \beta$, which is true by assumption (1). $\square$

The conclusion of this is that in fact

$$V \subset (2x + 2\beta, 1 - 3\alpha). \quad (10)$$

Similarly as before, we are assuming V is non-empty and therefore $1 - 3\alpha \notin R_3$. By inequalities (8), and since $1 - 3\alpha > 2x - 3\alpha$ by (7), we have that

$$1 - 3\alpha > 2x + 2\beta \quad \text{i.e.} \quad x < \frac{1 - 3\alpha - 2\beta}{2}. \quad (11)$$

and that the interval in (10) is in proper form. Again, we can obtain a weaker but more convenient inequality from (11):

$$x < \frac{1 - 3\alpha - 2\beta}{2} \leq \frac{1 - 2/5 - 1/10}{2} = 1/4. \quad (12)$$

The proof will be completed once we show that R_4 and R_5 cover the interval $(2x + 2\beta, 1 - 3\alpha)$. We first show that the union of these two intervals is itself an interval:

Claim 3. *The ends of the intervals R_4 and R_5 satisfy the following inequalities:*

$$-3x - 3\beta \leq -2x - 3\beta \leq -3x + 2\alpha \leq \alpha + \beta - 2x.$$

and therefore

$$R_4 \cup R_5 = [-3x - 3\beta, \alpha + \beta - 2x]. \quad (13)$$

Furthermore, this interval is in proper form.

Proof. The first inequality is equivalent to $x \geq 0$, which is true by assumption (1). The second inequality is equivalent to $x \leq 2\alpha + 3\beta$, which is precisely inequality (9). The third inequality is equivalent to $\alpha \leq x + \beta$, which is true by assumption (1). To see that it is in proper form, note that since both R_4 and R_5 are in proper form, and since they have interleaving ends (the middle inequality above), we have that

$$\begin{aligned} & (\alpha + \beta - 2x) - (-3x - 3\beta) \\ & \leq ((\alpha + \beta - 2x) - (-2x - 3\beta)) + ((-3x + 2\alpha) - (-3x - 3\beta)) \\ & = (\alpha + 4\beta) + (2\alpha + 3\beta) = 3\alpha + 7\beta \leq 5(\alpha + \beta) = 1. \end{aligned} \quad \square$$

The end is near. We only need to show that the interval in (13) covers the interval in (10):

Claim 4. $R_4 \cup R_5 \supset (2x + 2\beta, 1 - 3\alpha)$.

Proof. We have calculated the ends of both intervals in proper form. Therefore, it suffices to show that

$$-3x - 3\beta \leq (2x + 2\beta) - 1 \leq (1 - 3\alpha) - 1 \leq \alpha + \beta - 2x,$$

where we have shifted both points to check by the integer -1 . We have already established the middle inequality. The first inequality is equivalent to

$5(x + \beta) \geq 1$, which is true because $x + \beta \geq \alpha + \beta = 1/5$. The last inequality is equivalent to $2x \leq 4\alpha + \beta = 3\alpha + 1/5$. Since $x \leq 1/4$ by (12), and we have $\alpha \geq 1/10$, we get

$$2x \leq 1/2 = 3/10 + 1/5 \leq 3\alpha + 1/5,$$

and we are done. □

We have therefore shown that V is empty, so the proof is complete. □