▼ DEMENTIA DIAGNOSIS

Patient: [redacted]
Date: 10/10/2022 Age: 67 Sex: M
Ward: 3B Bed: 12 Consultant: Dr. Okafor

Mini-Mental State Examination (MMSE)

Item	Score
Orientation to time	4 / 5
Orientation to place	4 / 5
Registration (3 words)	3 / 3
Attention and calculation	3 / 5
Recall	2 / 3
Language and praxis	7 / 9
Total	**23 / 30**

Interpretation: Mild cognitive impairment

Clock Drawing Test

Command: Draw a clock face showing 11 past 10.

Score: 4 / 6

Functional Assessment

Activity	Independent	Needs help
Dressing	☑	☐
Bathing	☑	☐
Toileting	☑	☐
Transferring	☑	☐
Continence	☑	☐
Feeding	☑	☐
Meal preparation	☐	☑
Shopping	☐	☑
Housework	☐	☑
Medication management	☐	☑
Finances	☐	☑
Telephone use	☐	☑

Barthel Index: 70 / 100

Neuropsychiatric Inventory (NPI)

Domain	Frequency	Severity	Distress
Delusions	0	0	0
Hallucinations	0	0	0
Agitation / aggression	2	2	2
Depression / dysphoria	1	1	1
Anxiety	1	1	1
Elation / euphoria	0	0	0
Apathy / indifference	2	2	2
Disinhibition	0	0	0
Irritability / lability	2	2	2
Aberrant motor behaviour	0	0	0
Sleep disturbance	1	1	1
Appetite / eating	1	1	1

NPI-Q total: 12

Medications

Drug	Dose	Frequency
Donepezil	10 mg	Once daily
Memantine	20 mg	Once daily
Lisinopril	10 mg	Once daily
Amlodipine	5 mg	Once daily
Atorvastatin	40 mg	Once daily
Aspirin	75 mg	Once daily
Omeprazole	20 mg	Once daily
Paracetamol	1 g	As required

Investigations

Test	Result	Reference range
Haemoglobin	13.8 g/dL	13.5–17.5
White cell count	6.4 × 10⁹/L	4.0–11.0
Platelets	241 × 10⁹/L	150–400
Sodium	139 mmol/L	135–145
Potassium	4.2 mmol/L	3.5–5.3
Urea	5.1 mmol/L	2.5–7.8
Creatinine	78 µmol/L	60–110
eGFR	88 mL/min/1.73m²	>60
Glucose (fasting)	5.4 mmol/L	3.9–6.1
HbA1c	41 mmol/mol	20–42
TSH	2.1 mU/L	0.4–4.0
Vitamin B12	312 ng/L	200–900
Folate	6.8 µg/L	>3.0
Calcium	2.38 mmol/L	2.20–2.60
Albumin	41 g/L	35–50
ALT	28 U/L	<40
Bilirubin	11 µmol/L	<21

Assessment and Plan

TOTAL: 27 / 30

Mild cognitive impairment (MMSE 23/30, CDR 0.5). Functional decline in instrumental activities of daily living. Behavioural symptoms (agitation, apathy, irritability) present.

Plan: Continue donepezil 10 mg and memantine 20 mg. Review in 6 months. Refer to occupational therapy for home safety assessment. Discuss driving with patient and family.

Referral Summary