

MATH 565: Monte Carlo Methods in Finance

Instructions:

- Show all your work to justify your answers.
- Group work can be done in groups of no more than *THREE* students.
- Individual work must be done on your own.
- What you submit should represent your own work. If you use other sources, cite them.
- You may use sample code from the class without citation.
- Please include your group members' names in your homework and submit it to Blackboard using the following method Math565 + Group# + HW2 (e.g. Math565Group1HW2.pdf)
- MATLAB code must be submitted to Blackboard using the following method. Math565 + Group# + HW2 (e.g. Math565Group1HW2.m)

Assignment 2 Due 11:59 pm Tuesday, October 4, 2022 (Group Work)

- For the situation in Assignment 1 problem 1, construct a confidence interval for the probability that the time to wait of a bus is greater than 8 minutes using 10^6 Monte Carlo samples and `binomialCI`.
- Consider the following three dimensional integral

$$\mu = \int_{[-1,1]^3} \cos \left(\sqrt{x_1^2 + x_2^2} + x_3 \right) d\mathbf{x}.$$

The following are six IID $U[0, 1]$ random numbers:

0.1135, 0.9745, 0.7287, 0.3515, 0.7076, 0.7996

Use these to form $\hat{\mu}_n$, a Mont Carlo estimate of μ . Granted, n cannot be very large given only six IID $U[0, 1]$ random numbers, how large can you n be?

- Computer Problem: Consider the following integral, which arises in the pricing of an Asian arithmetic mean call option:

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \max \left(\frac{1}{2} [S_1(\mathbf{z}) + S_2(\mathbf{z})] - 100, 0 \right) \frac{\exp(-\mathbf{z}^T \Sigma^{-1} \mathbf{z} / 2)}{\sqrt{4\pi^2 \det(\Sigma)}} d\mathbf{z}, \quad \Sigma = \begin{pmatrix} 1/2 & 1/2 \\ 1/2 & 1 \end{pmatrix},$$

$$S_1(\mathbf{z}) = 100 \exp(-0.0225 + 0.3z_1), \quad S_2(\mathbf{z}) = 100 \exp(-0.045 + 0.3z_2).$$

Approximate the integral using `meanMC_g` with an error tolerance of 0.02.

- Y has a triangular probability density function (PDF) of

$$\rho(y) = \begin{cases} \frac{y-10}{100}, & 10 \leq y \leq 20, \\ \frac{30-y}{100}, & 20 < y \leq 30, \\ 0, & \text{otherwise.} \end{cases}$$

Verify that ρ is a PDF. Use the inverse transform method to generate Y using the following six IID $U[0, 1]$ random numbers:

0.1135, 0.9745, 0.7287, 0.3515, 0.7076, 0.7996

5. Computer Problem: Consider a distribution with density function as

$$\rho(x) = \frac{1}{B(\alpha, \beta)} x^{\alpha-1} (1-x)^{\beta-1}, \quad 0 \leq x \leq 1,$$

where $B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$, $\Gamma(\cdot)$ is the gamma function. Derive an acceptance-rejection method for generating random variables with above pdf with $\alpha = 2, \beta = 2$ from $U[0, 1]$ random variables. Generate 1000 such random numbers.