

PROBABILITY & STATISTICS

Continuous R.V Pdf Cdf Chapter 4: Continuous Random Variables and Probability Distributions

Mean & Variance

LEARNING OBJECTIVES

Common Distri.
Uniform
Normal

1. Continuous random variable:

Normal approxi.

(a) Probability density function

- (b) Cumulative distribution function
- (c) Mean and Variance
- 2. Common distribution: uniform and normal, exponential
- 3. Normal approximation to the Binomial and Poisson.



CONTINUOUS RANDOM VARIABLE

Continuous R.V

Pdf Cdf

Mean & Variance


Common Distri.
Uniform
Normal


Normal approxi.

Summary

Definition

A continuous random variable is a random variable whose possible values includes in an interval of real numbers.





The time spent studying can be any number between 0 and 24.



Continuous R.V

Pdf

Cdf

Mean & Variance

Common Distri. Uniform Normal

Normal approxi.

Summary

Definition

The **probability density function (pdf)** of a continuous random variable *X* is a function such that

$$(1) \quad f(x) \ge 0 \qquad \forall \quad x$$

$$(2) \int_{-\infty}^{+\infty} f(x) dx = 1$$

(3)
$$P(a \le X \le b) = \int_{a}^{b} f(x) dx \quad \text{for an}$$

for any a and b.



Continuous R.V

Pdf

Cdf

Mean & Variance

Common Distri. Uniform Normal

Normal approxi.

Summary




Figure 4-2 Probability determined from the area under f(x).

Property

If X is a continuous random variable then for any x_1 and x_2 we have $P(x \le X \le x_1) - P(x \le X \le x_2)$

$$P(x_1 \le X \le x_2) = P(x_1 \le X < x_2)$$

$$= P(x_1 < X \le x_2)$$

$$= P(x_1 < X < x_2)$$



Continuous R.V

Pdf

Cdf

Mean & Variance

Common Distri. Uniform Normal

Normal approxi.

Summary

Example

Let the continuous random variable X denote the diameter of a hole drilled in a sheet metal component. The target diameter is 12.5 millimeters. Most random disturbances to the process result in larger diameters. Historical data show that the distribution of X can be modeled by a probability density function

$$f(x) = 20e^{-20(x-12,5)}, x \ge 12.5$$

- (a) If a part with a diameter larger than 12.60 millimeters is scrapped, what proportion of parts is scrapped?
- (b) What proportion of parts is between 12.5 and 12.6 millimeters?



Continuous R.V

Pdf

Cdf

Mean & Variance

Common Distri. Uniform Normal

Normal approxi.

Summary


Solution

(a)
$$P(X > 12.60) = \int_{12.6}^{+\infty} f(x) dx = 20 \int_{12.6}^{+\infty} e^{-20(x-12.5)} dx$$

= $-e^{-20(x-12.5)} \Big|_{12.6}^{+\infty} = 0.135$

(b)
$$P(12.5 < X < 12.6) = \int_{12.5}^{12.6} 20e^{-20(x-12.5)} dx$$

= $-e^{-20(x-12.5)} \Big|_{12.5}^{12.6} = 0.865$





CUMULATIVE DISTRIBUTION FUNCTION

Continuous R.V Pdf

Cdf

Mean & Variance

Common Distri. Uniform Normal

Normal approxi.

Summary

Definition

The **cumulative distribution function (cdf**) of a continuous random variable X is

$$F(x) := \int_{-\infty}^{x} f(t)dt$$

for $-\infty < x < +\infty$.

Let us return to the above example

$$f(x) = \begin{cases} 20e^{-20(x-12,5)} & x \ge 12.5\\ 0 & x < 12.5 \end{cases}$$

$$F(x) = \begin{cases} 1 - e^{-20(x - 12, 5)} & x \ge 12.5 \\ 0 & x < 12.5 \end{cases}$$



MEAN AND VARIANCE

Continuous R.V
Pdf
Cdf

Mean & Variance

Common Distri. Uniform Normal

Normal approxi.

Summary

Definition

Suppose X is a continuous random variable with probability density function f(x).

The **mean** or **expected value** of *X* is defined by

$$\mu = E(X) := \int_{-\infty}^{+\infty} x f(x) dx$$

The **variance** of *X* is defined by

$$\sigma^2 = V(X) := \int_{-\infty}^{+\infty} (x - \mu)^2 f(x) dx = \int_{-\infty}^{+\infty} x^2 f(x) dx - \mu^2$$

The **standard deviation** of *X* is $\sigma = \sqrt{\sigma^2}$.



MEAN AND VARIANCE

Continuous R.V Pdf Cdf

Mean & Variance

Common Distri. Uniform Normal

Normal approxi.

Summary

Example

Assume that X is a continuous random variable with the following probability density function

$$f(x) = \begin{cases} 20e^{-20(x-12.5)} & x \ge 12.5\\ 0 & x < 12.5 \end{cases}$$

Mean:
$$EX = \int_{12.5}^{+\infty} x f(x) dx = \int_{12.5}^{+\infty} x 20 e^{-20(x-12.5)} dx$$

Integration by parts can be used to show that

$$EX = \left(-xe^{-20(x-12.5)} - \frac{e^{-20(x-12.5)}}{20}\right)_{12.5}^{+\infty} = 12.55$$



MEAN AND VARIANCE

Continuous R.V Pdf Cdf Variance:

$$V(X) = \int_{12.5}^{+\infty} x^2 f(x) dx - (EX)^2 = 0.0025$$

Mean & Variance

Common Distri. Uniform Normal

Expected Value of a Function of a Continuous Random Variable

Normal approxi.

$$E h(X) = \int_{-\infty}^{+\infty} h(x) f(x) dx$$



CONTINUOUS UNIFORM RANDOM VARIABLE

Continuous R.V Pdf Cdf

Continuous uniform random variable over interval [a, b]

pdf:
$$f(x) = \begin{cases} \frac{1}{b-a}, & a \le x \le b \\ 0 & otherwise \end{cases}$$

Mean & Variance

Common Distri.

Uniform

Normal

Normal approxi.

Mean and Variance:

$$\mu = EX = \frac{a+b}{2}$$
 , $\sigma^2 = V(X) = \frac{(b-a)^2}{12}$

cdf:
$$F(x) = \begin{cases} 0 & x < a \\ \frac{x-a}{b-a} & a \le x < b \\ 1 & b \ge b \end{cases}$$



Continuous R.V Pdf Cdf

Mean & Variance

Common Distri. Uniform

Normal

Normal approxi.

Summary


Normal random variable $X \sim N(\mu, \sigma^2)$

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{\frac{-1}{2}(\frac{x-\mu}{\sigma})^2}, -\infty < x < +\infty$$

Mean and Variance:

$$E(X) = \mu$$

$$E(X) = \mu$$
 $V(X) = \sigma^2$





Continuous R.V Pdf Cdf


Mean & Variance

Common Distri.

<u>Uniform</u>

Normal

Normal approxi.





Continuous R.V Pdf Cdf

Mean & Variance

Common Distri.
Uniform

Normal

Normal approxi.

Summary

Standard Normal Random Variable $Z \sim N(0,1)$

$$f(x) = \frac{1}{\sqrt{2\pi}} e^{\frac{-x^2}{2}}$$
, $-\infty < x < +\infty$

$$\Phi(z) = \int_{-\infty}^{z} \frac{e^{\frac{-x^2}{2}}}{\sqrt{2\pi}} dx.$$

Standardizing

If X is a normal random variable $X \sim N(\mu, \sigma^2)$ then

$$Z = \frac{X - \mu}{\sigma}$$

is a standard normal random variable N(0, 1).



Continuous R.V
Pdf
Cdf

Mean & Variance


Common Distri.
Uniform

Normal

Normal approxi.

Summary

Standardizing





Continuous R.V

Pdf Cdf

Mean & Variance

Common Distri. Uniform

Normal

Normal approxi.

Summary

Finding Probabilities

Problem 1: x is given known, find P(X < x).

Problem 2: P(X < x) = p is given known, find x.

Using Excel

1. $Z \sim N(0,1)$

To find P(Z < z) when given z: =NORMSDIST(z)

To find z when P(Z < z) = p: =NORMSINV(p)

#Using R pnorm(z) qnorm(p)

2.
$$X \sim N(\mu, \sigma^2)$$

To find P(X < x) when given x: =NORMDIST(x, μ , σ ,1)

To find x when P(X < x) = p: =NORMINV(p, μ , σ).



Continuous R.V Pdf Cdf

Example

(a) Let $X \sim N(34, 144)$. Find P(X < 43) and P(24 < X < 37).

Mean & Variance

(b) Let $Z \sim N(0,1)$. Find the value of z to P(Z > z) = 0.95

Common Distri.
Uniform

Normal

Normal approxi.

(a)
$$P(X < 43) = NORMDIST(43, 34, 12,1) = 0.7734$$

$$P(24 < X < 37) = P(X < 37) - P(X < 24)$$

$$= 0.5987 - 0.2023 = 0.3964$$

(b)
$$P(Z < z) = 1 - P(Z > z) = 1 - 0.95 = 0.05$$

$$z = NORMSINV(0.05) = -1.65$$



Continuous R.V Pdf Cdf

Mean & Variance

Common Distri.
Uniform

Normal

Normal approxi.

Summary

Example

- (a) Let $X \sim N(34, 144)$. Find P(X < 43) and P(24 < X < 37).
- (b) Let $Z \sim N(0,1)$. Find the value of z to P(Z > z) = 0.95

```
> pnorm(43,34,sqrt(144))
[1] 0.7733726
> pnorm(37,34,12)-pnorm(24,34,12)
[1] 0.3963779
> qnorm(1-0.95)
[1] -1.644854
```



Continuous R.V
Pdf
Cdf

Mean & Variance

Common Distri.
Uniform

Normal

Normal approxi.

Summary

Finding Probabilities: Using Table II

 Table II Cumulative Standard Normal Distribution (continued)

Z	0.00	0.01	0.02	0.03	0.04	0.05
0.0	0.500000	0.50 <mark>3</mark> 989	0.507978	0.511967	0.515953	0.519939
0.1	0.539828	0.54 <mark>3</mark> 795	0.547758	0.551717	0.555760	0.559618
0.2	0.579260	0.583166	0.587064	0.590954	0.594835	0.598706
0.3 -	0.617911	0.621719	0.625516	0.62 <mark>9</mark> 300	0.633072	0.636831
0.4	0.655422	0.659097	0.662757	0.666402	0.670031	0.673645
0.5	0.691462	0.694974	0.698468	0.701944	0.705401	0.708840
0.6	0.725747	0.729069	0.732371	0.73 <mark>5</mark> 653	0.738914	0.742154
0.7	0.758036	0.761148	0.764238	0.767305	0.770350	0.773373
0.8	0.788145	0.791030	0.793892	0.796731	0.799546	0.802338
0.9	0.815940	0.818589	0.821214	-0.823815	0.826391	0.828944
1.0	0.841345	0.843752	0.846136	0.848495	0.850830	0.853141
1.1	0.864334	0.866500	0.868643	0.870762	0.872857	0.874928
1.2	0.884930	0.886860	0.888767	0.890651	0.892512	0.894350



Continuous R.V
Pdf
Cdf


Mean & Variance

Common Distri.
Uniform
Normal

Normal approxi.

Summary

Under certain conditions, the normal distribution can be used to approximate the binomial distribution and the Poisson distribution.





Continuous R.V Pdf Cdf Example

Let $X \sim B(16 \times 10^6, 10^{-5})$. Find the probability P(X > 150).

Mean & Variance

Solution

Common Distri. Uniform Normal

 $P(X > 150) = 1 - P(X \le 150)$

$$=1-\sum_{x=0}^{150}C_{16\times10^6}^x(10^{-5})^x(1-10^{-5})^{16\times10^6-x}$$

Normal approxi.

Summary

Clearly, this probability is difficult to compute.



Continuous R.V Pdf Cdf

Mean & Variance

Common Distri. Uniform Normal

Normal approxi.

Summary

Normal Approximation to the Binomial Distribution

If $X \sim B(n, p)$ then random variable

$$Z = \frac{X - np}{\sqrt{np(1-p)}}$$

is approximately a standard random variable N(0,1).

$$P(X \le x) \approx P(Z \le \frac{x + 0.5 - np}{\sqrt{np(1-p)}})$$

$$P(X \ge x) \approx P(Z \ge \frac{x - 0.5 - np}{\sqrt{np(1-p)}})$$

Remark: The approximation is good for np > 5 and n(1-p) > 5.



Continuous R.V Pdf Cdf

Mean & Variance

Common Distri. Uniform Normal

Normal approxi.

Summary

Let us return to the above example

$$P(X > 150) = P(X \ge 151) = P(Z > \frac{150.5 - 160}{\sqrt{160(1 - 10^{-5})}})$$
$$= P(Z > -0.75) = 1 - P(Z < -0.75) = 0.773$$

Here, we use the result $np = 16 \times 10^{6} \times 10^{-5} = 160$.



Continuous R.V
Pdf
Cdf

Mean & Variance

Common Distri.
Uniform
Normal

Normal approxi.

Summary

Normal Approximation to the Poisson Distribution

If X is a Poisson random variable $P(\lambda)$ then

$$Z = \frac{X - \lambda}{\sqrt{\lambda}}$$

is approximately a standard random variable N(0,1).

$$P(X \le x) = P(Z \le \frac{x - \lambda}{\sqrt{\lambda}}) = \Phi(\frac{x - \lambda}{\sqrt{\lambda}}).$$

The approximation is good for $\lambda > 5$.



Continuous R.V Pdf Cdf

Mean & Variance

Common Distri. Uniform Normal

Normal approxi.

Summary

Example

Assume that the number of asbestos particles in a squared meter of dust on a surface follows a Poisson distribution with a mean of 1000. If a squared meter of dust is analyzed, what is the probability that less than 950 particles are found?

Let X = the number of asbestos particles in a squared meter of dust on a surface, then $X \sim P(1000)$

$$P(X \le 950) \cong P(Z \le \frac{950 - 1000}{\sqrt{1000}}) = P(Z \le -1.58) = 0.057$$



Continuous R.V Pdf Cdf

Mean & Variance

Common Distri.
Uniform

Normal

Normal approxi.

Summary

The random variable X that equals the distance between successive counts of a Poisson process with mean $\lambda > 0$ is an **exponential random** variable with parameter λ .

$$f(x) = \lambda e^{-\lambda x}$$
 for $0 \le x < \infty$

$$\mu = E(X) = \frac{1}{\lambda}$$
 and $\sigma^2 = V(X) = \frac{1}{\lambda^2}$



Continuous R.V
Pdf
Cdf

Mean & Variance

Common Distri.
Uniform

Normal

Normal approxi.




Figure 4-22 Probability density function of exponential random variables for selected values of λ .



Continuous R.V Pdf Cdf

Mean & Variance

Common Distri.
Uniform

Normal

Normal approxi.

Summary

EXAMPLE. In a large corporate computer network, user log-ons to the system can be modeled as a Poisson process with a mean of λ = 25 log-ons per hour. What is the probability that there are no logons in an interval of 6 minutes?



Continuous R.V Pdf Cdf

Mean & Variance

Common Distri.
Uniform

Normal

Normal approxi.

Summary

EXAMPLE. In a large corporate computer network, user log-ons to the system can be modeled as a Poisson process with a mean of λ = 25 log-ons per hour. What is the probability that there are no logons in an interval of 6 minutes?

- X: time in hours from the start of the interval until the first log-on.
- X: exponential distribution
- 6 minutes=0.1 hour.

$$P(X > 0.1) = \int_{0.1}^{\infty} 25e^{-25x} dx = e^{-25(0.1)} = 0.082$$




Continuous R.V
Pdf
Cdf

Mean & Variance

Common Distri.
Uniform

Normal

Normal approxi.



$$P(X > 0.1) = \int_{0.1}^{\infty} 25e^{-25x} dx = e^{-25(0.1)} = 0.082$$





Continuous R.V Pdf Cdf

Mean & Variance

Common Distri. Uniform Normal

Normal approxi.

Summary

We have studied:

- 1. Continuous random variable:
- (a) Probability density function
- (b) Cumulative distribution function
- (c) Mean and Variance
- 2. Common distribution: uniform and normal, exponential
- 3. Normal approximation to the Binomial and Poisson.

Homework: Read slides of the next lecture.