

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Chapter 7: Point Estimation of Parameters

Learning objectives

1. Introduction
2. Sampling Distributions
3. General Concepts Of Point
4. Estimation Methods Of Point Estimation

Introduction

Sampling Distributions

General concepts of point estimation

Methods of point estimation

Method of moments

Method of maximum likelihood

Summary

Let X is a random variable with probability distribution $f(x)$, which is characterized by the unknown $\theta = (\theta_1, \theta_2, \dots, \theta_k)$

For example, $X \sim N(\mu, \sigma^2)$ then $\theta = (\mu, \sigma^2)$.

How to “determine” the values of θ ?

Introduction

Sampling Distributions

General concepts of point estimation

Methods of point estimation


Method of moments

Method of maximum likelihood

Summary

19/05/2021

Example



Introduction

Sampling Distributions

General concepts of point estimation

Methods of point estimation

Method of moments Method of maximum likelihood

Summary

In engineering, we often need to estimate

1. The mean μ of a single population.
2. The variance σ^2 (or standard deviation) of a single population.
3. The proportion p of items in a population that belong to a class of interest.
4. The difference in means of two populations, $\mu_1 - \mu_2$.
5. The difference in two population proportions, $p_1 - p_2$.

Introduction

Sampling Distributions

General concepts of point estimation

Methods of point estimation

Method of moments

Method of maximum likelihood

Summary

The important results on point estimation

1. For μ , the estimate is the $\hat{\mu} = \bar{x}$, **sample mean.**
2. For σ^2 , the estimate is $\hat{\sigma}^2 = s^2$, **the sample variance.**
3. For p , the estimate is $\hat{p} = x/n$, **the sample proportion.**
4. For $\mu_1 - \mu_2$, the estimate is $\hat{\mu}_1 - \hat{\mu}_2 = \bar{x}_1 - \bar{x}_2$
5. For $p_1 - p_2$, the estimate is $\hat{p}_1 - \hat{p}_2$

Introduction

Sampling Distributions

General concepts of point estimation

Methods of point estimation

Method of moments

Method of maximum likelihood

Summary

Definition

Random Sample

The random variables X_1, \dots, X_n are called a **random sample** of size n if

- The X_i 's are independent
- Every X_i has the same probability distribution

Definition

Statistic

- A **statistic** $\hat{\Theta}$ is any function of the observations X_1, \dots, X_n :

$$\hat{\Theta} = h(X_1, \dots, X_n)$$

- The probability distribution of a statistic is called a **sampling distribution**.

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Example

Two important statistic

- Sample mean \bar{X}

$$\bar{X} = \frac{X_1 + \dots + X_n}{n}$$

- Sample variance S^2

$$S^2 = \frac{(X_1 - \bar{X})^2 + \dots + (X_n - \bar{X})^2}{n-1}$$

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Theorem

In above example, if (X_1, \dots, X_n) is a random sample of size n take from a normal distribution $N(\mu, \sigma^2)$ then

- \bar{X} has a normal distribution $N(\mu, \sigma^2/n)$

- $\frac{(n-1)S^2}{\sigma^2}$ has a chi-square distribution with $n-1$ degrees of freedom (see pages 273-274).

Introduction

Sampling
DistributionsGeneral concepts
of point
estimationMethods of point
estimationMethod of
momentsMethod of
maximum
likelihood

Summary

EXAMPLE 7-1 Resistors

An electronics company manufactures resistors that have a mean resistance of 100 ohms and a standard deviation of 10 ohms. The distribution of resistance is normal. Find the probability that a random sample of $n = 25$ resistors will have an average resistance less than 95 ohms.

Note that the sampling distribution of \bar{X} is normal, with mean $\mu_{\bar{X}} = 100$ ohms and a standard deviation of

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}} = \frac{10}{\sqrt{25}} = 2$$

Introduction

Sampling Distributions

General concepts of point estimation

Methods of point estimation

- Method of moments
- Method of maximum likelihood

Summary

$$z = \frac{95 - 100}{2} = -2.5$$

$$\begin{aligned} P(\bar{X} < 95) &= P(Z < -2.5) \\ &= 0.0062 \end{aligned}$$




Figure 7-2 Probability for Example 7-1.

```
> #Example 7.1
> #In R, you can compute easily
> pnorm(95,100,2)
[1] 0.006209665
```

Introduction

Sampling Distributions

General concepts of point estimation

Methods of point estimation

Method of moments

Method of maximum likelihood

Summary

Theorem

Central Limit Theorem

- If we are sampling from a population that has an unknown probability distribution, **the sampling distribution of the \bar{X} sample mean is approximately normal if the sample size is large.**
- This is one of the most useful theorems in statistics, called the **central limit theorem.**

Example: Distributions of sample means when rolling 2,3,10,...dices

Use R to simulate the rolling a dices 2,3,5,10 times and study the distributions of sample means

- **Note that rolling a fair dice has a uniform distribution with the mean 3.5 and variance 35/12**
- **Study \bar{X} sampling distributions of sample**

$$\mu = E(X) = (a + b)/2 \quad \sigma^2 = \frac{(b - a + 1)^2 - 1}{12}$$

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Example: Distributions of sample means when rolling 2,3,10,...dices

Simulation with R:

1. Define a command to roll a dice **n times**
2. Make **random samples of rolling a dice n times and compute sample means.**
3. Showing the **histogram of the sample means**
4. Compute the mean and variance of the sample mean variable.

Introduction

Sampling
Distributions

General concepts
of point
estimation


Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Figure 7-1
Distributions of
average scores
from throwing
dice.



Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

```
RollDice=function(n) sample(1:6,n,replace=T)
means = vars = numeric(1000)
for (i in 1:1000) {
  samp = RollDice(2)
  #print(samp)
  means[i] = mean(samp)
  vars[i] = var(samp)
}
means
vars
hist(means)
hist(vars)
```

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Theorem

Central Limit Theorem

Let (X_1, \dots, X_n) is a random sample of size n take from a population with mean μ and finite variance σ^2 , and if \bar{X} is the sample mean, the limiting form of the distribution of

$$Z = \frac{\bar{X} - \mu}{\sigma / \sqrt{n}}$$

as $n \rightarrow \infty$, is the standard normal distribution.

Remark: The normal approximation for \bar{X} depends on the sample size n .

Introduction

Sampling Distributions

General concepts of point estimation

Methods of point estimation

Method of moments

Method of maximum likelihood

Summary

EXAMPLE 7-2 Central Limit Theorem

Suppose that a random variable X has a continuous uniform distribution

$$f(x) = \begin{cases} 1/2, & 4 \leq x \leq 6 \\ 0, & \text{otherwise} \end{cases}$$

Find the distribution of the sample mean of a random sample of size $n = 40$.

The mean and variance of X are $\mu = 5$ and $\sigma^2 = (6 - 4)^2/12 = 1/3$. The central limit theorem indicates that the distribution of \bar{X} is approximately normal with mean $\mu_{\bar{X}} = 5$ and variance $\sigma_{\bar{X}}^2 = \sigma^2/n = 1/[3(40)] = 1/120$. The




Figure 7-3 The distributions of X and \bar{X} for Example 7-2.

Approximate Sampling Distribution of a Difference in Sample Means

Introduction

Sampling Distributions

General concepts of point estimation

Methods of point estimation

Method of moments

Method of maximum likelihood

Summary

If we have two independent populations with means μ_1 and μ_2 and variances σ_1^2 and σ_2^2 , and if \bar{X}_1 and \bar{X}_2 are the sample means of two independent random samples of sizes n_1 and n_2 from these populations, then the sampling distribution of

$$Z = \frac{\bar{X}_1 - \bar{X}_2 - (\mu_1 - \mu_2)}{\sqrt{\sigma_1^2/n_1 + \sigma_2^2/n_2}} \quad (7-4)$$

is approximately standard normal, if the conditions of the central limit theorem apply. If the two populations are normal, the sampling distribution of Z is exactly standard normal.

Approximate Sampling Distribution of a Difference in Sample Means

Introduction

Sampling Distributions

General concept of point estimation

Methods of point estimation

Method of moments

Method of maximum likelihood

Summary

EXAMPLE 7-3 Aircraft Engine Life

The effective life of a component used in a jet-turbine aircraft engine is a random variable with mean 5000 hours and standard deviation 40 hours. The distribution of effective life is fairly close to a normal distribution. The engine manufacturer

introduces an improvement into the manufacturing process for this component that increases the mean life to 5050 hours and decreases the standard deviation to 30 hours. Suppose that a random sample of $n_1 = 16$ components is selected from the “old” process and a random sample of $n_2 = 25$ components is selected from the “improved” process. What is the probability that the difference in the two sample means $\bar{X}_2 - \bar{X}_1$ is at least 25 hours? Assume that the old and improved processes can be regarded as independent populations.

Approximate Sampling Distribution of a Difference in Sample Means

Introduction

Sampling Distributions

General concepts of point estimation


Methods of point estimation

Method of moments

Method of maximum likelihood

Summary

Figure 7-4 The sampling distribution of $\bar{X}_2 - \bar{X}_1$ in Example 7-3.



To solve this problem, we first note that the distribution of \bar{X}_1 is normal with mean $\mu_1 = 5000$ hours and standard deviation $\sigma_1/\sqrt{n_1} = 40/\sqrt{16} = 10$ hours, and the distribution of \bar{X}_2 is normal with mean $\mu_2 = 5050$ hours and standard deviation $\sigma_2/\sqrt{n_2} = 30/\sqrt{25} = 6$ hours. Now the distribution of $\bar{X}_2 - \bar{X}_1$ is normal with mean $\mu_2 - \mu_1 = 5050 - 5000 = 50$ hours and variance $\sigma_2^2/n_2 + \sigma_1^2/n_1 = (6)^2 + (10)^2 = 136$ hours². This sampling distribution is shown in Fig. 7-4. The probability that $\bar{X}_2 - \bar{X}_1 \geq 25$ is the shaded portion of the normal distribution in this figure.

Approximate Sampling Distribution of a Difference in Sample Means

Introduction

Sampling Distributions

General concepts of point estimation


Methods of point estimation

Method of moments

Method of maximum likelihood

Summary

Figure 7-4 The sampling distribution of $\bar{X}_2 - \bar{X}_1$ in Example 7-3.



Corresponding to the value $\bar{x}_2 - \bar{x}_1 = 25$ in Fig. 7-4, we find that

$$z = \frac{25 - 50}{\sqrt{136}} = -2.14$$

and consequently,

$$\begin{aligned} P(\bar{X}_2 - \bar{X}_1 \geq 25) &= P(Z \geq -2.14) \\ &= 0.9838 \end{aligned}$$

Therefore, there is a high probability (0.9838) that the difference in sample means between the new and the old process will be at least 25 hours if the sample sizes are $n_1 = 16$ and $n_2 = 25$.

Introduction

Sampling Distributions

General concepts of point estimation

Methods of point estimation

Method of moments

Method of maximum likelihood

Summary

Definition

Point Estimate

- A **point estimate** of some population parameter θ is a single numerical value $\hat{\theta}$ of a statistic $\hat{\Theta}$.
- The statistic $\hat{\Theta}$ is called the **point estimator**.

Two steps to find point estimation:

Step 1. Determine $\hat{\Theta}$ by using the theoretical results.

Steps 2. Calculate $\hat{\theta}$ from the experimental data.

Introduction

Sampling Distributions

General concepts of point estimation

Methods of point estimation

Method of moments

Method of maximum likelihood

Summary

Example

Market researchers use the number of sentences per advertisement as a measure of readability for magazine advertisements. The following represents a random sample of the number of sentences found in 15 advertisements. Find a point estimate of the population mean, μ .

9 20 18 16 9 9 11 13 22 16 5 18 6 6 5

$$\text{Step 1. } \hat{\Theta} = \frac{X_1 + \dots + X_{15}}{15}$$

Step 2. A point estimate for μ is

$$\hat{\mu} = \frac{9 + 20 + \dots + 6 + 5}{15} = \frac{183}{15} = 12.2$$

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Definition

Unbiased Estimator

The point estimator $\hat{\Theta}$ is an **unbiased estimator** for the parameter θ if

$$E(\hat{\Theta}) = \theta$$

If the estimator is not unbiased, then the difference

$$E(\hat{\Theta} - \theta)$$

is called the **bias** of the estimator $\hat{\Theta}$.

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Example

Suppose that X is a random variable with mean μ and variance σ^2 . Let X_1, \dots, X_n be a random sample of size n from the population represented by X . Show that the sample mean \bar{X} and sample variance S^2 are unbiased estimators of μ and σ^2 , respectively.

For the sample mean.

$$E(\bar{X}) = E \frac{X_1 + \dots + X_n}{n} = \frac{EX_1 + \dots + EX_n}{n} = \mu$$

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Now consider the **sample variance**. We have

$$\begin{aligned}
 E(S^2) &= E \left[\frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1} \right] = \frac{1}{n-1} E \sum_{i=1}^n (X_i - \bar{X})^2 \\
 &= \frac{1}{n-1} E \sum_{i=1}^n (X_i^2 + \bar{X}^2 - 2\bar{X}X_i) = \frac{1}{n-1} E \left(\sum_{i=1}^n X_i^2 - n\bar{X}^2 \right) \\
 &= \frac{1}{n-1} \left[\sum_{i=1}^n E(X_i^2) - nE(\bar{X}^2) \right]
 \end{aligned}$$

Hence $E(S^2) = \sigma^2$

MINIMUM VARIANCE UNBIASED ESTIMATOR

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Definition


Minimum Variance Unbiased Estimator

If we consider all unbiased estimators of θ , the one with the smallest variance is called the **minimum variance unbiased estimator** (MVUE).

Figure 7-5

The sampling
distributions of
two unbiased
estimators

$\hat{\Theta}_1$ and $\hat{\Theta}_2$.



Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Theorem

Let (X_1, \dots, X_n) is a random sample of size n take from a normal distribution $N(\mu, \sigma^2)$, the sample mean \bar{X} is the MVUE for μ .

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Definition Minimum Variance Unbiased Estimator

The **standard error** of an estimator $\hat{\Theta}$ is its standard deviation, given by

$$\sigma_{\hat{\Theta}} = \sqrt{V(\hat{\Theta})}$$

If the standard error involves unknown parameters that can be estimated, substitution of those values into $\sigma_{\hat{\Theta}}$ produces an **estimated standard error**, denoted by $\hat{\sigma}_{\hat{\Theta}}$

Remark

If the random sample of size n take from a normal distribution $N(\mu, \sigma^2)$ then \bar{X} has a normal distribution $N(\mu, \sigma^2/n)$.

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments
Method of
maximum
likelihood

Summary

Standard error of \bar{X} is

$$\sigma_{\hat{X}} = \frac{\sigma}{\sqrt{n}}$$

If we did not know σ then the estimated standard error of \bar{X} would be

$$\hat{\sigma}_{\hat{X}} = \frac{s}{\sqrt{n}}$$

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Exercise

A new method of measuring the thermal conductivity of Armco iron: Using a temperature of 100°F and a power input of 550 watts, the following 10 measurements of thermal conductivity (in $\text{Btu/hr-ft-}^{\circ}\text{F}$) were obtained:

41.60, 41.48, 42.34, 41.95, 41.86,
42.18, 41.72, 42.26, 41.81, 42.04

a, Find a point estimate of the mean thermal conductivity at 100°F and 550 watts.

b, Find the estimated standard error of \bar{X}

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Definition

The k th moments

Let (X_1, \dots, X_n) be a random sample from random variable X with the probability distribution $f(x)$

- The k th moment of X are denoted by $M_k := E(X^k)$
- The corresponding k th **sample moment** is

$$\bar{X}^k = \frac{X_1^k + \dots + X_n^k}{n}$$

Introduction

Sampling

Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Definition

The k th moments

Let (X_1, \dots, X_n) be a random sample from either a probability mass function or probability density function with m unknown parameters $(\theta_1, \theta_2, \dots, \theta_m)$. The moment estimators $\hat{\Theta}_1, \dots, \hat{\Theta}_m$ are found the following system

$$\begin{cases} M_1 = \bar{X}^1 \\ \dots\dots\dots \dots \\ M_m = \bar{X}^m \end{cases}$$

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Example

Suppose that X_1, \dots, X_n be a random sample of size n from a normal distribution $N(\mu, \sigma^2)$. Find the moment estimators of μ and σ^2 .

We have to solve following system

$$\begin{cases} M_1 = \bar{X} \\ M_2 = \bar{X}^2 \end{cases} \quad \text{or} \quad \begin{cases} EX = \frac{X_1 + \dots + X_n}{n} \\ EX^2 = \frac{X_1^2 + \dots + X_n^2}{n} \end{cases}$$

Here $E(X) = \mu$ and $E(X^2) = V(X) + (EX)^2 = \sigma^2 + \mu^2$

Introduction

Sampling

Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Hence, by solving the above system we have

$$\mu = \bar{X}, \quad \sigma^2 = \frac{\sum_{i=1}^n X_i^2 - n(\bar{X})^2}{n} = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n}$$

Finally, the moment estimators are

$$\hat{\mu} = \bar{X},$$

Unbiased
estimator

$$\hat{\sigma}^2 = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n}$$

Biased
estimator

Introduction

Sampling

Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Definition

Maximum likelihood Estimator

Suppose that X is a random variable with probability distribution $f(x, \theta)$, where θ is a single unknown parameter. Let x_1, x_2, \dots, x_n be the observed values in a random sample of size n . Then the **likelihood function** of the sample is

$$L(\theta) = f(x_1, \theta) f(x_2, \theta) \dots f(x_n, \theta)$$

The maximum likelihood estimator (MLE) of θ is the value of θ that maximizes the $L(\theta)$.

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Example

Let X is a Bernoulli random variable. The probability mass function is

$$f(x, p) = \begin{cases} p^x (1-p)^{1-x}, & x = 0, 1 \\ 0 & \text{otherwise} \end{cases}$$

Find the maximum likelihood estimator of p .

The likelihood function of a random sample of size n is

$$\begin{aligned} L(p) &= p^{x_1} (1-p)^{1-x_1} p^{x_2} (1-p)^{1-x_2} \cdots p^{x_n} (1-p)^{1-x_n} \\ &= \prod_{i=1}^n p^{x_i} (1-p)^{1-x_i} = p^{\sum_{i=1}^n x_i} (1-p)^{n - \sum_{i=1}^n x_i} \end{aligned}$$

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

$$\ln L(p) = \left(\sum_{i=1}^n x_i \right) \ln p + \left(n - \sum_{i=1}^n x_i \right) \ln(1 - p)$$

$$\frac{d \ln L(p)}{dp} = \frac{\sum_{i=1}^n x_i}{p} - \frac{\left(n - \sum_{i=1}^n x_i \right)}{1 - p}$$

The maximum likelihood estimator of p is

$$\hat{P} = \frac{X_1 + \dots + X_n}{n}$$

Is this a unbiased estimator?

Introduction

Sampling

Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Exercise

Let X is a random variable, which has probability mass function is

$$f(x) = \begin{cases} p & , \quad x = -1 \\ 1 - p & , \quad x = 1 \\ 0 & , \quad otherwise \end{cases}$$

Find the maximum likelihood estimator of p .

Solution:

$$f(x) = \begin{cases} p^{\frac{1-x}{2}} (1-p)^{\frac{1+x}{2}} & , \quad x = -1, 1 \\ 0 & , \quad otherwise \end{cases}$$

$$\hat{p} = \frac{1}{2} - \frac{X_1 + \dots + X_n}{2n}$$

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Invariance Property

Let $\hat{\theta}_1, \dots, \hat{\theta}_k$ be the maximum likelihood estimators of the parameters $\theta_1, \dots, \theta_k$. Then the maximum likelihood estimator of any function $h(\theta_1, \dots, \theta_k)$ of these parameters is the same function $h(\hat{\theta}_1, \dots, \hat{\theta}_k)$ of the estimators $\hat{\theta}_1, \dots, \hat{\theta}_k$.

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

Exercise

Let X be normally distributed $N(\mu, \sigma^2)$, where both μ and σ^2 are unknown.

- (a) Find the maximum likelihood estimator of μ and σ^2 .
- (b) Find the maximum likelihood estimator of $\theta = \mu^2\sigma^2$.

Solution:

$$\hat{\mu} = \bar{X}, \quad \hat{\sigma}^2 = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n}$$

$$\hat{\theta} = (\bar{X})^2 \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n}$$

Introduction

Sampling
Distributions

General concepts
of point
estimation

Methods of point
estimation

Method of
moments

Method of
maximum
likelihood

Summary

We have studied:

1. Sampling distributions
 2. Central Limit Theorem
 3. General concepts of point estimation: Unbiased Estimator
 4. Methods of point estimation
 - Method of Moments
 - Method of Maximum Likelihood
- Homework: Read slides of the next lecture.