

The Office DVD Problem

Dave Fetterman

Obviously Unemployed

7/10/22

Screensavers have captivated [this] man since the 1990s. If watched long enough, what will the spirits of the machine tell us?

Specifically, the question of whether a bouncing rectangle will slide *exactly* into the corner of the screen, for a satisfying, perfectly diametric rebound, was even addressed on *The Office* (link).

However, though these characters reportedly watched this sleep-mode drama play out for years until payoff, we ask - under what conditions will the rectangle *definitely* perfectly bounce into the screen's corner?

0.1 Statement

Suppose we have a screen of length l , height h , containing an axis-aligned rectangle of length j and height k centered at point (x, y) .

Suppose this rectangle is launched at direction $\langle 1, m \rangle$ ¹ and “bounces” according to billiard

¹Think of this as slope m




Figure 1: The Office DVD problem's most generic setup




Figure 2: Success for $m = \frac{2}{3}, j, k = 0, h, l = 1$ (not to scale)

rules ².

Given $l, h, j, k, m \in \mathbb{R}$, can we tell whether the rectangle ever bounce perfectly into a corner?

We can approach this problem from the simplest version to the most complex.

0.2 Problem 1

Suppose $j = k = 0$ and $x = y = 0$. In other words, suppose we have a *point* starting at the bottom left corner (origin). Under what conditions (i.e. choice of m) does this bounce into a corner?

0.3 Problem 2

Suppose $j, k > 0, x = \frac{j}{2}, y = \frac{k}{2}$. In other words, suppose we have a rectangle starting at the bottom left corner. Under what conditions does this bounce into a corner?

0.4 Problem 3

Suppose we have maximally open (reasonable) conditions, with $x \in [\frac{j}{2}, l - \frac{j}{2}], y \in [\frac{k}{2}, h - \frac{k}{2}]$ (that is, a $j \times k$ rectangle fitting entirely in the screen). Under what conditions does this bounce into a corner?

²Glancing off a horizontal boundary, our trajectory goes from $\langle 1, m \rangle$ to $\langle 1, -m \rangle$, with $\langle \pm 1, m \rangle$ to $\langle \mp 1, m \rangle$ for a horizontal one