

BADG 2: A Perfect Game

Dave Fetterman

August 25, 2025

Abstract

An earlier paper (<http://fettermania.com/math/bdice.pdf>) focused on a near-optimal strategy for maximizing expected value in gameplay for *Bitches: A Dice Game*. This paper focuses on a different question: what is the probability of playing a perfect game? Using standard recursive calculation, we calculate the probabilities with the out-of-box game setup. Then we venture into a related question: What's the expected win probability with n standard dice of side count s as $n \rightarrow \infty$? We give poor bounds, prove convergence, and find surprising results along the way.

1 Introduction and the Commercial Game

Outside of ideological patron Larry Waldman, the publication of my original b. a. d. g. paper (<http://fettermania.com/math/bdice.pdf>) received little fanfare from humans. However, the robots somehow picked up on this and brought it to the attention of Samuel Cox, creator of the game. After dispensing mild praise about the paper and strong confusion about my interest in it, Mr. Cox suggested another problem.

Instead of “how does one play the game with high/maximum expected value?” (addressed in the first paper), Mr. Cox wondered how often a *perfect game* (one in which the best possible score is achieved at the end) occurs if played optimally. Though optimal play looks obvious (pick up all your good rolls), Mr. Cox has also debated with his class: “is it always best to pick up all your maximum-roll dice?”

I took this on and a day or so later, the solution to this particular question surfaced. As usual, twinkly subproblems appeared along the way, and I left to find a solution to a related problem: *does the probability of a perfect game converge as the number of dice increase to infinity?*

We'll briefly review the game's rules and the solution method for the commercially offered game before diving in to a discovered second quest: asymptotic behavior of infinite dice. We'll note surprising or interesting findings in ***bold-italic***.

1.1 The Game

Though the rules to the multi-player game are detailed in the original paper (<http://fettermania.com/math/bdi>) we will reformulate this for a solitaire-style perfect game, which follows this pattern:

1. Begin with 12 standard dice of six sides, along with one eight-, one ten-, and one twelve-sided die.
2. On each turn, roll all remaining dice in play.
3. If there are no dice with a maximum side showing (e.g. a six-sided die showing six pips), you have lost (“busted”).
4. Set aside some number of dice which have achieved their maximum roll.
5. Continue until no dice remain (you have a perfect game) or you have busted out.

The question “*how likely am I to get a perfect score?*” therefore relies on the *strategy* one employs in picking up dice.

1.2 Mistake 1: The Strategy Assumption

Proposition 1.1 (Smoke ’em if you got ’em). *The best strategy to achieve a perfect game is to always pick up as many dice with maximum face value as the turn allows.*

This proposition is:

1. Simple
2. Obvious
3. The strategy any reasonable person would employ
4. Assumed for the calculations in the next section
5. ***Mathematically false***

In the next section, we assume this simple, deterministic strategy in calculating the expected value of any state reachable from the standard b. a. d. g. starting configuration and show where it goes wrong.

1.3 Solution Method for the Commercial Game

The following definitions will help us write a program to get our solution for the commercial game:

- The “state” $\vec{s} := [a, b, c, d]$ is how many dice of size 6, 8, 10, and 12 remain, respectively. We start at $\vec{S} = [12, 1, 1, 1]$. A perfect game ends at $\vec{s} = \vec{0}$. A failure to roll any max roll ends the game in an unlisted ‘losing’ state $\emptyset$.

- The “probability vector” $\vec{p} := [1/6, 1/8, 1/10, 1/12]$ is the chance of getting a max roll from each of the dice sizes, again here assumed to be 6, 8, 10, 12.
- With a win defined as value 1 ($F(\vec{0}) = F([0, 0, 0, 0]) = 1$) and a loss as value 0 ($F(\emptyset) = 0$), the “expected game value” of any other state $\vec{s}$ is denoted as $F(\vec{s})$.
- Said another way, $F(\vec{s})$ is the probability of ending up in winning state $\vec{0}$ from state $\vec{s}$.
- We call any state $\vec{t} := [t_1, t_2, \dots, t_z]$ *reachable* from state $\vec{s} := [s_1, s_2, \dots, s_z]$ if $t_i \leq s_i$ for all $1 \leq i \leq z$, but $\vec{s} \neq \vec{t}$. More intuitively, if you can start at state $\vec{s}$ and get to state $\vec{t}$ in one roll (and corresponding pickup) of your dice, then $\vec{t}$ is reachable from $\vec{s}$. This is just any non-bust state “smaller than” where you are now.

1.4 Probability Lemmas

First, let’s limit ourselves to a game (or state) with only a single flavor of dice, say d -sided, collapsing state vector $\vec{s}$ to a single integer r (r dice of size d left). Refer to Fig. 1 for a visualization of the states starting at r dice left of size d , with $p := 1/d, q := 1 - p$.




Figure 1: Transition graph of the dice game showing expected values $F(r)$ and transition probabilities.

Lemma 1.2 (EV of r d -sided dice). *The chance of a completing perfect game from a state of r standard dice of d sides each, with definitions $p := \frac{1}{d}, P_{single}(r, k, p) := \binom{r}{k} p^k (1-p)^{r-k}$, and $F(0) = 1$ is defined by the recurrence $F(r) = \sum_{k=1}^r P_{single}(r, k, p) F(r-k)$.*

Proof: This is a clear application of the binomial theorem. With the Smoke 'Em if you Got 'Em Strategy, the chance of transitioning to the state with $r-k$ remaining dice from that with r is the chance of rolling exactly k max rolls and $r-k$ non-max rolls among your d -sided dice, or $P_{single}(r, k, p)$. The expected value of being in this new state is defined as $F(r-k)$. The value of this state transition possibility is $P_{single}(r, k, p) F(r-k)$. The total expected value of being in state r is therefore the sum of the values of all possible state transitions. Note that $k=0$ implies a roll with no maxes, and thus a loss (value 0), and

$F(0) = 1$ since no dice remaining (outside of a loss case) means a victory, or unit value of expectation.

Extending this to dice of z different sizes is clear:

Lemma 1.3 (State Transition lemma). $P_{transition}(\vec{s}, \vec{t}, \vec{p})$, the chance of transitioning from state $\vec{s} = [s_1, s_2, \dots, s_z]$ to state $\vec{t} = [t_1, t_2, \dots, t_z]$ with probability vector $\vec{p} = [p_1, p_2, \dots, p_z]$ is equal to $\prod_{i=1}^z P_{single}(\vec{s}_i, \vec{t}_i, p_i)$.

Proof: Divide the turn's roll into z independent subrolls, each rolling all s_i dice, $1 \leq i \leq z$, each with p_i chance of hitting a max. The outcomes of these z subrolls are all mutually independent, so $P_{transition}$ can be expressed as the product of P_{single} quantities.

1.5 Simple recursive algorithm

With these easy proofs and definitions in hand, the algorithm follows readily - build up each state from $F(\vec{0})$ to $F(\vec{S})$ by applying the State Transition Lemma and previously computed values of F in Algorithm 1. Notes:

- Steps 1 - 5 are definitions from the lemmas above.
- Step 6 creates a set of all states reachable from state $\vec{s}$, including $\vec{s}$ itself.
- Thinking of the states as a directed graph where a state points to all the states reachable from it, Step 7 creates an ordered list of states $s_1, s_2, s_3, \dots$, like $[0, 0, 0, 0], [0, 0, 1, 0], [0, 1, 0, 0], [0, 1, 1, 0]$, where, when s_i is encountered in the list, all states reachable from s_i occur before s_i in the list. There are many such possible orders. This is required for step 9's $F(\vec{t})$ values to be defined.
- Step 9 uses Lemma 1.3 to calculate $F(\vec{s})$ from the values of previously calculated substates.

1.6 Results for Commercial Game

Using the Smoke 'Em if you Got 'Em strategy, the chance of winning¹ $F(\vec{s})$ for every state $\vec{s}$, with $\vec{s}$ ranging from $\vec{0}$ to $\vec{S} = [12, 1, 1, 1]$ is listed in Figures 2 and 3. These are computed with the simple python program at <https://github.com/fettermania/mathnotes/blob/main/bdice2/bdice2.py>.

Note that these are *four-dimensional* results, with each dimension i corresponding to the count of dice s_i in each state. Since the number of 8-, 10-, and 12-sided dice is either zero or one, we've collapsed these into four 12 by 2 tables, where d_k notes the number of dice of size k .

Our main *initial* result is in the bottom-right corner of Fig. 3.

¹Alternatively, expected value of being in state $F(\vec{s})$, with $F(\vec{0}) = 1, F(\emptyset) = 0$

Algorithm 1 Dynamic program for $F(\vec{S})$

```
1:  $\vec{S} \leftarrow [12, 1, 1, 1]$ 
2:  $\vec{p} \leftarrow [\frac{1}{6}, \frac{1}{8}, \frac{1}{10}, \frac{1}{12}]$ 
3:  $P_{\text{single}}(r, k, p) := \binom{r}{k} p^k (1-p)^{r-k}$ 
4:  $P_{\text{transition}}(\vec{s}_1, \vec{s}_2, \vec{p}) := \prod_{i=0}^{\dim(\vec{s})} P_{\text{single}}(\vec{s}_{1i}, \vec{s}_{2i}, \vec{p}_i)$ 
5:  $F(\vec{0}) \leftarrow 1$ 
6:  $\text{substates}(\vec{s}) := \prod_{i=0}^{\dim(s)} [0, s_i]$ 
7:  $\text{states} \leftarrow \text{topological\_sort}(\text{substates}(\vec{S}) \setminus \{[0, 0, 0, 0]\})$ 
8: for  $\vec{s} \in \text{states}$  do
9:    $F(\vec{s}) \leftarrow \sum_{\vec{t} \in (\text{substates}(\vec{s}) \setminus \{\vec{s}\})} P_{\text{transition}}(\vec{s}, \vec{t}) \cdot F(\vec{t})$ 
10: end for
```

1.7 Observations

Observations from Figs. 2 and 3:

- Most of the time, starting with some state $\vec{s}$ and picking up one or more dice will increase the expected value of winning.
- However, we do note some non-monotonicity: there are two surprising “critical points” at $[12, 0, 1, 1]$ and $[9, 1, 1, 1]$, where instead of the EV decreasing as d_6 increases, the EV increases instead. This means that, say, rolling two max-sixes and no other max dice in state $[10, 1, 1, 1]$ suggests the best strategy is to **leave one max die on the table** so as to end up in slightly higher EV state $[9, 1, 1, 1]$ over $[8, 1, 1, 1]$. We will revisit these in the next section.
- At all states reachable from $[12, 0, 1, 1]$, the best strategy is confirmed: pick up as many max-rolled dice as possible. Deviating from this strategy produces an F -value smaller than those in the table. *Note:* The “bend” at $[11, 0, 1, 1]$ does not affect this; you have no alternative from $[12, 0, 1, 1]$ than to pick up if $[11, 0, 1, 1]$ is your new roll state.
- At all states reachable from $[10, 1, 1, 1]$, the simple strategy remains best.
- We can confirm that holding d_6 steady, it’s always better to have zero than one of each of the 8, 10, and 12-sided dice.

What The Creator Asked For: The chance of getting a perfect run the commercial game using the Smoke ’Em strategy is about 0.004413, or a little more than 4.4%.

$d_6 \backslash d_8$	0	1	$d_6 \backslash d_8$	0	1
0	1	0.125	0	0.1	0.03469
1	0.1667	0.05642	1	0.04556	0.02046
2	0.07407	0.03244	2	0.02644	0.01425
3	0.04192	0.02206	3	0.01815	0.01109
4	0.02809	0.01679	4	0.01392	0.009299
5	0.02107	0.01378	5	0.01152	0.008218
6	0.01708	0.01195	6	0.01006	0.007537
7	0.01463	0.01076	7	0.009124	0.0071
8	0.01303	0.009966	8	0.008506	0.006818
9	0.01195	0.009423	9	0.00809	0.006638
10	0.01119	0.009044	10	0.007809	0.006529
11	0.01065	0.008779	11	0.007618	0.006468
12	0.01025	0.008591	12	0.007491	0.006441

Figure 2: Expected values for $(d_{10}, d_{12}) = (0, 0)$ (left) and $(1, 0)$ (right).

1.8 Re-examining The optimal strategy

From examining the table it becomes clear that there are expected values for states $\vec{s}, \vec{t}$ such that $\vec{t}$ is reachable from $\vec{s}$ yet $F(\vec{t}) < F(\vec{s})$.

This means that for state, say, $\vec{s} + e_1$, that even when rolling enough maxes to get to $\vec{t}$, picking up fewer dice than the Smoke 'em if you Got 'em strategy may be best.

This means we need to refine our formula to choose the best selectable EV state from state $\vec{s}$, not just the furthest from $\vec{s}$. With this definition in hand:

$$selectable_states(\vec{s}, \vec{t}) := (substates(\vec{s}) - \vec{s}) - substates(\vec{t}) + \vec{t}$$

Update Step 9 of Algorithm 1 above to:

$$F(\vec{s}) \leftarrow \sum_{\vec{t} \in (substates(\vec{s}) - \{\vec{s}\})} P_{transition}(\vec{s}, \vec{t}) \cdot \max_{\vec{u} \in selectable_states(\vec{s}, \vec{t})} F(\vec{u})$$

Because F monotonically decreases as all of d_6, d_8, d_{10}, d_{12} increase up to these critical points, these two functions will be equal on states reachable from $[10, 1, 1, 1]$ and $[12, 0, 1, 1]$.

We then need to recursively reevaluate F on “bigger” states than $[9, 1, 1, 1]$, shown in Fig. 4.

What The Creator Really Asked For: The chance of getting a perfect run the commercial game using the very best strategy (smoke 'em except leave a second and final

$d_6 \backslash d_8$	0	1	$d_6 \backslash d_8$	0	1
0	0.08333	0.02908	0	0.02347	0.01087
1	0.03819	0.01726	1	0.01406	0.007809
2	0.02231	0.01209	2	0.009946	0.006252
3	0.0154	0.009464	3	0.007849	0.005385
4	0.01188	0.00798	4	0.006672	0.004879
5	0.009888	0.007088	5	0.005971	0.00458
6	0.008677	0.006532	6	0.005542	0.004408
7	0.007907	0.006181	7	0.005278	0.004318
8	0.007404	0.00596	8	0.00512	0.004283
9	0.007071	0.005825	9	0.005033	0.004284
10	0.006851	0.00575	10	0.004994	0.004312
11	0.006707	0.005715	11	0.004988	0.004356
12	0.006618	0.005709	12	0.005005	0.004413

Figure 3: Expected values for $(d_{10}, d_{12}) = (0, 1)$ (left) and $(1, 1)$ (right).

unsupported six on $[d_6 \in \{10, 11, 12\}, d_8 = 1, d_{10} = 1, d_{12} = 1]$) is about 0.004446, or a little more than 4.4%, a 0.73% relative improvement on the smoke 'em strategy.

So it turns out Mr. Cox's students are correct - you may indeed want to "save" future six rolls if your rarer "onesies" remain. The difference on the commercial game begins at $[10, 1, 1, 1]$, and occurs exactly when you roll exactly 2 sixes on 6-sided dice and no other maxes.

- $[10, 1, 1, 1]$: Naive: Take both sixes (EV = 0.0043116537). **Optimal**: Take only 1 six (EV = 0.0043120029).
- $[11, 1, 1, 1]$: Naive: Take both sixes (EV = 0.0043564817). **Optimal**: Take only 1 six (EV = 0.0043662025).
- $[12, 1, 1, 1]$: Naive: Take both sixes (EV = 0.0044132805). **Optimal**: Take only 1 six (EV = 0.0044456381).

It's nice to see the game is complex and non monotonic even in this constrained scenario. There are likely higher configurations yielding other critical points in the space e , especially as more dimensions (dice types) are added. Though the difference is minuscule in the commercial game, expanding $\vec{S}$ to something like $S^* = [24, 2, 2, 2]$ yields a wider EV difference between simple and optimal strategies².

² $F(S^*) = 0.0033078600$ using the basic strategy and 0.0038614244 using the optimal, a 17% improvement

$d_6 \backslash d_8$	0	1
0	0.02347	0.01087
1	0.01406	0.007809
2	0.009946	0.006252
3	0.007849	0.005385
4	0.006672	0.004879
5	0.005971	0.00458
6	0.005542	0.004408
7	0.005278	0.004318
8	0.00512	0.004283
9	0.005033	0.004284
10	0.004994	0.004312 $\rightarrow$ 0.004312
11	0.004988	0.004356 $\rightarrow$ 0.004366
12	0.005005	0.004413 $\rightarrow$ 0.004446

Figure 4: Expected values for $(d_{10}, d_{12}) = (0, 1)$ (left) and $(1, 1)$ (right).

2 The Second Quest: $d_6 \rightarrow \infty$

But can we say nothing more definitive than that? Looking at the left column on the first table (Fig. 2) with only one type of die, it looks like a higher dice count both decreases our chance of winning, but may also do so asymptotically.

Let's run $\vec{0} + 100e_1$ and see.

It certainly looks like there is an asymptote to which $f(s)$ converges. This will be the thrust of the rest of this paper: examining the infinite game of $F(s)$ with one type of die.

Lemma 2.1 (Easy Cases Lemma). *If $p = 0$, $F(r) = 0$. If $p = 1$, $F(r) = 1$. If $s = 2$, for any $r > 0$, $F(r) = \frac{1}{2}$.*

Proof: Though we won't be flipping infinite ($p = 0$) or 1-sided ($p = 1$), dice, these algebraic identities are obvious (can't succeed, can't fail, respectively) and will be useful later.

Base case for $s=2$: $F(1) = \frac{1}{2}$ since a perfect game occurs if and only if the coin flips 2 instead of 1.

r	$P(r)$
0	1.000000000000000
1	0.166666666666667
2	0.074074074074074
3	0.041923868312757
4	0.028091341512981
5	0.021074519393544
6	0.017084426737672
7	0.014629278661270
8	0.013031406311202
9	0.011947904221376
10	0.011190349681739
$\vdots$	$\vdots$
90	0.009033527660227
91	0.009033527656758
92	0.009033527653985
93	0.009033527651768
94	0.009033527649995
95	0.009033527648578
96	0.009033527647444
97	0.009033527646538
98	0.009033527645813
99	0.009033527645234
100	0.009033527644770

Table 1: Values of $P(n)$ for selected indices

Inductive case for $s=2$: Assuming $F(k) = \frac{1}{2}$ for all $0 < k < r$, and $F(0) = 1$:

$$\sum_{k=1}^{r-1} \binom{r}{k} p^k q^{r-k} F(r-k) + \binom{r}{r} p^r q^0 F(0) = \quad (1)$$

$$(2^r - 2) \frac{1}{2^r} \left(\frac{1}{2}\right) + \frac{1}{2^r} 2 \left(\frac{1}{2}\right) = \quad (2)$$

$$(2^r) \frac{1}{2^r} \left(\frac{1}{2}\right) = \frac{1}{2} \quad (3)$$

table of contents TODO

- Initial Problem Statement: Perfect games
- Solution for BADG commercial version

- One die type, $n \rightarrow \infty$
- Easy Cases
- Lazy Upper bound
- Lazy Lower bound (Pochhammer)
- Extremely Lazy Lower bound (closed form)
- Wrong attempt: $F(t) < F(t - 1)$?
- $F(n)$ exhibits provably asymptotic behavior

Theorem 2.2. *For an unbounded set of $r > 0$ standard s -sided dice, $s \geq 2$, the probability $F(r)$ of getting a perfect game has a lower bound of $\exp(-\frac{q}{(1-q)^2})$, if we define $q := 1 - \frac{1}{s}$.*

Proof:

- First, while at stage k , where k dice remain, the probability $P_s(k)$ of continuing to some stage $j < k$ is clearly $1 - q^k$.
- The probability $F(k)$ of passing all such stages is greater than or equal to $P^* = \prod_{i=1}^k P_s(k)$, since a successful traversal may involve skipping one or more stages, and P^* assumes we no stages.
- Therefore, $F(k) \geq \prod_{i=1}^k P_s(k) = \prod_{i=1}^k (1 - q^k)$.
- This means $\log(F(k)) \geq \sum_{i=1}^k \log(P_s(k)) = \sum_{i=1}^k \log(1 - q^k)$.

This $F(k)$ is apparently defined as the **q-Pochhammer symbol** $(q; q)_\infty = \prod_{i=1}^k (1 - q^k)$, with known convergence properties. I'll instead attempt to tackle convergence without the aid of these theorems.

Often, $\log(1 - z)$ is approximated by $-z$ near zero, but we are most concerned with z near one, since $1 - z$ approaches 0, and $\log(1 - z)$ explodes. Therefore, let's look at the expansion of $\log(1 - z)$:

- Identity 1: $f(z) = -\frac{1}{1-z} = -1 - z - z^2 - \dots$
- Identity 2: $\int f(z) = \log(1 - z) = -z - \frac{z^2}{2} - \frac{z^3}{3} - \dots$

So when summing $\sum_{i=1}^k \log(1 - q^i)$, we can instead sum

$$\log(F(k)) = \log(1 - q^1) + \log(1 - q^2) + \log(1 - q^3) + \dots \quad (4)$$

$$\log(F(k)) = (-q - \frac{q^2}{2} - \frac{q^3}{3} + \dots) + (-q^2 - \frac{q^4}{2} - \frac{q^6}{3} + \dots) + (-q^3 - \frac{q^6}{2} - \frac{q^9}{3} + \dots) + \dots \quad (5)$$

$$> (-q - q^2 - q^3 + \dots) + (-q^2 - q^4 - q^6 + \dots) + (-q^3 - q^6 - q^9 + \dots) + \dots \quad (6)$$

$$= \frac{-q}{1-q} + \frac{-q^2}{1-q^2} + \frac{-q^3}{1-q^3} + \dots \quad (7)$$

$$> \frac{-q}{1-q} + \frac{-q^2}{1-q} + \frac{-q^3}{1-q} + \dots \quad (8)$$

$$= \frac{1}{1-q} [-q - q^2 - q^3 - \dots] \quad (9)$$

$$= \frac{-q}{(1-q)^2} \quad (10)$$

- (2) follows from (1) by substitution using Identity1 above.
- (3) follows from (2) since the denominators of (2) are larger.
- (4) follows from (3) when dividing each sequence by $-q^k$ and applying identity 2.
- (5) follows from (4) since the denominators of (4) are larger.
- (6) and (7) follow from (5) by dividing out $\frac{1}{1-q}$ and applying Identity 1.

Since $\log(F(k)) > \frac{-q}{(1-q)^2}$, $F(k) > \exp(\frac{-q}{(1-q)^2})$.

s	p	q	bound
2	$\frac{1}{2}$	$\frac{1}{2}$	$\exp(-2) \approx .135$
3	$\frac{1}{3}$	$\frac{1}{3}$	$\exp(-6) \approx .00248$
4	$\frac{1}{4}$	$\frac{1}{4}$	$\exp(-12) \approx 6.14 \times 10^{-6}$
10	$\frac{1}{10}$	$\frac{1}{10}$	$\exp(-90) \approx 8.19 \times 10^{-40}$

Table 2: Very loose lower bounds on $F(k = \infty)$

This is a thoroughly unsatisfying lower bound, as we know $F(2) = \frac{1}{2}$.

Definition 2.3 $((q; q)_\infty$: The q -pochhammer symbol). *The q -pochhammer symbol $(q; q)_\infty = \prod_{k=1}^{\infty} (1 - q^k)$ is a known convergent quantity.*

q	$(q; q)_\infty$	$\exp\left(-\frac{q}{(1-q)^2}\right)$
1/2	2.8879×10^{-1}	1.3534×10^{-1}
2/3	6.9272×10^{-2}	2.4788×10^{-3}
3/4	1.5545×10^{-2}	6.1442×10^{-6}
4/5	3.3680×10^{-3}	2.0612×10^{-9}
5/6	7.1400×10^{-4}	9.3576×10^{-14}
6/7	1.4913×10^{-4}	5.7495×10^{-19}
7/8	3.0815×10^{-5}	4.7809×10^{-25}
8/9	6.3155×10^{-6}	5.3802×10^{-32}
9/10	1.2861×10^{-6}	8.1940×10^{-40}

However, this does provide a positive lower bound, and brings up this interesting fact: **Though increasing the number of sides of the dice can drive $F(k)$ to zero, increasing k to ∞ will never set $F(k)$ to zero.**

If we can couple this with the proof of the next theorem, we can state that $F(k)$ has an asymptotic limit as $k \rightarrow \infty$ given some q .

Calculating by hand:

- $F(0) = 1$
- $F(1) = p$
- $F(2) = p^2(1 + 2q) = p^2(3 - 2p)$
- $F(3) = p^3(13 - 27p + 21p^2 - 6p^3)$
- $F(4) = p^4(75 - 316p + 606p^2 - 664p^3 + 432p^4 - 156p^5 + 24p^6)$
- $F(1) - F(0) = p - 1 = -(1 - p)$
- $F(2) - F(1) = p^2(3 - 2p) - p = (-1)p(1 - 2p)(1 - p)$
- $F(3) - F(2) = (-1)3p^2(1 - p)^3(1 - 2p)$
- $F(4) - F(3) = -p^3(1 - 2p)(p - 1)^2(12p^4 - 48p^3 + 72p^2 - 50p + 13)$

Theorem 2.4 (Incorrect Blind Alley: $F(n) < F(n - 1)$). *If $p < \frac{1}{2}$, $F(n - 1) > F(n)$ for $n > 1$.*

- TODO: Show calculation of various functions directly.
- Show calculation of $p = \frac{1}{2} - \epsilon$, $q = \frac{1}{2} - \epsilon$ with $p = 0.495$.

- Show interesting observations: growth of polynomial
- $R(t)$ is starting at position t , throwing $t - 1$ dice.
- $F(t) = p(1 + q^{t-1})F(t - 1) + qR(t) = pF(t - 1) + pq^{t-1}F(t - 1) + qR(t)$
- If $qR(t) < qF(t - 1) - pq^{t-1}F(t - 1)$, same as $F(t - 1) - R(t) > pq^{t-2}F(t - 1)$, then $F(t) < F(t - 1)$.
- In the special case $q = p = \frac{1}{2}$, then $F(t - 1) - R(t) = p^{t-1}(1) - p^{t-1} \cdot p + \text{other zero factors} = p^t = pq^{t-2} \cdot \frac{1}{2}$.
- So we're satisfied since $F(t) = \frac{1}{2}$ for $t \geq 1$.
- If $p \leq \frac{1}{3}$, $F(t - 1) - R(t) = [p^{t-1}F(0) - p^{t-1}F(1) + \text{other positive factors}] = p^{t-1}q + \epsilon$.

$$t = 3, F(3) - R(4) > pq^{4-1}F(4 - 1) \quad (11)$$

$$p^3F(0) - p^3F(1) \quad (12)$$

$$\binom{3}{2}p^2qF(1) - \binom{3}{2}p^2qF(2) \quad (13)$$

$$\binom{3}{1}p^2qF(2) - \binom{3}{2}p^2qF(3) \quad (14)$$

$$(15)$$

3 Interesting Findings

- Calculating the asymptotic behavior of a combination of s-sided dice and t-sided dice isn't particularly interesting; if $s > t$, this less than $F_{1/t}(\infty)$ and more than $F_{1/s}(\infty)$.
- no closed form - triangle exponent
- non monotone commercial game
- non monotone single game
- pochhammer bounds, ultimate convergence
- can be made zero in p, not in n
- note: different than best strategy (since others have knowledge)