

## Review

## Analytical and mathematical methods for revealing hidden details in ancient manuscripts and paintings: A review



Anna Tonazzini<sup>a</sup>, Emanuele Salerno<sup>a</sup>, Zienab A. Abdel-Salam<sup>b</sup>, Mohamed Abdel Harith<sup>b</sup>, Luciano Marras<sup>c</sup>, Asia Botto<sup>d</sup>, Beatrice Campanella<sup>d</sup>, Stefano Legnaioli<sup>d</sup>, Stefano Pagnotta<sup>d</sup>, Francesco Poggiali<sup>d</sup>, Vincenzo Palleschi<sup>d,\*</sup>

<sup>a</sup> National Research Council of Italy, Institute of Information Science and Technologies "Alessandro Faedo", Via G. Moruzzi, 1, Pisa, Italy

<sup>b</sup> National Institute of Laser Enhanced Sciences, Cairo University, Cairo, Egypt

<sup>c</sup> Art Test Studio di Luciano Marras, via del Martello 14, 56121 Pisa, Italy

<sup>d</sup> National Research Council of Italy, Applied and Laser Spectroscopy Laboratory, Institute of Chemistry of Organometallic Compounds, Via G. Moruzzi, 1, Pisa, Italy

## HIGHLIGHTS

- Methods for revealing hidden details in ancient manuscripts and paintings are presented.
- Different experimental approaches are described.
- The most effective techniques of image analysis are introduced.
- Special attention is given to multi-spectral imaging and blind separation methods.
- Several case studies are presented.

## GRAPHICAL ABSTRACT



## ARTICLE INFO

## Article history:

Received 18 October 2018

Revised 7 January 2019

Accepted 8 January 2019

Available online 12 January 2019

## Keywords:

Image analysis

Cultural heritage

Archaeology

Multispectral imaging

Ancient manuscripts

Blind separation techniques

## ABSTRACT

In this work, a critical review of the current nondestructive probing and image analysis approaches is presented, to revealing otherwise invisible or hardly discernible details in manuscripts and paintings relevant to cultural heritage and archaeology. Multispectral imaging, X-ray fluorescence, Laser-Induced Breakdown Spectroscopy, Raman spectroscopy and Thermography are considered, as techniques for acquiring images and spectral image sets; statistical methods for the analysis of these images are then discussed, including blind separation and false colour techniques. Several case studies are presented, with particular attention dedicated to the approaches that appear most promising for future applications. Some of the techniques described herein are likely to replace, in the near future, classical digital photography in the study of ancient manuscripts and paintings.

© 2019 The Authors. Published by Elsevier B.V. on behalf of Cairo University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

This review is focused on the analytical techniques and methods that have been used to date and are likely to be used extensively in the near future to reveal hidden details in cultural heritage artefacts. Technically, all techniques used in archaeometry (the discipline that applies scientific methods to the study of cul-

Peer review under responsibility of Cairo University.

\* Corresponding author.

E-mail address: [vincenzo.palleschi@cnr.it](mailto:vincenzo.palleschi@cnr.it) (V. Palleschi).

<https://doi.org/10.1016/j.jare.2019.01.003>

2090-1232/© 2019 The Authors. Published by Elsevier B.V. on behalf of Cairo University.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

tural heritage and archaeology) are aimed at revealing what is not evident and cannot be determined without the use, as a matter of fact, of specific analytical techniques and methods. To further define the scope of this paper, the discussion will be focused on the techniques that may help improve the interpretation and understanding of the manuscripts and paintings, not considering techniques such as radiography or X-ray tomography, which, although extremely interesting for their applications, are typically used for acquiring bulk information, well below the visible surface of the objects under study. These techniques would require, because of their complexity and importance, a full separate review. In the following, probing methods, instrumentation and digital processing techniques for the analysis of the artefact surface are described and discussed. Particular attention is devoted to spectrally resolved imaging methods (reflectometry, fluorescence), although methods based on thermal or elemental analysis of artefacts are also considered when functional to the recovery of surface information. Among the processing techniques, only those that operate on sets of images (separation techniques, false colour imaging, etc.), rather than those operating on single greyscale images (image enhancement technique, segmentation, etc.) are discussed. In the conclusion, a brief discussion of the most promising approaches in the field is presented.

Keywords, including image analysis, cultural heritage, archaeology, multispectral imaging, ancient manuscripts, and blind separation techniques were searched through database “Scopus” from 1968 to 2018.

## Probing techniques

### Multispectral imaging – MSI

Multispectral imaging is one of the most popular techniques for the study of cultural heritage and archaeological findings. One main advantage of MSI is that it is a non-invasive technique and therefore can be applied to any artwork, despite its possible fragility. Although the spectral resolution of this type of analysis is, in general, limited (typical bandwidths are of the order of 50 nm or even larger in multispectral imaging and of the order of 10–20 nm in so-called hyperspectral imaging), the amount of information that can be obtained is extremely high, considering the high spatial resolution of the images that can be obtained through very simple experimental setups.

MSI, originally developed for remote sensing applications, began to be applied extensively in art conservation and art history in the early 1990s [1–7], as it can reveal information in an artwork that cannot be seen by the human eye.

A multispectral image can be described as a set, or *cube*, of images of the same scene taken over different spectral ranges, i.e., at different wavelengths in the electromagnetic spectrum, including light outside of the visible range, such as infrared (IR) and ultraviolet (UV) light. Reflectance and fluorescence images can be independently acquired but treated simultaneously [8].

From an experimental point of view, an image in a multispectral cube (a channel) can either be isolated by specific filters [9] or using appropriate narrow-band illumination systems [10]. Scanning systems can also be used [11].

In the method's simplest realization, four images of the subject under study are acquired in the blue, green, red and infrared spectral bands. In most cases, the infrared image is the one carrying the most information because of the unique ability of infrared radiation to penetrate the object surface, allowing for the visualization of otherwise invisible details such as underdrawings and *pentimenti* in canvas and panel paintings [11–14]. Infrared imaging is also important for other applications because of the possible

enhancement of features deriving from the different infrared reflectivity of the subject's constituent materials. The improvement of readability of degraded manuscripts in the infrared image was demonstrated, for example, in the recovery of the burnt Heracleum scrolls [15] and in revealing several hidden characters obscured by exposure to moisture in the Dead Sea Scrolls [16]. In other cases, e.g., in palimpsests or archaeological wall paintings, imaging in the UV spectral range often succeeds in providing additional information [17,18].

In addition to highlighting hidden patterns, multispectral images and their further elaborations can also provide information on the materials used for the realization of a painting [19–22], on illumination conditions and pigment identification [23,24], and for monitoring the conservation of cultural heritage objects [25–27]. Grifoni *et al.* [28,29] recently proposed the use of spectrally resolved images as photogrammetric sources for building 3D models of paintings that would carry information about the painted surface in depth structure (see Fig. 1).

Multispectral and hyperspectral imaging, along with techniques for the digital processing of the acquired images, has been the focus of several national and international projects devoted to the study of precious artworks of great historical value. In most cases, dedicated imaging equipment has been devised and implemented. The study of ancient manuscripts and, among them, of palimpsests in particular, is one of the fields where multispectral imaging has demonstrated to give excellent results. The Archimedes palimpsest project [30] has been one of the most important efforts in this field, aimed to the recovery from a XIII century prayer book of the erased and overwritten text of a earlier copy of two lost treatises of Archimedes. In the framework of this project, Easton *et al.* [31–33] introduced an MSI acquisition system that makes use of narrow-band LEDs.

Other projects have been carried out regarding palimpsests in Europe, among which one of the most important and comprehensive has been the European Project “Digitale Palimpsestforschung” (2001–2004) [34]. The project was led by the University of Hamburg and gathered the efforts of more than 50 partner institutions from 26 countries to study a large part of existing Greek palimpsests, with the help of newly developed digital technologies. From a technological perspective, new multispectral capture systems were among the results of this project, along with a set of basic image enhancement techniques and computer tools for document archiving and cataloguing.

In the project “Critical Edition of the New Sinaitic Glagolitic Euchology (Sacramentary) Fragments with the Aid of Modern Technologies” [35,36], a portable MSI system has been used to image the Sinaitic Glagolitic manuscripts. This system consists of two multispectral LED panels and two different cameras, a greyscale camera with sensitivity from the UV to the near-infrared (NIR), and a traditional RGB camera utilized for UV fluorescence and visible-light imaging.

Also in the field of manuscript analysis, Bianco *et al.* [37] described an MSI apparatus that uses a filter wheel consisting of eight different optical filters and a monochromatic camera for simultaneous 3D acquisition. Lettner *et al.* [38] introduced a similar MSI system with an extra single-lens reflex camera. Rapantzikos and Balas [39] used a system with optical filters for imaging over 34 narrow spectral bands.

The efficacy of MSI for the analysis of texts was evaluated in [40].

### X-ray fluorescence (XRF)

X-ray fluorescence (XRF) can be used to support MSI for the non-destructive elemental analysis of those parts of the artwork in which MSI is ineffective. This technique consists of acquiring



Fig. 1. 3D multispectral reconstruction of the surface of a painting. (a) RGB; (b) UV-Vis fluorescence; (c) infrared [29].

the spatial distribution of the chemical elements [41,42] of large samples.

When used to probe ancient manuscripts, XRF can distinguish among different types of iron-gall inks due to its high sensitivity to iron concentration and the impurities (typically of copper and zinc) that characterize different batches of ink or inks of different periods [43].

Experiments on the use of XRF for reading palimpsests have been conducted within a project carried out by the Centre for the Study of Manuscript Cultures at the University of Hamburg and the University library of Leipzig, in cooperation with the Hamburg synchrotron radiation laboratory (HASYLAB) and the German electron synchrotron (DESY). Within that project, monochromatized, high-flux X-ray fluorescence techniques were employed [34].

Knox et al. [44] analysed the capabilities of MSI and XRF in revealing hidden characters in various types of damaged parchment manuscripts. The conclusions of this analysis were that the nature of the inks and the condition of the parchment would influence what regions of the optical spectrum would reveal characters. As general rules, it was concluded that infrared illumination is good for revealing carbon-based ink on blackened parchment, ultraviolet fluorescence (and sometimes reflectance) can enhance erased characters, and finally, X-ray fluorescence can detect iron gall ink that is completely covered by optically opaque materials.

### Thermography

Infrared thermography [45] can also be used effectively to reveal the presence of hidden patterns or structures in a large variety of objects. Multispectral imaging normally detects the near-IR radiation emerging from the objects under test (0.75–1.4  $\mu\text{m}$  wavelength range); the typical wavelengths used for thermography belong to the thermal IR range (3–15  $\mu\text{m}$ ). Techniques based on infrared thermography are capable of detecting subsurface features in the investigated object by mapping the temperature distribution at its surface and can be implemented in different experimental arrangements [46]. A first distinction can be made on the possible presence of an artificial illumination system: *passive* techniques evaluate temperature differences naturally occurring at the investigated surface, whereas *active* techniques rely on the temporal evolution of surface temperature induced by suitably timed and filtered artificial heating systems (usually flash

lamps). Both of these approaches have already been used to investigate many classes of objects relevant to cultural heritage, such as historical stone and masonry artefacts [47–49], archaeological findings and ancient documents [46,50–52]. In particular, active pulsed thermography has been successfully applied to non-invasively highlight the presence of ancient texts in parchment book bindings, to characterize the status of conservation of painted decorations and to reveal the presence of possible *pentimenti* under the painted surfaces [46,52].

### Raman and LIBS imaging

The effectiveness of using micro-Raman imaging, a technique that provides information about the molecular structure of surfaces, together with MSI, was evaluated by Maybury et al. [53] in an analysis of Armenian manuscripts. Deneckere et al. [54] used micro-Raman imaging coupled with the elemental technique of micro-XRF to acquire elemental and molecular images of a Belgian porcelain card. Bicchieri et al. [55] used MSI, FT-IR spectroscopy, micro-Raman and micro-XRF for the analysis of a degraded 18th century manuscript. Finally, Botteon et al. [56] used a variation of Raman microscopy called spatially offset Raman spectroscopy (SORS) to demonstrate the possibility of recovering painted images hidden by, for example, graffiti or other types of overpainting. In fact, any experimental technique capable of reconstructing spectrally resolved images of the surface of cultural heritage artefacts can be used for recovering hidden information. Elemental images obtained using Laser-induced Breakdown Spectroscopy (LIBS), a micro-destructive spectroscopic technique, were reported in [57] and [58]. Among these, non-destructive approaches are obviously preferable, when applicable.

### Digital processing techniques

#### Statistical analysis and source separation

Among the image processing techniques typically explored using MSI data, statistical analysis and dimension reduction have proven to be powerful tools for further enhancing and detecting hidden patterns in artworks or removing unwanted interferences. Dimension reduction can be both unsupervised, as in blind source



separation (BSS) techniques [59,60], and supervised, as in Fisher linear discriminant analysis (LDA) [61].

Indeed, unsupervised dimension reduction techniques, such as principal component analysis (PCA) and independent component analysis (ICA), linearly combine highly correlated spectral images to produce a different set of images that are uncorrelated and show decreasing variance. Furthermore, the output channels of ICA are statistically independent. Thus, the main principle underlying the enhancement capabilities of dimension reduction techniques is that, while the spectral components of an image are usually spatially correlated, the individual patterns (or classes, or sources) superposed onto the image are usually much less correlated. Hence, decorrelating the colour components gives a different representation, where the now orthogonal components of the image could coincide with single classes [62–66]. For example (Fig. 2), for palimpsests containing mixtures of two different texts and possibly further information layers (parchment texture, mould, etc.), dimensionality reduction often results in images in which each shows a single layer separated from the others. Because the statistical independence requirement of ICA is a stronger condition than the assumption of uncorrelation of PCA, it is possible that signals that are not well segmented by PCA may be separable by ICA or by ICA applied to a set of principal components, as done in [67].

PCA, ICA and other orthogonalization methods, when applied to multispectral images, can increase the readability of degraded texts [68–70] or reveal hidden features not apparent in any of the individual input images, as in [66], in which a hidden text was shown to exist in a XVIII century painting, or in [71], in which the existence of many otherwise hidden details was demonstrated in wall paintings found in the Etruscan *Tomb of the Monkey* (Chiusi,


Italy, 5th Century BCE). BSS techniques were particularly important in investigating the lost mural paintings in the Etruscan *Tomb of Blue Demons* (Tarquinia, Italy, 5th Century BCE), as reported in a recent paper by Adinolfi et al. [18]. In that study, a set of visible, infrared and fluorescence images was treated statistically by BSS algorithms, revealing a magnificent hunting scene with three hunters, a wild boar, a deer, a dog, and two felids, where the naked eye could perceive only a white wall (a detail of the scene, depicting the wild boar and the head of a hunter, is shown in Fig. 3).

If the mutual independence assumption is not tenable, the ICA-based strategies for source separation may fail. One option in this case could be to rely on dependent component analysis (DCA), a class of model-based techniques that exploit other possible properties of sources or mixtures to reach their goal [72,73]. This type of technique has been applied extensively to fields such as remote sensing [74,75] and medical imaging [76]. Although some DCA approaches could be employed to analyse different types of cultural heritage-related images, only one proposal is present in the open literature in which a DCA approach is used for the digital restoration of colour images of double-sided documents [77].

Fisher linear discriminant analysis (LDA) can also be applied to reduce the dimensions of multispectral scans and to enhance degraded writings. Because Fisher LDA is a supervised dimension reduction tool, it is necessary to label a subset of multispectral data. To this end, in [78], a semi-automated label generation step was introduced based on an automated detection of text lines. This approach is thus based not only on spectral information, as in PCA and ICA, but also on spatial information and, when tested on two Slavonic manuscripts, has yielded better performance compared with that of unsupervised techniques.



**Fig. 2.** Folio 16v–17r of the Archimedes palimpsest. (a) RGB image under strobe lamp illumination. (b) Second component output (contrast-enhanced) from the  $2 \times 2$  PCA of the red and blue colour channels, revealing the underwritten text and drawings (© The owner of the Archimedes Palimpsest, licensed for use under creative Commons Attribution 3.0 Unported Access Rights. Image processing: The Institute of Information Science and Technologies, National Research Council of Italy).



**Fig. 3.** Detail of the hunting scene (a wild boar, running from right to left) recovered using BSS in the *Tomb of Blue Demons* in Tarquinia, using MSI and BSS. On the right, the visible image of the wall. Note the improvement in readability of the wild boar (muzzle with ear and fang is evidenced in the yellow circle) and of the head of one of the hunters (red circle) and vegetation at his right.

The self-organizing maps (SOMs) method, introduced by Kohonen at the end of last century [79], represents a completely different approach to the blind separation problem. SOMs are artificial neural networks that achieve separation through the similarity of the (optical) properties of materials, which are represented in an  $n$ -dimensional space by the coordinates of the corresponding (hyper)-colours. Unlike in BSS, the number of images that can thus be extracted by a multispectral set can be greater than the number of images in the original set. No hypothesis is made on the linearity of the model, and the information layers are separated through an iterative, competitive process between the neurons that “move” in the hypercolour space arriving, after convergence, to assume the coordinates of the centroid of the corresponding cluster. This method requires the definition of a metric that determines the similarity of the hypercolours defining different materials (Euclidean, Angular, Manhattan, etc. [80]). The number of neurons is also left to the decision of the operator, based on the expected number of different materials/optical responses in the physical object [65]. An important advantage of the SOM approach is that no dimensional reduction must be performed for the classification of materials; the position of the neurons in the hyperspace represents the “prototype” of the optical properties of the corresponding material. The hypercolour associated with each pixel in the MSI image may have components corresponding to visible and infrared reflectivity, fluorescence, and elemental or molecular information. Applications of SOMs to elemental images obtained by the LIBS technique were reported by Pagnotta et al. [57,58] (Fig. 4).

In addition to BSS, SOM and LDA, non-blind spectral unmixing has proven useful in text analysis. This approach, popular in remote-sensed hyperspectral image analysis, is based on the availability of a dictionary containing the typical spectral signatures of the materials of interest and unmixing strategies such as spectral angle mapping (SAM) [81] capable of labelling the different sensed regions as belonging to specified classes [82,83]. In document image analysis, pixel regions belonging to specific object classes, e.g., parchment, mould, overwriting, or erased text, are first identified by the user. An algorithm then computes the class membership of each pixel in the image based on the similarity of its spectrum to each of the specified classes. Although intensive both in terms of human interaction and computation time, this method was applied with success to the Archimedes Palimpsest [84–86]. Spectral unmixing for document image analysis can be particularly

useful in situations in which different feature spectra are known or can be determined a priori, as in remote sensing for earth observation, where the spectra are known from field or laboratory measurements.

#### Pseudocolour imaging

A simple approach for enhancing hidden features in an artwork when appropriate non-visible bands are available is a rendering technique called *false colour* or *pseudocolour*. Because only three spectral bands can be displayed in a colour image, three suitable images are selected from the multispectral set and superimposed in the form of a (false) colour image. The most common combination of the multispectral images is infrared, red and green (IrRG), although the combination infrared, green and blue (IrGB) [87] is also used. The procedure implies that one of the visible colour channels is discarded (the blue band in IrRG false colour imaging or the red band in IrGB imaging), and the information it contains is not present in the pseudocolour image (Fig. 5).

Pseudocolour imaging can be generalized in several ways. For example, to render the image data used in the study of the Archimedes Palimpsest, images captured through a blue filter under ultraviolet illumination, where the underwriting was mostly visible, and through a red filter under tungsten light, where the underwriting had nearly disappeared, were combined to render the overwriting in black and the underwriting in a reddish tint. In the resulting pseudocolour image, the two texts were then perceptually well separated because they featured highly contrasting colours, enabling the reader to distinguish between them [88].

When using data reduction methods, if layers are perfectly separated, each feature class would dominate the greyscale range in the related output channel, while pixels belonging to the other feature classes would exhibit the same grey value and thus merge with the background in that channel. More realistically, PCA or ICA may not succeed in separating features if the different feature patterns are not truly orthogonal or independent. Thus, in palimpsests, traces of overwriting usually appear in those channels in which the erased underwriting is most visible, and the variation in statistics across the scene, for example to variations in erasures, makes the erased text often appear in more than one output channel with varying intensity. This fact can, however, be exploited to generate pseudocolour rendering of extracted component images,






Fig. 4. SOM segmentation of a set of elemental images obtained on a Roman mortar sample using  $\mu$ -LIBS [57]. The yellow square in the figure indicates the zone analysed.




Fig. 5. Schematic representation of the procedure used for building IrRG and IrGB false colour images from a multispectral set of images.

where small variations in grey value may appear as large changes in colour, thus further improving the readability of the erased text [89].

A similar effect can be obtained by changing the pseudocolour rendering by varying the hue angle or by creating weighted combinations of images, including results from ICA and PCA and possibly original image bands.

Legnaioli et al. [71] introduced a false colour imaging technique called chromatic derivative imaging (ChromaDI), which exploits the subtraction of consecutive couples of 4 consecutive spectral images, namely, G-B, R-G and IR-R. This method was developed with the intent of building a false colour image that would take

into account the information from all multispectral images acquired, without excluding *a priori* one of the four images in the multispectral set. The ChromaDI image provides information on the changes in reflectivity of an object with wavelength. With respect to the canonical false colour image, the differences between the optical behaviour of various pigments are enhanced, taking into account the changes occurring while passing from short wavelengths (blue band, which is more sensitive to surface details) to longer ones (green and red bands) in the visible image (see Fig. 6).

ChromaDI has been successfully applied to images of a Roman painted sarcophagus, III century A.D., and to images of a mural




Fig. 6. Schematic representation of the procedure used to build a ChromaDI image.

painting of an Etruscan tomb in Chiusi (Siena, Italy), among artefacts. The method can easily be generalized to multispectral sets containing more than four images. For instance, in palimpsests, ChromaDI images could include one or more channels of UV fluorescence.

Another false colour imaging method, only experimented on paintings to date, aims at producing chromatically faithful pseudo-colour images, which maintain good readability of the information contained in the infrared band. Examples of the application of this technique include the multispectral images acquired for the *Pietà* of Agnolo Bronzino (1569, Florence) and the analysis and visualization of the multispectral data obtained from Etruscan mural paintings (*Tomb of the Monkey*, Siena, Italy, V century B.C.) [90]. The method is called gradient transfer and, through a regularization strategy, merges the information from the IR band into the RGB image, preserving at best the chromatic similarity with the visible image (Fig. 7).

A similar approach for image inpainting exploiting infrared information has been recently proposed by Calatroni et al. [91] for removing overpaintings in the visible image in the analysis of illuminated manuscripts and by Peng et al. [92] for mining patterns of painted cultural relics in ancient pottery and murals.

In the context of ancient manuscripts, e.g., palimpsests, the IR band could be substituted by the blue band of the UV fluorescence, where, presumably, the underwriting is best visible.


Even XRF data can take advantage of pseudocolour. For instance, in [93], a linear model was proposed to disentangle the four texts emerging from an XRF analysis of a recto-verso palimpsested manuscript. A pseudocolour rendering is then used to enhance the individual patterns in the resulting images. A non-linear model for the superposition of texts in recto-verso scanned manuscripts [94] is envisaged to further improve the result.

#### Colour spaces for RGB imaging

The potential of MSI and other imaging techniques for the analysis of cultural artefacts is currently widely recognized and demonstrated. Furthermore, portable and inexpensive equipment is available. Nevertheless, the efficient use of these instruments requires specialized operators and mechanical apparatuses for the correct alignment of the camera and the artwork. Thus, in the majority of cases, simpler-to-use acquisition devices operating in the visible spectrum alone are employed, and more specialized probing techniques are limited to artworks of particular importance. In recent years, extensive digitization campaigns have been conducted in most museums, libraries and archives around the world, mainly for conservation purposes. An enormous number of digital reproductions of artworks as high-resolution RGB images are thus available. This situation poses the problem of finding fast, efficient and easily deployed image processing techniques that can meet the two requirements of being suitable for routine use and effective in helping scholars in the study and analysis of the artwork at hand.

Manuscripts often contain patterns such as underwritings, stamps, or paper watermarks that represent the most significant information from a cultural and historical point of view for establishing authorship and origin. Such marks should thus be undisclosed and enhanced. As previously mentioned, in many cases, explorations in the near-infrared or UV band can be extremely useful in this respect, as can further elaborations of the multiplicity of multispectral/hyperspectral images.

However, sometimes representing the only available RGB images in different colour spaces can be an efficient tool for “simulating” views outside the visible range, and even without introducing additional information from, for example, infrared images,



**Fig. 7.** Detail of the *Pietà* of Agnolo Bronzino. (a) RGB image, (b) infrared Image and (c) merged “true colour infrared image” [90].

the method can disentangle interesting information contained in the visible spectrum from masking interferences.

Indeed, although the RGB colour representation is the most frequently used colour space in image processing, it presents some limitations in terms of maximization of image information content. Hence, in the literature, many different colour spaces have been developed for different image analysis tasks, such as object segmentation and edge detection [95]. Some of these spaces are particularly suitable for the analysis of degraded documents, often allowing for the enhancement of document content, the improvement of text readability, and the extraction of partially hidden features.

In 1989, Xerox Corporation proposed a colour encoding called YES [96]. The YES colour space is a linear transformation of the RGB vector that matches the physiology of the human visual system. The space separates colour information and intensity information. The three coordinates are an achromatic luminance channel that is a weighted sum of the RGB values, called  $Y$ , and two opponent-colour chromatic coordinates given by spectral differences: the  $E$  channel is proportional to red minus green, while the  $S$  channel is proportional to yellow minus blue. YES has been specifically employed for the enhancement of degraded ancient parchments. When imaging the Dead Sea Scrolls, it was found that the contrast in the  $E$  map was significantly augmented, and hidden characters were revealed [16]. Other authors claim that subtracting the green component from the red, hidden characters in charred documents can be revealed, exhibiting a performance similar to that obtained using the near-infrared band [97]. A possible explanation for this behaviour is that the red channel may have recorded some infrared information, which is separated from the rest by subtracting the “red” part contained in the green channel as well [98].

The OHTA colour space was derived to approximate the PCA of RGB components [99]. The fixed coefficients of the OHTA matrix were experimentally found by a statistical study of the uncorrelated colour components in a large population of images of typical real-world scenes. The three coordinates are an achromatic luminance channel that is a homogeneous weighted sum of the RGB

values, called  $O$ , and two chromatic coordinates given by spectral differences: the  $H$  channel is proportional to red minus blue, while the  $T$  channel is proportional to green minus magenta.

The YES and OHTA colour spaces and the red-minus-green and red-minus-blue operations were also useful for removing the bleed-through distortion in reddish documents [100]. The rationale for this application can be found by examining the histogram of this type of document, from which it can be observed that, in the background/bleed-through areas, red and green (or red and blue) are well separated, *i.e.*, their difference is large. Thus, red-green/red-blue return nearly equal, high values for both the background and the bleed-through pixels such that they merge; conversely, much lower values are obtained for the text, resulting in enhancement.

Note that dimension reduction techniques, such as PCA or ICA, when applied to RGB images, can be interpreted as adaptive colour representations, in which the new colours, *i.e.*, the components extracted, are mutually spatially uncorrelated or independent.

#### *Modelling Multispectral/Multiview images*

The use of statistical processing techniques to elaborate the multispectral/hyperspectral images of an artwork, with the aim of separating the various layers of information that it contains, implicitly assumes a linear, instantaneous data model. In other words, all available views of an artwork are considered linear combinations of a number of patterns. The recovery of individual patterns thus amounts to inverting this transformation. However, because the coefficients of the transformation are not known, *a priori* assumptions about the patterns must be exploited. Applying the various PCA and ICA operators corresponds to assuming mutual uncorrelation (or independence) between the patterns [62,63,101].

This basic linear instantaneous mixing model can be extended to account for nonlinearity, spatial non-stationarity, convolutional mixing, noise, *etc.*, to better adhere to the physical characteristics of specific instances of pattern superposition in artworks. For example, some of the abovementioned variants have been explored to model the phenomenon of text overlap in recto-verso



manuscripts affected by show-through or bleed-through distortion, with the aim of correcting the distortion. In such cases, solutions exploiting nonlinear ICA, non-negative matrix factorization, variational approaches, regularization, dependent component analysis, or other ad hoc strategies have been proposed [77,94,110–112,102–109].

## Conclusions and future perspectives

In this paper, experimental methods and analytical techniques that can help in recovering hidden details in cultural heritage artefacts are presented and discussed. These methods are particularly suited for the analysis of degraded texts, palimpsests and paintings but can also be applied, for example, to the study of geological materials, pottery and mortars.

Regardless of the experimental technique used, if a representative set of images can be obtained, processing methods can be applied to treat these images and extract meaningful information. Blind source separation techniques, self-organizing maps, and linear discriminant analysis provide statistical algorithms that can reveal hidden features that, although present in the input set, might not be observable in the individual channel images. These techniques can also be applied to simple RGB images, possibly with the help of freely available software, such as the D-stretch ImageJ plugin [113]. Once the image set is obtained, pseudocolour images can be obtained or, using new techniques based on the gradient transfer method, even colour faithful images, embedding otherwise invisible information, can be obtained. 3D multispectral models can also be recovered using digital photogrammetry. Many examples of the application of the above described techniques in restoration, archiving and documentation processes can already be found in recent literature [114–118].

With the progress of instrumentation (improved CCD cameras, illuminators, and non-optical imaging systems such as micro or macro XRF/LIBS elemental imaging, Raman molecular imaging, etc.) and the introduction of simpler, faster and more performant statistical algorithms for the treatment of large image sets, it is reasonable to expect that in the near future multispectral imaging and the related techniques described here will likely replace colour digital photography for quick and information-rich documentation and study of cultural heritage.

## Conflict of interest

*The authors have declared no conflict of interest*

## Compliance with Ethics Requirements

*This article does not contain any studies with human or animal subjects.*

## References

- [1] Baronti S, Casini A, Lotti F, Porcinai S. Multispectral imaging system for the mapping of pigments in works of art by use of principal-component analysis. *Appl Opt* 1998;37(8):1299.
- [2] Casini A, Lotti F, Piccolo M, Stefani L, Buzzegoli E. Image spectroscopy mapping technique for noninvasive analysis of paintings. *Stud Conserv* 1999;44(1):39–48.
- [3] Balas C, Papadakis V, Papadakis N, Papadakis A, Vazgiouraki E, Themelis G. A novel hyper-spectral imaging apparatus for the non-destructive analysis of objects of artistic and historic value. *J Cult Herit* 2003;4:330s–7s.
- [4] Fischer C, Kakoulli I. Multispectral and hyperspectral imaging technologies in conservation: current research and potential applications. *Rev Conserv* 2006;7:3–16.
- [5] Vilaseca M, Pujol J, Arjona M, de Lasarte M. Multispectral system for reflectance reconstruction in the near-infrared region. *Appl Opt* 2006;45(18):4241.
- [6] Cristoforetti G, Legnaioli S, Palleschi V, Salvetti A, Tognoni E. Optical chemical sensors for cultural heritage. In: *Optical Chemical Sensors NATO Science Series II: Mathematics, Physics and Chemistry*. p. 515–26.
- [7] Bonifazzi C, Carcagnì P, Fontana R, Greco M, Mastrianni M, Materazzi M, et al. A scanning device for VIS–NIR multispectral imaging of paintings. *J Opt A Pure Appl Opt* 2008;10(6):064011.
- [8] Marras L, Pelagotti A, Castaldi M, Carmagnola R, Adinolfi G. La Signora dei Demoni Azzurri. *Archeo* 2009;296:80–5.
- [9] Grifoni E, Briganti L, Marras L, Orsini S, Colombini MP, Legnaioli S, et al. The chemical-physical knowledge before the restoration: The case of “The Plague in Lucca”, a masterpiece of Lorenzo Viani (1882–1936). *Herit Sci* 2015;3(1).
- [10] Marengo E, Manfredi M, Zerbinati O, Robotti E, Mazzucco E, Gosetti F, et al. Technique based on LED multispectral imaging and multivariate analysis for monitoring the conservation state of the dead sea scrolls. *Anal Chem* 2011;83(17):6609–18.
- [11] Daffara C, Pampaloni E, Pezzati L, Barucci M, Fontana R. Scanning multispectral IR reflectography SMIRR: An advanced tool for art diagnostics. *Acc Chem Res* 2010;43(6):847–56.
- [12] Fontana R, Gambino MC, Greco M, Marras L, Materazzi M, Pampaloni E, et al. New high-resolution IR-color reflectography scanner for painting diagnosis. In: Salimbeni R, editor. *International Society for Optics and Photonics*; 2003. p. 108–15. Available from: <http://proceedings.spiedigitallibrary.org/proceeding.aspx?articleid=767715>.
- [13] van Asperen de Boer JRJ. A note on the use of an improved infrared vidicon for reflectography of paintings. *Stud Conserv* 1974;19:97–9.
- [14] Hollmann JC, Crause K. Digital imaging and the revelation of “hidden” rock art: Vaalekop Shelter, KwaZulu-Natal. *South African Humanit* 2011;23(1):55–76.
- [15] Chabries DM, Booras SW, Bearman GH. Imaging the past: recent applications of multispectral imaging technology to deciphering manuscripts. *Antiquity* 2003;77(296):359–72.
- [16] Knox K, Johnston R, Easton RL. Imaging the dead sea scrolls. *Opt Photonics News* 1997;8(8):30.
- [17] Bloechl K, Hamlin H, Easton, Jr. RL. Text recovery from the ultraviolet-fluorescent spectrum for treatises of the Archimedes Palimpsest. In: Stork DG, Coddington J, Bentkowska-Kafel A, editors. 2010. p. 753109. Available from: <http://proceedings.spiedigitallibrary.org/proceeding.aspx?doi=10.1117/12.838886>.
- [18] Adinolfi G, Carmagnola R, Cataldi M, Marras L, Palleschi V. Recovery of a lost wall painting at the Etruscan Tomb of the Blue Demons in Tarquinia (Viterbo, Italy) by multispectral reflectometry and UV fluorescence imaging. *Archaeometry* 2018. <http://doi.wiley.com/10.1111/arcm.12423>.
- [19] Toque JA, Sakatoku Y, Ide-Ektessabi A. Pigment identification by analytical imaging using multispectral images. In: 2009 16th IEEE International Conference on Image Processing (ICIP). IEEE; 2009. p. 2861–4. Available from: <http://ieeexplore.ieee.org/document/5414508/>.
- [20] Melis M, Miccoli M, Quarta D. Multispectral hypercolorimetry and automatic guided pigment identification: some masterpieces case studies. In: Pezzati L, Targowski P, editors. 2013. p. 87900W. <http://proceedings.spiedigitallibrary.org/proceeding.aspx?doi=10.1117/12.2020643>.
- [21] Comelli D, Nevin A, Valentini G, Osticioli I, Castellucci EM, Toniolo L, et al. Insights into Masolino's wall paintings in Castiglione Olona: Advanced reflectance and fluorescence imaging analysis. *J Cult Herit* 2011;12(1):11–8.
- [22] Mazzeo R, Palazzi CE, Rocchetti M, Sciuotto G. Computer-assisted pigment identification in artworks. In: *Proceedings of the IASTED European Conference on Internet and Multimedia Systems and Applications*. p. 266–71.
- [23] Paviotti A, Forsyth DA, Cortelazzo GM. Lightness recovery for pictorial surfaces. *Int J Comput Vis* 2011;94(1):54–77.
- [24] Ricciardi P, Delaney JK, Glinesman L, Thoury M, Facini M, de la Rie ER. Use of visible and infrared reflectance and luminescence imaging spectroscopy to study illuminated manuscripts: pigment identification and visualization of underdrawings. In: Pezzati L, Salimbeni R, editors. *International Society for Optics and Photonics*; 2009. p. 739106. <http://proceedings.spiedigitallibrary.org/proceeding.aspx?doi=10.1117/12.827415>.
- [25] Marengo E, Manfredi M, Zerbinati O, Robotti E, Mazzucco E, Gosetti F, et al. Development of a technique based on multi-spectral imaging for monitoring the conservation of cultural heritage objects. *Anal Chim Acta* 2011;706(2):229–37.
- [26] Toque JA, Komori M, Murayama Y, Ide-Ektessabi A. Analytical Imaging of Traditional Japanese Paintings Using Multispectral Images. 2010. p. 119–32. [http://link.springer.com/10.1007/978-3-642-11840-1\\_9](http://link.springer.com/10.1007/978-3-642-11840-1_9).
- [27] Rogerio-Candelera MA, Jurado V, Laiz L, Saiz-Jimenez C. Laboratory and in situ assays of digital image analysis based protocols for biodeteriorated rock and mural paintings recording. *J Archaeol Sci* 2011;38(10):2571–8.
- [28] Grifoni E, Legnaioli S, Lorenzetti G, Pagnotta S, Palleschi V. Image based recording of three-dimensional profiles of paint layers at different wavelengths. *Eur J Sci Theol* 2017;13(2).
- [29] Grifoni E, Legnaioli S, Nieri P, Campanella B, Lorenzetti G, Pagnotta S, et al. Construction and comparison of 3D multi-source multi-band models for cultural heritage applications. *J Cult Herit* 2018.
- [30] The Archimedes Palimpsest Project. <http://www.archimedespalimpsest.org/>.
- [31] Easton RL, Knox KT, Christens-Barry WA. Multispectral imaging of the Archimedes palimpsest. In: 32nd Applied Imagery Pattern Recognition Workshop, 2003 Proceedings. IEEE; 2003. p. 111–6.
- [32] Easton RL, Christens-Barry WA, Knox KT. Ten years of lessons from imaging of the archimedes palimpsest. In: *ΕΙΚΟΝΟΛΟΓΙΑ Symposium on Digital Imaging of Ancient Textual Heritage: Technological Challenges and Solutions*. p. 3–26.

- [33] Easton RL, Knox KT, Christens-Barry WA, Boydston K, Toth MB, Emery D, et al. Standardized system for multispectral imaging of palimpsests. In: Stork DG, Coddington J, Bentkowska-Kafel A, editors. 2010. p. 75310D. <http://proceedings.spiedigitallibrary.org/proceeding.aspx?doi=10.1117/12.839116>.
- [34] Digitale Palimpsestforschung. Available from: <http://www.teuchos.uni-hamburg.de/palimpsestuntersuchung>.
- [35] Gau M, Miklas H, Lettner M, Sablatnig R. Image acquisition and processing routines for damaged manuscripts. *Digit Mediev* 2010;6:715–36.
- [36] Čamba A, Gau M, Hollaus F, Fiel S, Sablatnig R. Multispectral imaging, image enhancement, and automated writer identification in historical manuscripts. *Manuscr Cult* 2014;7:83–91.
- [37] Bianco G, Bruno F, Tonazzini A, Salerno E, Savino P, Sroubek F, et al. A framework for virtual restoration of ancient documents by combination of multispectral and 3D imaging. *Proc 8th Eurographics Italian Chapter Conference*, 2010.
- [38] Lettner M, Diem M, Sablatnig R, Miklas H. Registration of multispectral manuscript images as a prerequisite for computer-aided script description. In: 12th Computer Vision Winter Workshop. p. 51–8.
- [39] Rapantzikos K, Balas C. Hyperspectral imaging: potential in non-destructive analysis of palimpsests. In: IEEE International Conference on Image Processing 2005. IEEE; 2005. p. II-618.
- [40] Giacometti A, Campagnolo A, MacDonald L, Mahony S, Robson S, Weyrich T, et al. The value of critical destruction: Evaluating multispectral image processing methods for the analysis of primary historical texts. *Digit Scholarch Humanit* 2015;32(1):fqv036.
- [41] Janssens K, Dik J, Cotte M, Susini J. Photon-based techniques for nondestructive subsurface analysis of painted cultural heritage artifacts. *Acc Chem Res* 2010;43(6):814–25.
- [42] Bruker. M6 Jetstream, Large Area Micro X-ray Fluorescence Spectrometer brochure [Internet]. Available from: <https://www.bruker.com/products/x-ray-diffraction-and-elemental-analysis/micro-xrf-and-txrf/m6-jetstream/overview.html>.
- [43] Bruni S, Caglio S, Guglielmi V, Poldi G. The joined use of n.i. spectroscopic analyses – FTIR, Raman, visible reflectance spectrometry and EDXRF – to study drawings and illuminated manuscripts. *Appl Phys A*. 2008;92(1):103–8.
- [44] Knox KT, Easton RL, Christens-Barry WA. Image restoration of damaged or erased manuscripts. *Proceedings of the 16th European Signal Processing Conference (EUSIPCO)*, 2008.
- [45] Meola C, Carlomagno GM. Recent advances in the use of infrared thermography. *Meas Sci Technol* 2004;15(9):R27–58. , <http://stacks.iop.org/0957-0233/15/i=9/a=R01?key=crossref.8744e902a0758ad6dfd1b2f1a5168ac>.
- [46] Mercuri F, Orazi N, Paoloni S, Cicero C, Zammit U, Mercuri F, et al. Thermography applied to the study of cultural heritage. *Appl Sci* 2017;7(10):1010.
- [47] Grinzato E, Bison PG, Marinetti S. Monitoring of ancient buildings by the thermal method. *J Cult Herit* 2002;3(1):21–9.
- [48] Avdelidis N, Moropoulou A. Applications of infrared thermography for the investigation of historic structures. *J Cult Herit* 2004;5(1):119–27.
- [49] Moropoulou A, Labropoulos KC, Delegou ET, Karoglou M, Bakolas A. Non-destructive techniques as a tool for the protection of built cultural heritage. *Constr Build Mater* 2013;48:1222–39.
- [50] Colombo G, Mercuri F, Scudieri F, Zammit U, Marinelli M, Volterri R. Thermographic analysis of parchment bindings. *Restauratorology* 2005;26(2):92–104.
- [51] Marinelli M, Mercuri F, Scudieri F, Zammit U, Colombo G. Thermographic study of microstructural defects in deteriorated parchment sheets. *J Phys IV* 2005;125:527–9.
- [52] Mercuri F, Zammit U, Orazi N, Paoloni S, Marinelli M, Scudieri F. Active infrared thermography applied to the investigation of art and historic artefacts. *J Therm Anal Calorim* 2011;104(2):475–85.
- [53] Maybury IJ, Howell D, Terras M, Viles H. Comparing the effectiveness of hyperspectral imaging and Raman spectroscopy: a case study on Armenian manuscripts. *Herit Sci* 2018;6(1):42.
- [54] Deneckere A, Vekemans B, Voorde L, Paeppe P, Vincze L, Moens L, et al. Feasibility study of the application of micro-Raman imaging as complement to micro-XRF imaging. *Appl Phys A* 2012;106(2):363–76.
- [55] Bicchieri M, Monti M, Piantanida G, Sodo A. All that is iron-ink is not always iron-gall! *J Raman Spectrosc* 2008;39(8):1074–8.
- [56] Botteon A, Conti C, Realini M, Colombo C, Matousek P. Discovering hidden painted images: subsurface imaging using microscale spatially offset raman spectroscopy. *Anal Chem* 2017;89(1):792–8.
- [57] Pagnotta S, Lezzerini M, Ripoll-Seguer L, Hidalgo M, Grifoni E, Legnaioli S, et al. Micro-Laser-Induced Breakdown Spectroscopy (Micro-LIBS) Study on Ancient Roman Mortars. *Appl Spectrosc* 2017;71(4):721–7.
- [58] Pagnotta S, Legnaioli S, Campanella B, Grifoni E, Lezzerini M, Lorenzetti G, et al. Micro-chemical evaluation of ancient potsherds by  $\mu$ -LIBS scanning on thin section negatives. *Mediterr Archaeol Archaeom* 2018;18(5):171–8.
- [59] Bell AJ, Sejnowski TJ. An information-maximization approach to blind separation and blind deconvolution. *Neural Comput* 1995;7(6):1129–59.
- [60] Cichocki A, Amari S. Adaptive Blind Signal and Image Processing. Chichester, UK: John Wiley & Sons, Ltd; 2002.
- [61] McLachlan GJ. Discriminant Analysis and Statistical Pattern Recognition (Wiley Series in Probability and Statistics). Hoboken, NJ, USA: John Wiley & Sons, Inc.; 1992.
- [62] Tonazzini A, Salerno E, Mochi M, Bedini L. Blind Source Separation Techniques for Detecting Hidden Texts and Textures in Document Images. Berlin, Heidelberg: Springer; 2004. p. 241–8.
- [63] Tonazzini A, Bedini L, Salerno E. Independent component analysis for document restoration. *Doc Anal Recognit* 2004;7(1).
- [64] Tonazzini A, Bianco G, Salerno E. Registration and Enhancement of Double-Sided Degraded Manuscripts Acquired in Multispectral Modality. In: 2009 10th International Conference on Document Analysis and Recognition [Internet]. IEEE; 2009. p. 546–50.
- [65] Salerno E, Tonazzini A, Grifoni E, Lorenzetti G, Legnaioli S, Lezzerini M, et al. Analysis of multispectral images in cultural heritage and archaeology. *J Appl Laser Spectrosc* 2014;1:22–7.
- [66] Legnaioli S, Grifoni E, Lorenzetti G, Marras L, Pardini L, Palleschi V, et al. Enhancement of hidden patterns in paintings using statistical analysis. *J Cult Herit* 2013;14(3):S66–70.
- [67] Salerno E, Tonazzini A, Bedini L. Digital image analysis to enhance underwritten text in the Archimedes palimpsest. *Int J Doc Anal Recognit* 2007;9(2–4):79–87.
- [68] Tonazzini A, Bedini L, Salerno E. Digital analysis of damaged documents by ICA techniques. In: Gori M, Marinai S, editors. Artificial Neural Networks in Pattern Recognition. p. 33–8.
- [69] Hollaus F, Gau M, Sablatnig R. Multispectral image acquisition of ancient manuscripts. Berlin, Heidelberg: Springer; 2012. p. 30–9.
- [70] Knox KT, Easton RL, Christens-Barry WA, Boydston K. Recovery of handwritten text from the diaries and paper s of David Livingstone. 2011. p. 786909. Available from: <http://proceedings.spiedigitallibrary.org/proceeding.aspx?doi=10.1117/12.877135>.
- [71] Legnaioli S, Lorenzetti G, Cavalcanti GH, Grifoni E, Marras L, Tonazzini A, et al. Recovery of archaeological wall paintings using novel multispectral imaging approaches. *Herit Sci* 2013;1.
- [72] Xiang Y, Peng D, Yang Z. Blind Source Separation (SpringerBriefs in Electrical and Computer Engineering). Singapore: Springer Singapore; 2015.
- [73] Li R, Li H, Wang F. Dependent component analysis: concepts and main algorithms. *J Comput* 2010;5(4):589–97.
- [74] Caiafa CF, Salerno E, Proto AN, Fiumi L. Blind spectral unmixing by local maximization of non-Gaussianity. *Signal Process* 2008;88(1):50–68.
- [75] Du Q, Kopriva I. Dependent component analysis for blind restoration of images degraded by turbulent atmosphere. *Neurocomputing* 2009;72(10–12):2682–92.
- [76] Kopriva I, Peršin A, Puizina-Ivić N, Mirić L. Dependent component analysis based approach to robust demarcation of skin tumors. In: Plum JPW, Dawant BM, editors. 2009. p. 72594Q. Available from: <http://proceedings.spiedigitallibrary.org/proceeding.aspx?doi=10.1117/12.806404>.
- [77] Tonazzini A, Bedini L. Restoration of recto-verso colour documents using correlated component analysis. *EURASIP J Adv Signal Process* 2013;2013(1):58.
- [78] Hollaus F, Gau M, Sablatnig R. Enhancement of Multispectral Images of Degraded Documents by Employing Spatial Information. In: 2013 12th International Conference on Document Analysis and Recognition. IEEE; 2013. p. 145–9. Available from: <http://ieeexplore.ieee.org/document/6628601>.
- [79] Kohonen T. The self-organizing map. *Neurocomputing* 1998;21(1–3):1–6.
- [80] McCune B, Grace JB, Urban DL. Analysis of ecological communities. *MjM Software Design*; 2002. 300 p.
- [81] Kruse FA, Lefkoff AB, Boardman JW, Heidebrecht KB, Shapiro AT, Barloon PJ, et al. The spectral image processing system (SIPS)—interactive visualization and analysis of imaging spectrometer data. *Remote Sens Environ* 1993;44(2–3):145–63.
- [82] Schwarz J, Staenz K. Adaptive threshold for spectral matching of hyperspectral data. *Can J Remote Sens* 2001;27(3):216–24.
- [83] Dennison PE, Halligan KQ, Roberts DA. A comparison of error metrics and constraints for multiple endmember spectral mixture analysis and spectral angle mapper. *Remote Sens Environ* 2004;93(3):359–67.
- [84] Easton RL, Christens-Barry WA, Knox KT. Spectral image processing and analysis of the Archimedes Palimpsest. In: Proceedings of the 19th European Signal Processing Conference, EUSIPCO 2011. p. 1440–4.
- [85] Noel W. Archimedes and company. *Br Acad Rev* 2007;10.
- [86] Walvoord DJ, Easton RL. Digital transcription of the archimedes palimpsest [applications corner]. *IEEE Signal Process Mag* 2008;25(4):100–4.
- [87] Moon T, Schilling MR, Thirkettle S. A note on the use of false-color infrared photography in conservation. *Stud Conserv* 1992;37(1):42.
- [88] Easton RL, Kelbe D. Statistical processing of spectral imagery to recover writings from erased or damaged manuscripts. *Manuscr Cult* 2014;7:35–46.
- [89] Salerno E, Tonazzini A. Extracting erased text from palimpsests by using visible light. In: Proceedings of the 4-th Intl Congr Science and Technology for the Safeguard of Cultural Heritage in the Mediterranean Basin. p. 532–5.
- [90] Grifoni E, Campanella B, Legnaioli S, Lorenzetti G, Marras L, Pagnotta S, et al. A new Infrared True-Color approach for visible-infrared multispectral image analysis. *ACM J Comp Cult Her* 2018 (in press).
- [91] Calatroni L, d'Autume M, Hocking R, Panayotova S, Parisotto S, Ricciardi P, et al. Unveiling the invisible: mathematical methods for restoring and interpreting illuminated manuscripts. *Herit Sci* 2018;6(1):56. <https://doi.org/10.1186/s40494-018-0216-z>.
- [92] Peng J, Yu K, Wang J, Zhang Q, Wang L, Fan P. Mining painted cultural relic patterns based on principal component images selection and image fusion of hyperspectral images. *J Cult Herit* 2018. <https://doi.org/10.1016/j.culher.2018.09.008>.


- [93] Bergmann U, Knox KT. Pseudo-color enhanced x-ray fluorescence imaging of the Archimedes Palimpsest. In: Berkner K, Likforman-Sulem L, editors. 2009. p. 724702. <http://proceedings.spiedigitallibrary.org/proceeding.aspx?doi=10.1117/12.806053>.
- [94] Sharma G. Show-through cancellation in scans of duplex printed documents. *IEEE Trans Image Process*. 2001;10(5):736–54.
- [95] Sharma G. Show-through cancellation in scans of duplex printed documents. *IEEE Trans Image Process* 2001;10(5):736–54.
- [96] Corporation X. The Xerox Color Encoding Standard; 1989.
- [97] Swift R. Analysis of the Spectra of Degraded Documents. Technology. 2001. Available from: <https://ritdml.rit.edu/handle/1850/5773>.
- [98] Knox KT. Enhancement of overwritten text in the Archimedes Palimpsest. In: Stork DG, Coddington J, editors. 2008. p. 681004. <http://proceedings.spiedigitallibrary.org/proceeding.aspx?doi=10.1117/12.766679>.
- [99] Ohta Y-I, Kanade T, Sakai T. Color information for region segmentation. *Comput Graph Image Process* 1980;13(3):222–41.
- [100] Tonazzini A. Color space transformations for analysis and enhancement of ancient degraded manuscripts. *Pattern Recognit Image Anal* 2010;20(3):404–17.
- [101] Tonazzini A, Salerno E, Bedini L. Fast correction of bleed-through distortion in grayscale documents by a blind source separation technique. *Int J Doc Anal Recognit* 2007;10(1):17–25.
- [102] Tan Chew Lim, Cao R, Shen Peiyi. Restoration of archival documents using a wavelet technique. *IEEE Trans Pattern Anal Mach Intell* 2002;24(10):1399–404.
- [103] Huang Yi, Brown MS, Dong Xu. User-Assisted Ink-Bleed Reduction. *IEEE Trans Image Process* 2010;19(10):2646–58.
- [104] Rowley-Brooke R, Pitie F, Kokaram A. A Non-parametric Framework for Document Bleed-through Removal. In: 2013 IEEE Conference on Computer Vision and Pattern Recognition [Internet]. IEEE; 2013. p. 2954–60.
- [105] Tonazzini A, Gerace I, Martinelli F. Multichannel blind separation and deconvolution of images for document analysis. *IEEE Trans Image Process* 2010;19(4):912–25.
- [106] Merrikh-Bayat F, Babaie-Zadeh M, Jutten C. Using non-negative matrix factorization for removing show-through. 2010. p. 482–9. Available from: [http://link.springer.com/10.1007/978-3-642-15995-4\\_60](http://link.springer.com/10.1007/978-3-642-15995-4_60).
- [107] Martinelli F, Salerno E, Gerace I, Tonazzini A. Nonlinear model and constrained ML for removing back-to-front interferences from recto-verso documents. *Pattern Recognit* 2012;45(1):596–605.
- [108] Salerno E, Martinelli F, Tonazzini A. Nonlinear model identification and see-through cancelation from recto-verso data. *Int J Doc Anal Recognit* 2013;16(2):177–87.
- [109] Merrikh-Bayat F, Babaie-Zadeh M, Jutten C. Linear-quadratic blind source separating structure for removing show-through in scanned documents. *Int J Doc Anal Recognit* 2011;14(4):319–33.
- [110] Almeida MSC, Almeida LB. Nonlinear separation of show-through image mixtures using a physical model trained with ICA. *Signal Process* 2012;92(4):872–84.
- [111] Moghaddam RF, Cheriet M. A variational approach to degraded document enhancement. *IEEE Trans Pattern Anal Mach Intell* 2010;32(8):1347–61.
- [112] Tonazzini A, Savino P, Salerno E. A non-stationary density model to separate overlapped texts in degraded documents. *Signal, Image Video Process* 2015;9(S1):155–64.
- [113] Harman J. <http://www.dstretch.com/>
- [114] tanco F, Battiato S, Gallo G. Digital imaging for cultural heritage preservation : analysis, restoration, and reconstruction of ancient artworks [Internet]. CRC Press; 2011. 487 p. Available from: [https://books.google.it/books/about/Digital\\_Imaging\\_for\\_Cultural\\_Heritage\\_Pr.html?id=QHnBxQ2xhGQC&redir\\_esc=y](https://books.google.it/books/about/Digital_Imaging_for_Cultural_Heritage_Pr.html?id=QHnBxQ2xhGQC&redir_esc=y).
- [115] McCarthy J. Multi-image photogrammetry as a practical tool for cultural heritage survey and community engagement. *J Archaeol Sci* 2014;43:175–85.
- [116] Dimoulas CA, Kalliris GM, Chatzara EG, Tsipias NK, Papanikolaou GV. Audiovisual production, restoration-archiving and content management methods to preserve local tradition and folkloric heritage. *J Cult Herit* 2014;15(3):234–41.
- [117] Tuna G, Zogo R, Çiftçi EE, Demirelli B, Tuna A. Identification, preservation and management of cultural heritage of Edirne, Turkey by means of a web-based application. *J Balk Libr Union* 2015;3(2):36–41.
- [118] Zacharopoulos A, Hatzigiannakis K, Karamaounas P, Papadakis VM, Andrianakis M, Melessanaki K, et al. A method for the registration of spectral images of paintings and its evaluation. *J Cult Herit* 2018;29:10–8.



**Anna Tonazzini** is a senior researcher at the Institute of Information Science and Technologies, National Research Council of Italy, in Pisa. She coordinated several Projects in Image Processing and Analysis, Neural Networks and Learning, Computational Biology and Document Analysis, and is co-author of over 100 peer reviewed papers. She was also the ISTI responsible for the UE Project ISYREADET and the national Flagship Project InterOmics. She chaired the EUSIPCO2008 Special Session on restoration of degraded document images, and edited the special Issue on image and video processing for cultural heritage of the *Eurasip Journal on Image and Video Processing* in 2009. She supervised many theses in computer science, mathematics and information engineering, two Ercim fellowships, and various post-doctoral positions. She was an editor of *Digital Signal Processing*, a member of the IASTED Technical Committee on Image Processing, and a program committee member in several international conferences.



**Emanuele Salerno** graduated from electronic engineering at the University of Pisa, Italy, and joined the National Research Council of Italy in 1987. Currently, he is a senior researcher at the Institute of Information Science and Technologies in Pisa, Italy. He has been working in microwave tomography, monitoring of combustion processes, computer vision for robot guidance, and astrophysical imaging. His present scientific interests are in applied inverse problems, image processing, IT for cultural heritage, and computational biology. He has been teaching instrumentation and measurements and microwaves at the university of Pisa, and is a senior member of the IEEE, the Signal Processing Society, a member of the Italian Federation of Electrical and Information Engineering (AEIT), and a fellow of the Electromagnetics Academy.



**Zienab A. Abdel-Salam** Associate Professor at the National Institute of Laser Enhanced Science (NILES), Cairo University. Obtained her M.Sc. and Ph.D. degrees in Laser Applications in Biotechnology from NILES, Cairo University. Her current research interests include Applied laser spectroscopy in animal production, animal health and nutrition, archaeology, microbiology and chemometrics. Dr. Abdel-Salam has published many papers in internationally reputable journals.




**Mohamed Abdel Harith** is a Professor of Laser Physics at the National Institute of Laser Enhanced Science (NILES). He obtained his degree of Ph.D. in experimental physics from Dresden Technical University in Germany. He founded a scientific school in laser spectroscopy at NILES where 32 students obtained their master degrees and 25 students have had their doctorate degrees. He served as the head of department Egyptian Society of Optical Sciences and Applications (ESOSA) and then the institute's vice Dean. From 2003 to 2005 he was the Dean of NILES. He has more than 128 publications with more than 1500 citations and *h*-index 24. Member of the European Society of Physics EPS and senior member of the Optical Society (OSA). Chairman of the Board of Directors of the Awarded the University of Cairo award of Appreciation in 2009 and the Cairo University award of Excellence for the year 2016.





Luciano Marras is a professional in art diagnostics, business owner of Art-Test di Luciano Marras professional studio in Pisa, specialized on optical techniques on cultural heritage. He completed his Ph.D. in physics at Padua University and afterwards a research grant on multispectral techniques at CNR-INOA, Florence. He is author or co-author of nearly 45 articles, congress contributions and book chapters. He is Physics teacher in high schools, and he has other teaching experience in Master courses for Cultural Heritage. His interests range on all the imaging and spectrometric techniques in diagnostics applied on cultural heritage, and currently he is focused on the recovery of hardly damaged archaeological wall paintings.



Asia Botto has recently obtained a MSc degree in chemistry at the University of Pisa. Her work focused on the characterization and analysis of natural organic colorants dyes, used to dye textile materials, using the SERS (surface enhanced Raman spectroscopy) technique.




Beatrice Campanella received her PhD in analytical chemistry from the University of Pisa in 2016. She is currently post-doctoral fellow at the National Research Council – Institute of Chemistry of Organometallic Compounds (CNR – ICCOM) in Pisa. Her research is focused on the application of non/micro-invasive spectroscopic techniques (Raman, SERS, LIBS, XRF, and multispectral imaging) to the chemical characterization of modern and contemporary artworks.



Stefano Legnaioli has a master's degree in physics and PhD in Chemistry. He's a researcher at the Italian National Council for Research - Institute for the Chemistry of OrganoMetallic Compounds since 2008. He is co-author of more than 115 peer reviewed papers (h-index: 30 source Scopus), and has participated in various national and international conferences. Based on the classification of the European Research Council, its research activity can be classified within the following sectors: PE2, PE4, SH5\_1, SH6\_1. His research experience lies in the field of laser spectroscopy, particularly LIBS (Laser Induced Breakdown Spectroscopy), Raman, SERS, XRF, and Multispectral Imaging techniques. The fields of application concern the analysis of materials, environmental protection, the study and conservation of

cultural heritage. Over the years he has shown special aptitude for laboratory activity, both in the development of new instrumentation and in measurement procedures and data processing with chemometric techniques.



Stefano Pagnotta is a PhD Candidate in Earth Sciences at University of Florence with a research project in “ $\mu$ -LIBS scanner for Cultural Heritage Geomaterials”. He is an Anthropology Open Journal editorial board member. He has a Md in Archaeology (Prehistory and Archaeometry Studies) and a Bd in Conservation in Cultural Heritage (Techniques and Diagnostics for Cultural Heritage Materials). Dr. Pagnotta publications include 38 peer-reviewed journal articles with 180 citations, and 2 book chapters. His interests and research fields are wide: Laser-Induced Breakdown Spectroscopy, XRF, XRD, micro-Raman, Multispectral Imaging and Optical Microscopy, Artificial Intelligence developments, Features extraction from Images. He has numerous skills: digital photography, Matlab scripting, Data Mining, Multispectral Imaging, 3D printing, Algorithms, and Scientific Writing.



Francesco Poggialini is a PhD candidate in Chemistry at Scuola Normale Superiore in Pisa, working with the ALS Lab group of the Italian National Research Council in Pisa. He received his MSc in Analytical Chemistry at the University of Pisa. His research interests focus on laser spectroscopies, nanomaterials, and cultural heritage.



Vincenzo Palleschi is a Physicist, Senior Researcher at the Institute of Chemistry of Organometallic Compounds and Head of the Applied and Laser Spectroscopy Laboratory at Research Area of CNR in Pisa (Italy). Dr. Palleschi is a world-renowned expert in LIBS; he organized the first LIBS International Conference in Pisa, in year 2000 and was the Chairman and organizer of the 9th EMSLIBS Conference in Pisa, in June 2017. Besides LIBS, Dr. Palleschi has also experience in X-Ray Fluorescence analysis, micro-Raman spectroscopy, Multispectral Imaging, 3D photogrammetry and Chemometrics. He has published more than 200 papers in ISI journals, which received more than 6000 citations, and the book 'Laser-Induced Breakdown Spectroscopy: Principles and Applications' (eds. A.Miziolek, I. Schechter and V.Palleschi, CUP 2006). His h-index is 40 (Source: Scopus). He teaches at the University of Pisa the Courses of Analytical Chemistry III, Solid State Physicochemical Methods and Archaeometry and at the University of Turin the Course of Physical Methods for Restoration/Multispectral Analysis. In 2012 he had obtained the qualification as Full Professor in Experimental Physics of the Matter.