

The complexity of homomorphisms of signed graphs

Florent Foucaud
(Universitat Politècnica de Catalunya, Barcelona)

joint work (in progress) with:

Richard Brewster (Thompson Rivers U., Kamloops)
Pavol Hell (Simon Fraser U., Vancouver)
Reza Naserasr (U. Paris-Sud, Orsay)


JGA 2013

Graph homomorphisms


Definition - Graph homomorphism from G to H

Mapping from $V(G)$ to $V(H)$ which **preserves adjacency**.

If it exists, we note $G \rightarrow H$.



Target graph: $H = C_5$




Graph homomorphisms


Definition - Graph homomorphism from G to H

Mapping from $V(G)$ to $V(H)$ which **preserves adjacency**.

If it exists, we note $G \rightarrow H$.



Target graph: $H = C_5$




Graph homomorphisms


Definition - Graph homomorphism from G to H

Mapping from $V(G)$ to $V(H)$ which **preserves adjacency**.

If it exists, we note $G \rightarrow H$.



Target graph: $H = C_5$




Graph homomorphisms


Definition - Graph homomorphism from G to H

Mapping from $V(G)$ to $V(H)$ which **preserves adjacency**.

If it exists, we note $G \rightarrow H$.



Target graph: $H = C_5$




Graph homomorphisms


Definition - Graph homomorphism from G to H

Mapping from $V(G)$ to $V(H)$ which **preserves adjacency**.

If it exists, we note $G \rightarrow H$.



Target graph: $H = C_5$




Graph homomorphisms


Definition - Graph homomorphism from G to H

Mapping from $V(G)$ to $V(H)$ which **preserves adjacency**.

If it exists, we note $G \rightarrow H$.



Target graph: $H = C_5$




Graph homomorphisms


Definition - Graph homomorphism from G to H

Mapping from $V(G)$ to $V(H)$ which **preserves adjacency**.

If it exists, we note $G \rightarrow H$.



Target graph: $H = C_5$




Graph homomorphisms


Definition - Graph homomorphism from G to H

Mapping from $V(G)$ to $V(H)$ which **preserves adjacency**.

If it exists, we note $G \rightarrow H$.



Target graph: $H = C_5$




Graph homomorphisms


Definition - Graph homomorphism from G to H

Mapping from $V(G)$ to $V(H)$ which **preserves adjacency**.

If it exists, we note $G \rightarrow H$.



Target graph: $H = C_5$




Graph homomorphisms


Definition - Graph homomorphism from G to H

Mapping from $V(G)$ to $V(H)$ which **preserves adjacency**.

If it exists, we note $G \rightarrow H$.



Target graph: $H = C_5$




Graph homomorphisms


Definition - Graph homomorphism from G to H

Mapping from $V(G)$ to $V(H)$ which **preserves adjacency**.

If it exists, we note $G \rightarrow H$.



Target graph: $H = C_5$




Graph homomorphisms


Definition - Graph homomorphism from G to H

Mapping from $V(G)$ to $V(H)$ which **preserves adjacency**.

If it exists, we note $G \rightarrow H$.



Target graph: $H = C_5$




Graph homomorphisms


Definition - Graph homomorphism from G to H

Mapping from $V(G)$ to $V(H)$ which **preserves adjacency**.

If it exists, we note $G \rightarrow H$.



Target graph: $H = C_5$




Graph homomorphisms


Definition - Graph homomorphism from G to H

Mapping from $V(G)$ to $V(H)$ which **preserves adjacency**.

If it exists, we note $G \rightarrow H$.



Target graph: $H = C_5$




Graph homomorphisms


Definition - Graph homomorphism from G to H

Mapping from $V(G)$ to $V(H)$ which **preserves adjacency**.

If it exists, we note $G \rightarrow H$.



Target graph: $H = C_5$




Remark: Homomorphisms generalize **proper vertex-colourings**

$$G \rightarrow K_k \iff G \text{ is } k\text{-colourable}$$

Signed graphs: definitions


Signature Σ of graph G : assignment of $+$ or $-$ sign to each edge of G .
 Σ : set of $-$ edges.



$$\Sigma = \{12, 34\}$$


Signed graphs: definitions

Signature Σ of graph G : assignment of $+$ or $-$ sign to each edge of G .
 Σ : set of $-$ edges.




$$\Sigma = \{12, 34\}$$

Re-signing operation at v : switch sign of each edge incident to v




Signed graphs: definitions

Signature Σ of graph G : assignment of $+$ or $-$ sign to each edge of G .
 Σ : set of $-$ edges.



$$\Sigma = \{12, 34\}$$


Re-signing operation at v : switch sign of each edge incident to v



Signatures Σ, Σ' are **equivalent** ($\Sigma \equiv \Sigma'$) if one can be obtained from the other with **re-signings**. (equivalently: changing signs along an edge-cut)


Signed graphs: definitions

Signature Σ of graph G : assignment of $+$ or $-$ sign to each edge of G .
 Σ : set of $-$ edges.




$$\Sigma = \{12, 34\}$$

Re-signing operation at v : switch sign of each edge incident to v




Signatures Σ, Σ' are **equivalent** ($\Sigma \equiv \Sigma'$) if one can be obtained from the other with **re-signings**. (equivalently: changing signs along an edge-cut)




Signed graphs: definitions

Signature Σ of graph G : assignment of $+$ or $-$ sign to each edge of G .
 Σ : set of $-$ edges.




$$\Sigma = \{12, 34\}$$

Re-signing operation at v : switch sign of each edge incident to v




Signatures Σ, Σ' are **equivalent** ($\Sigma \equiv \Sigma'$) if one can be obtained from the other with **re-signings**. (equivalently: changing signs along an edge-cut)




Signed graphs: definitions

Signature Σ of graph G : assignment of $+$ or $-$ sign to each edge of G .
 Σ : set of $-$ edges.





$$\Sigma = \{12, 34\}$$

Re-signing operation at v : switch sign of each edge incident to v




Signatures Σ, Σ' are **equivalent** ($\Sigma \equiv \Sigma'$) if one can be obtained from the other with **re-signings**. (equivalently: changing signs along an edge-cut)




Signed graphs: definitions

Signature Σ of graph G : assignment of $+$ or $-$ sign to each edge of G .
 Σ : set of $-$ edges.





$$\Sigma = \{12, 34\}$$

Re-signing operation at v : switch sign of each edge incident to v




Signatures Σ, Σ' are **equivalent** ($\Sigma \equiv \Sigma'$) if one can be obtained from the other with **re-signings**. (equivalently: changing signs along an edge-cut)




Signed graphs: definitions

Signature Σ of graph G : assignment of $+$ or $-$ sign to each edge of G .
 Σ : set of $-$ edges.






$$\Sigma = \{12, 34\}$$

Re-signing operation at v : switch sign of each edge incident to v




Signatures Σ, Σ' are **equivalent** ($\Sigma \equiv \Sigma'$) if one can be obtained from the other with **re-signings**. (equivalently: changing signs along an edge-cut)




Signed graphs: definitions

Signature Σ of graph G : assignment of $+$ or $-$ sign to each edge of G .
 Σ : set of $-$ edges.






$$\Sigma = \{12, 34\}$$

Re-signing operation at v : switch sign of each edge incident to v




Signatures Σ, Σ' are **equivalent** ($\Sigma \equiv \Sigma'$) if one can be obtained from the other with **re-signings**. (equivalently: changing signs along an edge-cut)




Signed graphs: definitions

Signature Σ of graph G : assignment of $+$ or $-$ sign to each edge of G .
 Σ : set of $-$ edges.







$$\Sigma = \{12, 34\}$$

Re-signing operation at v : switch sign of each edge incident to v




Signatures Σ, Σ' are **equivalent** ($\Sigma \equiv \Sigma'$) if one can be obtained from the other with **re-signings**. (equivalently: changing signs along an edge-cut)




Signed graphs: definitions

Signature Σ of graph G : assignment of $+$ or $-$ sign to each edge of G .
 Σ : set of $-$ edges.




$$\Sigma = \{12, 34\}$$

Re-signing operation at v : switch sign of each edge incident to v




Signatures Σ, Σ' are **equivalent** ($\Sigma \equiv \Sigma'$) if one can be obtained from the other with **re-signings**. (equivalently: changing signs along an edge-cut)




Signed graphs: definitions

Signature Σ of graph G : assignment of $+$ or $-$ sign to each edge of G .
 Σ : set of $-$ edges.




$$\Sigma = \{12, 34\}$$

Re-signing operation at v : switch sign of each edge incident to v




Signatures Σ, Σ' are **equivalent** ($\Sigma \equiv \Sigma'$) if one can be obtained from the other with **re-signings**. (equivalently: changing signs along an edge-cut)




Signed graphs: definitions

Signature Σ of graph G : assignment of $+$ or $-$ sign to each edge of G .
 Σ : set of $-$ edges.





$$\Sigma = \{12, 34\}$$

Re-signing operation at v : switch sign of each edge incident to v




Signatures Σ, Σ' are **equivalent** ($\Sigma \equiv \Sigma'$) if one can be obtained from the other with **re-signings**. (equivalently: changing signs along an edge-cut)




Signed graphs: definitions

Signature Σ of graph G : assignment of $+$ or $-$ sign to each edge of G .
 Σ : set of $-$ edges.





$$\Sigma = \{12, 34\}$$

Re-signing operation at v : switch sign of each edge incident to v




Signatures Σ, Σ' are **equivalent** ($\Sigma \equiv \Sigma'$) if one can be obtained from the other with **re-signings**. (equivalently: changing signs along an edge-cut)




Signed graphs: definitions

Signature Σ of graph G : assignment of $+$ or $-$ sign to each edge of G .
 Σ : set of $-$ edges.






$$\Sigma = \{12, 34\}$$

Re-signing operation at v : switch sign of each edge incident to v




Signatures Σ, Σ' are **equivalent** ($\Sigma \equiv \Sigma'$) if one can be obtained from the other with **re-signings**. (equivalently: changing signs along an edge-cut)




Signed graphs: definitions

Signature Σ of graph G : assignment of $+$ or $-$ sign to each edge of G .
 Σ : set of $-$ edges.



$$\Sigma = \{12, 34\}$$

Re-signing operation at v : switch sign of each edge incident to v



Signatures Σ, Σ' are **equivalent** ($\Sigma \equiv \Sigma'$) if one can be obtained from the other with **re-signings**. (equivalently: changing signs along an edge-cut)



Signed graph: Graph G with an equivalence class \mathcal{C} of signatures.

Notation: (G, Σ) with any $\Sigma \in \mathcal{C}$.

Signed graphs: (un)balanced cycles

Definition - Unbalanced cycle



Cycle with an odd number of negative edges.



Signed graphs: (un)balanced cycles

Definition - Unbalanced cycle

Cycle with an odd number of negative edges.





Remark

Re-signing always preserves the balance of a cycle.

Signed graphs: (un)balanced cycles

Definition - Unbalanced cycle

Cycle with an odd number of negative edges.



Remark

Re-signing always preserves the balance of a cycle.

Theorem (Zaslavsky, 1982)

Two signatures are equivalent if and only if they induce the same set of unbalanced cycles.

Why signed graphs?

Introduced by Harary (1953): notion of **balanced** signed graphs (each cycle is balanced)

→ **Social psychology**: “like” and “dislike” relations in a social network. Balanced networks are socially stable. (Cartwright and Harary, 1956)

Why signed graphs?

Introduced by Harary (1953): notion of **balanced** signed graphs (each cycle is balanced)

→ **Social psychology:** “like” and “dislike” relations in a social network. Balanced networks are socially stable. (Cartwright and Harary, 1956)

→ **Graph theory**

Conjecture (Hadwiger, 1943)

If G has no K_k as a minor, $\chi(G) \leq k - 1$.

Very difficult; proved up to $k = 6$.

Why signed graphs?

Introduced by Harary (1953): notion of **balanced** signed graphs (each cycle is balanced)

→ **Social psychology**: “like” and “dislike” relations in a social network. Balanced networks are socially stable. (Cartwright and Harary, 1956)

→ **Graph theory**

Conjecture (Hadwiger, 1943)

If G has no K_k as a minor, $\chi(G) \leq k - 1$.

Very difficult; proved up to $k = 6$.

Conjecture (“Odd Hadwiger” - Seymour; Gerards, 1993)

If $(G, E(G))$ has no $(K_k, E(K_k))$ as a minor, $\chi(G) \leq k - 1$.

Extends the previous one; proved up to $k = 5$.

Signed graph homomorphisms

Definition - Signed graph homomorphism from (G, Σ_G) to (H, Σ_H)

Homomorphism $f : G \rightarrow H$ such that there exists $\Sigma'_G \equiv \Sigma_G$ for which the signs are preserved with respect to Σ'_G, Σ_H .

Recently introduced by Naserasr, Rollova and Sopena (2012)

Signed graph homomorphisms

Definition - Signed graph homomorphism from (G, Σ_G) to (H, Σ_H)

Homomorphism $f : G \rightarrow H$ such that there exists $\Sigma'_G \equiv \Sigma_G$ for which the signs are preserved with respect to Σ'_G, Σ_H .

Recently introduced by Naserasr, Rollova and Sopena (2012)




Target: $(K_2, E(K_2))$

Signed graph homomorphisms

Definition - Signed graph homomorphism from (G, Σ_G) to (H, Σ_H)

Homomorphism $f : G \rightarrow H$ such that there exists $\Sigma'_G \equiv \Sigma_G$ for which the signs are preserved with respect to Σ'_G, Σ_H .

Recently introduced by Naserasr, Rollova and Sopena (2012)




Target: $(K_2, E(K_2))$

Signed graph homomorphisms

Definition - Signed graph homomorphism from (G, Σ_G) to (H, Σ_H)

Homomorphism $f : G \rightarrow H$ such that there exists $\Sigma'_G \equiv \Sigma_G$ for which the signs are preserved with respect to Σ'_G, Σ_H .

Recently introduced by Naserasr, Rollova and Sopena (2012)




Target: $(K_2, E(K_2))$

Signed graph homomorphisms



Definition - Signed graph homomorphism from (G, Σ_G) to (H, Σ_H)

Homomorphism $f : G \rightarrow H$ such that there exists $\Sigma'_G \equiv \Sigma_G$ for which the signs are preserved with respect to Σ'_G, Σ_H .

Recently introduced by Naserasr, Rollova and Sopena (2012)



Target: $(K_2, E(K_2))$




Target:
 UC_4

Signed graph homomorphisms



Definition - Signed graph homomorphism from (G, Σ_G) to (H, Σ_H)

Homomorphism $f : G \rightarrow H$ such that there exists $\Sigma'_G \equiv \Sigma_G$ for which the signs are preserved with respect to Σ'_G, Σ_H .

Recently introduced by Naserasr, Rollova and Sopena (2012)



Target: $(K_2, E(K_2))$




Target:
 UC_4

Signed graph homomorphisms



Definition - Signed graph homomorphism from (G, Σ_G) to (H, Σ_H)

Homomorphism $f : G \rightarrow H$ such that there exists $\Sigma'_G \equiv \Sigma_G$ for which the signs are preserved with respect to Σ'_G, Σ_H .

Recently introduced by Naserasr, Rollova and Sopena (2012)



Target: $(K_2, E(K_2))$




Target:
 UC_4

Signed graph homomorphisms



Definition - Signed graph homomorphism from (G, Σ_G) to (H, Σ_H)

Homomorphism $f : G \rightarrow H$ such that there exists $\Sigma'_G \equiv \Sigma_G$ for which the signs are preserved with respect to Σ'_G, Σ_H .

Recently introduced by Naserasr, Rollova and Sopena (2012)



Target: $(K_2, E(K_2))$




Target:
 UC_4

Signed graph homomorphisms



Definition - Signed graph homomorphism from (G, Σ_G) to (H, Σ_H)

Homomorphism $f : G \rightarrow H$ such that there exists $\Sigma'_G \equiv \Sigma_G$ for which the signs are preserved with respect to Σ'_G, Σ_H .

Recently introduced by Naserasr, Rollova and Sopena (2012)



Target: $(K_2, E(K_2))$




Target:
 UC_4

Signed graph homomorphisms



Definition - Signed graph homomorphism from (G, Σ_G) to (H, Σ_H)

Homomorphism $f : G \rightarrow H$ such that there exists $\Sigma'_G \equiv \Sigma_G$ for which the signs are preserved with respect to Σ'_G, Σ_H .

Recently introduced by Naserasr, Rollova and Sopena (2012)



Target: $(K_2, E(K_2))$




Target:
 UC_4

Signed graph homomorphisms






Definition - Signed graph homomorphism from (G, Σ_G) to (H, Σ_H)

Homomorphism $f : G \rightarrow H$ such that there exists $\Sigma'_G \equiv \Sigma_G$ for which the signs are preserved with respect to Σ'_G, Σ_H .

Recently introduced by Naserasr, Rollova and Sopena (2012)



Target: $(K_2, E(K_2))$



Target:
 UC_4

Complexity: classical homomorphisms

Definition - H -COLOURING

INSTANCE: A graph G .

QUESTION: does $G \rightarrow H$?

Complexity: classical homomorphisms

Definition - H -COLOURING

INSTANCE: A graph G .

QUESTION: does $G \rightarrow H$?

Theorem (Karp, 1972)

K_3 -COLOURING is NP-complete.

Complexity: classical homomorphisms

Definition - H -COLOURING

INSTANCE: A graph G .

QUESTION: does $G \rightarrow H$?

Theorem (Hell, Nešetřil, 1990)

H -COLOURING is NP-complete for every non-bipartite graph H .
Polynomial (trivial) if H is bipartite or has a loop.

Complexity: classical homomorphisms

Definition - H -COLOURING

INSTANCE: A graph G .

QUESTION: does $G \rightarrow H$?

Theorem (Hell, Nešetřil, 1990)

H -COLOURING is NP-complete for every non-bipartite graph H .
Polynomial (trivial) if H is bipartite or has a loop.

Conjecture (Feder-Vardi, 1998: Dichotomy conjecture)

For every **digraph** D , D -COLOURING is either NP-complete or polynomial-time solvable.

(Equivalent to dichotomy for CSP and MMSNP — tough conjecture!)

Complexity: questions for signed graphs

Definition - (H, Σ_H) -COLOURING

INSTANCE: A signed graph (G, Σ) .

QUESTION: does $(G, \Sigma) \rightarrow (H, \Sigma_H)$?

Complexity: questions for signed graphs

Definition - (H, Σ_H) -COLOURING

INSTANCE: A signed graph (G, Σ) .

QUESTION: does $(G, \Sigma) \rightarrow (H, \Sigma_H)$?

Remark

- checking if $\Sigma \equiv \Sigma'$: polynomial
 - $(G, \Sigma) \rightarrow (H, \emptyset)$ IFF $G \rightarrow H$ and $\Sigma \equiv \emptyset$.
 - $(G, \Sigma) \rightarrow (H, E(H))$ IFF $G \rightarrow H$ and $\Sigma \equiv E(G)$.
- If $\Sigma_H \equiv \emptyset$ or $\Sigma_H \equiv E(H)$, (H, Σ_H) -COLOURING has same complexity as H -COLOURING.

Complexity: questions for signed graphs

Definition - (H, Σ_H) -COLOURING

INSTANCE: A signed graph (G, Σ) .

QUESTION: does $(G, \Sigma) \rightarrow (H, \Sigma_H)$?

Remark

- checking if $\Sigma \equiv \Sigma'$: polynomial
 - $(G, \Sigma) \rightarrow (H, \emptyset)$ IFF $G \rightarrow H$ and $\Sigma \equiv \emptyset$.
 - $(G, \Sigma) \rightarrow (H, E(H))$ IFF $G \rightarrow H$ and $\Sigma \equiv E(G)$.
- If $\Sigma_H \equiv \emptyset$ or $\Sigma_H \equiv E(H)$, (H, Σ_H) -COLOURING has same complexity as H -COLOURING.


Polynomial cases:

- H bipartite, $\Sigma_H \equiv \emptyset \equiv E(H)$
- H has one vertex with both $+$ loop and $-$ loop
- H has a loop and $\Sigma_H \equiv \emptyset$ or $\Sigma_H \equiv E(H)$
- H is bipartite and contains a multi-edge ($+$ and $-$)

Reduction from classical H -COLOURING

Theorem (Brewster, F., Hell, 2013+)


$$(G, E(G)) \rightarrow (H, \Sigma) \text{ IFF } G \rightarrow ((H, \Sigma) \times K_2^{R-})^U$$



Reduction from classical H -COLOURING

Theorem (Brewster, F., Hell, 2013+)


$$(G, E(G)) \rightarrow (H, \Sigma) \text{ IFF } G \rightarrow ((H, \Sigma) \times K_2^{R-})^U$$



Reduction from classical H -COLOURING

Theorem (Brewster, F., Hell, 2013+)

$$(G, E(G)) \rightarrow (H, \Sigma) \text{ IFF } G \rightarrow ((H, \Sigma) \times K_2^{R-})^U$$



Corollary


If (H, Σ) has an **unbalanced odd** cycle, then (H, Σ) -COLOURING is NP-complete.

(symmetric result holds when (H, Σ) has *balanced* odd cycle)

Reduction from classical H -COLOURING

Theorem (Brewster, F., Hell, 2013+)

$$(G, E(G)) \rightarrow (H, \Sigma) \text{ IFF } G \rightarrow ((H, \Sigma) \times K_2^{R-})^U$$



Corollary


If (H, Σ) has an **unbalanced odd** cycle, then (H, Σ) -COLOURING is NP-complete.

(symmetric result holds when (H, Σ) has *balanced* odd cycle)

Reduction from classical H -COLOURING

Theorem (Brewster, F., Hell, 2013+)

$$(G, E(G)) \rightarrow (H, \Sigma) \text{ IFF } G \rightarrow ((H, \Sigma) \times K_2^{R-})^U$$



Corollary


If (H, Σ) has an **unbalanced odd** cycle, then (H, Σ) -COLOURING is NP-complete.

(symmetric result holds when (H, Σ) has *balanced* odd cycle)

Reduction from classical H -COLOURING

Theorem (Brewster, F., Hell, 2013+)

$$(G, E(G)) \rightarrow (H, \Sigma) \text{ IFF } G \rightarrow ((H, \Sigma) \times K_2^{R-})^U$$



Corollary


If (H, Σ) has an **unbalanced odd** cycle, then (H, Σ) -COLOURING is NP-complete.

(symmetric result holds when (H, Σ) has *balanced* odd cycle)

Reduction from classical H -COLOURING

Theorem (Brewster, F., Hell, 2013+)

$$(G, E(G)) \rightarrow (H, \Sigma) \text{ IFF } G \rightarrow ((H, \Sigma) \times K_2^{R-})^U$$



Corollary

If (H, Σ) has an **unbalanced odd** cycle, then (H, Σ) -COLOURING is NP-complete.

(symmetric result holds when (H, Σ) has *balanced* odd cycle)

Reduction from NAE-3SAT

Definition - UC_{2k} -COLOURING

INSTANCE: A (bipartite) signed graph (G, Σ) .

QUESTION: does $(G, \Sigma) \rightarrow UC_{2k}$?

Reduction from NAE-3SAT

Definition - UC_{2k} -COLOURING

INSTANCE: A (bipartite) signed graph (G, Σ) .

QUESTION: does $(G, \Sigma) \rightarrow UC_{2k}$?

Theorem (F., Naserasr, 2012+)

UC_{2k} -COLOURING is NP-complete for every $k \geq 2$.

Reduction from NAE-3SAT

Definition - UC_{2k} -COLOURING

INSTANCE: A (bipartite) signed graph (G, Σ) .

QUESTION: does $(G, \Sigma) \rightarrow UC_{2k}$?

Theorem (F., Naserasr, 2012+)



UC_{2k} -COLOURING is NP-complete for every $k \geq 2$.

Definition - MONOTONE NOT-ALL-EQUAL-3SAT

INSTANCE: A set of clauses of 3 Boolean variables from set X .

QUESTION: Is there a truth assignment $X \rightarrow \{0, 1\}$ s.t. each clause has variables with different values?


NAE-3SAT \leq_R UC_4 -COLOURING: clause gadget



Construction of $G(F)$: one clause gadget per clause of F .
All vertices with same labels (c or x_i) identified with each other.

Main idea: In a mapping, re-signing at $x_i \iff x_i = \text{TRUE}$

NAE-3SAT \leq_R UC_{2k} -COLOURING: clause gadget



(where P_k has length $k - 1$)

CSPs, signed CSPs

Constraint Satisfaction Problem (CSP) for relational system

$T = (X_T, V_T)$: domain X_T , set V relations R_1, \dots, R_k of arity a_1, \dots, a_k with $R_i \subseteq X^{a_i}$ (vocabulary).

Definition - T -CSP, $T = (X_T, V_T)$

INSTANCE: domain X with vocabulary $V \cong V_T$.

QUESTION: does $(X, V) \rightarrow T$?

Constraint Satisfaction Problem (CSP) for relational system

$T = (X_T, V_T)$: domain X_T , set V relations R_1, \dots, R_k of arity a_1, \dots, a_k with $R_i \subseteq X^{a_i}$ (vocabulary).

Definition - T -CSP, $T = (X_T, V_T)$

INSTANCE: domain X with vocabulary $V \cong V_T$.

QUESTION: does $(X, V) \rightarrow T$?

Examples:

- (Di)graph homomorphism to D : $X_T = V(D)$, V_T is one binary (non-)symmetric relation.
- 3SAT: $X_T = \{0, 1\}$, V_T : one ternary relation with all triples except 000.

CSPs, signed CSPs

Constraint Satisfaction Problem (CSP) for relational system

$T = (X_T, V_T)$: domain X_T , set V relations R_1, \dots, R_k of arity a_1, \dots, a_k with $R_i \subseteq X^{a_i}$ (vocabulary).

Definition - T -CSP, $T = (X_T, V_T)$

INSTANCE: domain X with vocabulary $V \cong V_T$.

QUESTION: does $(X, V) \rightarrow T$?

Conjecture (Feder-Vardi, 1998: Dichotomy conjecture)

For every T , T -CSP is either NP-complete or polynomial-time.

CSPs, signed CSPs

Constraint Satisfaction Problem (CSP) for relational system

$T = (X_T, V_T)$: domain X_T , set V relations R_1, \dots, R_k of arity a_1, \dots, a_k with $R_i \subseteq X^{a_i}$ (vocabulary).

Definition - T -CSP, $T = (X_T, V_T)$

INSTANCE: domain X with vocabulary $V \cong V_T$.

QUESTION: does $(X, V) \rightarrow T$?

Conjecture (Feder-Vardi, 1998: Dichotomy conjecture)

For every T , T -CSP is either NP-complete or polynomial-time.

Signed CSP: $+$ and $-$ tuples, re-signing allowed.

Proposition

Dichotomy for CSP \iff Dichotomy for signed CSP

- Prove dichotomy for (H, Σ) -COLOURING.
→ remaining cases: H bipartite or has both kinds of loops
- Study signed CSPs