

Examining the Interplay of Interoceptive and Exteroceptive Influences on Perception

Fan Gao¹

¹ University of Chicago

Author Note

For this apa 6 style pdf document, I used Tinytex
[<https://github.com/rstudio/tinytex-releases>].

The authors made the following contributions. Fan Gao: Data collection, Writing -
Original Draft Preparation, Writing - Review & Editing.

Correspondence concerning this article should be addressed to Fan Gao, University of
Chicago. E-mail: fgao38@uchicago.com

Abstract

This study explores the interplay between interoceptive stimuli, specifically the heartbeat, and exteroceptive visual perception. While existing research showed the impact of external stimuli on bodily functions, this investigation delves into the reverse relationship—whether interoceptive stimuli, often inaccessible to conscious awareness, influence visual perception. Recent studies have demonstrated continuous electrical activity in the heart and gastrointestinal tract, sending messages to the brain and shaping cognition (Azzalini et al., 2019). Emotional states derived from interoception can alter our perception of the external world (Smith et al., 2017), exemplified by perceiving neutral stimuli as threatening during anxious states. Despite prior investigations, the visual aspect of interoceptive-exteroceptive interaction remains understudied. This research addresses this gap by employing a binocular rivalry paradigm (Carmel et al., 2010), presenting competing visual stimuli with one synchronized to participants' real-time heartbeat. The objective is to examine if this synchronization influences the prioritization of visual stimuli in conscious awareness. Building on earlier findings, we hypothesize that stimuli matching the participant's heartbeat will dominate the visual field for longer durations. Importantly, we anticipate this effect to persist irrespective of participants' conscious awareness of their heartbeat sensations. This study contributes to a more comprehensive understanding of the dynamic processes underlying perception and consciousness, particularly in the context of visual bistable perceptual switching. The findings hold potential implications for elucidating the mechanisms through which the brain resolves perceptual ambiguities, shedding light on fundamental aspects of human perception.

Keywords: ECG,Binocular-rivalry-paradigm,heart-rate,vision

Word count: X

Examining the Interplay of Interoceptive and Exteroceptive Influences on Perception

Introduction

Though we are unconscious of most bodily sensations (e.g. immune system), in a place where the internal (i.e. self) and external (i.e. physical world) interact, interoceptive stimuli—the sensation that arises from an internal organ (e.g. heartbeat), have been found to yield an unexpected influence over how we see and sense the world (i.e. exteroceptive stimuli).


A substantial prior study has been dedicated to exploring how external stimuli affect our body and brain. For example, intentionally observing and recognizing external stimuli typically results in a deceleration of the heart rate, referred to as “bradycardia of attention” (J. I. Lacey et al., 1963). Such an effect is further examined in a follow-up study that showed subjects’ heart rate decreased following a ready signal (B. C. Lacey & Lacey, 1978). These findings have provided us with a novel understanding of how exteroceptive stimuli (e.g. a ready signal at a traffic light) influence our interoceptive stimuli (e.g. heart rate), but also raise the interesting question about the reverse effect: could interoceptive stimuli have an influence on exteroceptive stimuli? The question may seem counterintuitive at first since most of the interoceptive stimuli within one’s self are not accessible (e.g. immune system, heartbeat). For example, studies have suggested that only a quarter of the participants can perceive and judge their heart rate that is closely synchronized with external stimuli above chance (Brener & Ring, 2016). How can these interoceptive stimuli affect our perception of the world if we, for the most of time, do not have conscious access to them? Yet, recent research has shed light on this question. For instance, the heart rate and gastrointestinal tract (GI) are shown to be continuously producing electrical activity, thus sending messages to the brain, and eventually altering our perception and cognition (Azzalini et al., 2019; Monti et al., 2022). In addition, it is suggested that interoceptive stimuli may play a significant role in shaping emotions and cognition, and this process is derived from a low-level function—homeostatic regulation (Smith et al., 2017). These emotional states can,

in turn, affect how we perceive the world. For instance, we may perceive a neutral stimulus as threatening when we are in an anxious state. One study investigating the subjective experience of body ownership (EBO) presented even more compelling evidence of interaction between interoceptive and exteroceptive stimuli by showing that during the induction of a “fake” rubber hand, participants exhibited an increased sense of EBO if the heartbeat were synchronized with the color change of the rubber hand (Suzuki et al., 2013). These similar embodiment effect were also observed in breathing (Monti et al., 2020). Another study delves into examining how activation of certain cortical areas impacts subjects’ hits and misses on a visual signal detection task: participants were asked to identify whether or not they saw a faint annulus; the study showed that the activation of ventromedial prefrontal cortex bilaterally (vACC-vmPFC), the site known for receiving cardiac inputs, were more likely to have participants consciously perceive the faint annulus (Park et al., 2014). Recently, the impact of cardiac activity on exteroceptive perception gained significant attention, particularly fueled by recent theories emphasizing the crucial role of interoception in shaping the subjective sense of self (Park et al., 2014; Seth & Tsakiris, 2018). To our knowledge, despite these prior investigations on interoceptive and exteroceptive interaction, there is limited research that closely examines this effect visually. Also, there is still a notable gap in the existing literature, particularly in the context of visual bistable perceptual switching. Visual bistable perceptual switching refers to presenting participants with two visual stimuli, each of which dominates the visual field for a short period of time. This is usually achieved by using the binocular rivalry paradigm (Carmel et al., 2010). While prior studies have primarily investigated the realm of detection thresholds (A binary response: whether the signal or not), it is important to study this effect more comprehensively in a bistable perception. When perception oscillates between two ambiguous stimuli, it suggests dynamic processes at play in the brain. Understanding the mechanism of how the brain suppresses these perceptual ambiguities can shed light on fundamental aspects of perception and consciousness. To fill in the gap, our research plans to use a binocular rivalry paradigm

(Carmel et al., 2010), where one of two competing visual stimuli will be synchronized with the subjects' heartbeat (i.e. electrocardiogram ECG signals) in real-time. Our goal is to investigate whether the synchronization of interoceptive stimuli (i.e. heartbeat) will influence the prioritization of the visual stimuli in conscious awareness. Based on earlier studies that examined the effect of interoceptive stimuli on the brain (Azzalini et al., 2019) and homeostasis regulation (Smith et al., 2017), we hypothesized to find that the visual stimulus that matched with the participant's real-time heartbeat should overall dominate the visual field longer than the stimulus that was not synchronized. In addition, we also expect to see that this effect is not dependent on participants' conscious awareness of their heartbeat sensations.


Methods

As written in my authornote, my role in this study include data collection, writing the original draft preparation, as well as review and editing. Our experiment is going to be divided into two parts. In the first section, we are planning to use a binocular rivalry paradigm – presenting different visual stimuli, one to each eye of the participant; because the brain cannot process two visual stimuli simultaneously, one visual stimulus will dominate the other visual stimulus, see Figure 1.



106

107 The idea is to synchronize one of the visual stimuli with the participant's heartbeat
108 (measured by using an electrocardiogram ECG) in
109 real-time; the synchronization of the heartbeat and visual stimulus is randomized, see Figure 2.



110

111 Participants are not going to be told that one of the stimuli was synchronized with their
 112 real-time ECG; the Participants will identify which visual stimulus they are currently
 113 viewing by pressing the left (red) and right (blue) arrow keys. In the second section, we are
 114 going to measure whether the participants could judge the external stimulus that is
 115 synchronized with their own heartbeat correctly. This will be done by presenting two pulsing
 116 circles, one synchronizes with the participant's ECG (immediately followed at the R peak)
 117 and the other one does not (followed later after the R peak).

118 Participants

119 We aim to collect 60 undergraduate students taking Psychology courses at the
 120 University of Chicago. We are going to recruit participants through an online platform
 121 named SONA (Psychological and Brain Science Research System). Participants will need to

have normal color vision and see well without glasses, as well as consent to participate in our study. Our participants' sample may not be representative since our sample consists of only college students, specifically students who are taking introductory Psychology courses. The introductory Psychology courses include a diverse population of students with different majors and backgrounds, but it is biased toward college and well-educated students at University of Chicago. However, as mentioned above, we do not expect that our results will vary significantly across races and genders since this effect is mostly driven by biological factors within the body. We are going to send our study protocol to the University of Chicago institutional review board for approval.

Material

Electrocardiogram (ECG) signals were recorded with a TMSi SAGA amplifier (TMSi, Netherlands) at a sampling rate of 100 Hz. The ECG data, accessible in Python (refer to Data and Code Availability), underwent processing through the Lab Streaming Layer (LSL, labstreaminglayer.org). To enhance signal quality, the ECG data was subjected to bandpass filtering within the frequency range of 5-15 Hz. The detection of R-peaks was accomplished utilizing the Pan-Thompkins algorithm (Pan & Tompkins, 1985), adapted from a pre-existing implementation for compatibility with LabGraph (Michał Sznajder & Łukowska, 2017).

Data analysis

We used R [Version 4.2.2; R Core Team (2022)] for all our analyses.

Results

The mean duration of synchronous stimulus is 2.45, and the median is 1.96; the mean duration of asynchronous stimulus is 2.39, and the median is 1.91

The results of the generalized linear mixed-effects model (glmer) are presented in (Table 1). The glmer revealed significant effects for the dominant variable (Estimate=0.02,

t=2.64, $p<0.05$). This suggests that on average, the duration is expected to be increase by 0.02 seconds for synchronous stimulus compared to asynchronous stimulus.

Discussion

Although the research field has shifted attention to study interoceptive stimuli on exteroceptive stimuli, there are a limited number of studies that examined this effect in vision. In addition, previous studies tend to focus on investigating this effect with a detection threshold paradigm (A binary response: whether the signal or not), it is unclear the mechanism that the brain uses to suppress perceptual ambiguity.

The results of our study extend beyond a simple examination of interoceptive stimuli of exteroceptive stimuli. Understanding the influential effect of our unconscious internal sensations on our conscious perception has a broader meaning across various aspects of human life and the scientific field. An enhanced understanding of interoceptive cues could facilitate new and more effective ways of treating mental disorders such as depression and anxiety. For instance, if the accelerated heartbeat is causing a negative emotional state and thus leading to symptoms of depression and anxiety, therapies could develop new interventions that aim to help the patients become more aware of the interoceptive stimuli, making individuals manage the emotional impact led by interoceptive stimuli, thereby alleviate the depression and anxiety symptoms. Furthermore, interoceptive stimuli such as heartbeat, accompanied with other diagnostic criteria can be used as a sign to predict certain mental disorders. For example, anxiety is marked by excessive worries and fears that tend to trigger a fight or flight response. The heartbeat is likely to get accelerated when an individual is deciding whether to fight or flight. Therefore, internal sensations such as heartbeat, can also serve as a complementary diagnostic information to improve current therapeutic interventions. The present study was conducted within a controlled laboratory environment, which may limit the extent to which the findings can be extrapolated to real-world settings. While the controlled environment allowed for precise manipulation of

172 variables, the ecological validity of the results may be constrained. Future research could
173 consider incorporating more ecologically valid scenarios to enhance the external validity of
174 the study. Another potential limitation of this study is the specificity of the sample
175 population. Participants in this study were students from University of Chicago, and caution
176 should be exercised when generalizing the findings to a broader and more diverse population.
177 Future research could explore the generalizability of the observed effects across different
178 demographic groups to provide a more comprehensive understanding of the phenomenon
179 under investigation.

References

- Azzalini, D., Rebollo, I., & Tallon-Baudry, C. (2019). Visceral Signals Shape Brain Dynamics and Cognition. *Trends in Cognitive Sciences*, 23(6), 488–509. <https://doi.org/10.1016/j.tics.2019.03.007>
- Brener, J., & Ring, C. (2016). Towards a psychophysics of interoceptive processes: The measurement of heartbeat detection. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371(1708), 20160015. <https://doi.org/10.1098/rstb.2016.0015>
- Carmel, D., Arcaro, M., Kastner, S., & Hasson, U. (2010). How to Create and Use Binocular Rivalry. *Journal of Visualized Experiments*, 45, 2030. <https://doi.org/10.3791/2030>
- Lacey, B. C., & Lacey, J. I. (1978). Two-way communication between the heart and the brain: Significance of time within the cardiac cycle. *American Psychologist*, 33(2), 99–113. <https://doi.org/10.1037/0003-066X.33.2.99>
- Lacey, J. I., Kagan, J., Lacey, B. C., & Moss, H. A. (1963). *The visceral level: Situational determinants and behavioral correlates of autonomic response patterns*. 161–196.
- Michał Sznajder, & Łukowska, M. (2017). *Python Online and Offline ECG QRS Detector based on the Pan-Tomkins algorithm*. Zenodo. <https://doi.org/10.5281/ZENODO.583770>
- Monti, A., Porciello, G., Panasiti, M. S., & Aglioti, S. M. (2022). Gut markers of bodily self-consciousness in men. *Iscience*, 25(10).
- Monti, A., Porciello, G., Tieri, G., & Aglioti, S. M. (2020). The “embreathment” illusion highlights the role of breathing in corporeal awareness. *Journal of Neurophysiology*, 123(1), 420–427. <https://doi.org/10.1152/jn.00617.2019>
- Pan, J., & Tompkins, W. J. (1985). A Real-Time QRS Detection Algorithm. *IEEE Transactions on Biomedical Engineering*, BME-32(3), 230–236. <https://doi.org/10.1109/TBME.1985.325532>
- Park, H.-D., Correia, S., Ducorps, A., & Tallon-Baudry, C. (2014). Spontaneous fluctuations in neural responses to heartbeats predict visual detection. *Nature Neuroscience*, 17(4), 612–618. <https://doi.org/10.1038/nn.3671>

- 207 R Core Team. (2022). *R: A Language and Environment for Statistical Computing*. R
208 Foundation for Statistical Computing. <https://www.R-project.org/>
- 209 Seth, A. K., & Tsakiris, M. (2018). Being a Beast Machine: The Somatic Basis of Selfhood.
210 *Trends in Cognitive Sciences*, 22(11), 969–981. <https://doi.org/10.1016/j.tics.2018.08.008>
- 211 Smith, R., Thayer, J. F., Khalsa, S. S., & Lane, R. D. (2017). The hierarchical basis of
212 neurovisceral integration. *Neuroscience & Biobehavioral Reviews*, 75, 274–296.
213 <https://doi.org/10.1016/j.neubiorev.2017.02.003>
- 214 Suzuki, K., Garfinkel, S. N., Critchley, H. D., & Seth, A. K. (2013). Multisensory integration
215 across exteroceptive and interoceptive domains modulates self-experience in the
216 rubber-hand illusion. *Neuropsychologia*, 51(13), 2909–2917.
217 <https://doi.org/10.1016/j.neuropsychologia.2013.08.014>

Table 1

Generalized Linear Mixed-Effects Models subject random intercept

Effect	Estimate and SE		Model fit	
	Estimate	SEr	t.value	p.value
Intercept	0.927540	0.045024	20.601	< 2e-16
Dominant	0.019123	0.007245	2.639	0.00831**

Note:

*p < 0.05, **p < 0.01, ***p < 0.001




Figure 1. Average Duration for Syn and Asyn




Figure 2. Histogram for subtracted difference