Ch1 Covid-19

COVID-19 **pandemic** is the disease caused by a new **coronavirus** called SARS-CoV-2. This new virus was known to the public on 31 December 2019, following a report of a cluster of cases of '**viral pneumonia**' in Wuhan, China.
COVID-19 **流行病**是由一種名為 SARS-CoV-2 的新型**冠狀病毒**所引起的疾病 。這種新病毒在 2019 年 12 月 31 日中國武漢通報了一群「**病毒的 肺炎**」病例後為大眾所知 。

The most common **symptoms** of COVID-19 include **fever**, dry **cough**, and **fatigue**. Other **symptoms** that are less common and may affect some people include: loss of taste or **smell**, **nasal congestion**, **conjunctivitis** (also known as red eyes), **sore** throat, **headache**, **muscle** or **joint** pain, **diarrhea**, different types of skin **rash**, **chills** or **dizziness**, **nausea** or **vomiting**.
COVID-19 最常見的**癥狀**包括**發燒**、乾**咳**和**疲勞** 。其他較不常見且可能影響某些人的**癥狀**包括：味覺或**嗅覺**喪失、**鼻的 阻塞**（鼻塞）、**結膜炎**（也稱為紅眼症）、**痛的**喉嚨（喉嚨痛）、**頭痛**、**肌肉**或**關節**疼痛、**腹瀉**、不同類型的**皮疹**、**寒意**或**眩暈**、**噁心**或**嘔吐** 。

Symptoms of **severe** COVID-19 disease include **shortness of breath**, loss of **appetite**, **confusion**, persistent pain or pressure in the **chest**, and high temperature. Other less common **symptoms** include **irritability**, **confusion**, reduced **consciousness**, **anxiety**, **depression**, and sleep **disorders**. More **severe** and rare **neurological complications** include **strokes**, brain **inflammation**, **delirium** and nerve damage.
非常嚴重的 COVID-19 疾病**癥狀**包括**呼吸急促**、**食慾**不振、**困惑**（意識不清）、**胸部**持續疼痛或有壓迫感，以及高溫 。其他較不常見的**癥狀**包括**易怒性情**、**困惑**、**意識**降低、**焦慮**、**憂鬱**和睡眠**混亂**（障礙） 。更**非常嚴重的**和罕見的**神經系統的 併發症**包括**中風**、腦**炎**、**神志不清**和神經損傷 。

People of all **ages** who experience **fever** and/or **cough** associated with difficulty breathing or **shortness of breath**, **chest** pain or pressure, or loss of speech or movement should seek medical care immediately. Among those who develop **symptoms**, about 80% **recover** from the disease without needing hospital **treatment**. About 15% become seriously ill and require **oxygen** and 5% become critically ill and need intensive care.
所有**年齡**層的人，如果出現**發燒**和/或**咳嗽**，並伴隨呼吸困難或**呼吸急促**、**胸部**疼痛或壓迫感，或喪失言語或運動能力，應立即就醫 。在出現**癥狀**的患者中，約 80% 的人無需住院**治療**即可從疾病中**康復** 。約 15% 的人病情嚴重需要**氧氣**，5% 的人病情危急需要重症監護 。

Complications leading to death may include **respiratory failure**, **acute respiratory** distress **syndrome** (ARDS), **sepsis** and **septic shock**, **thromboembolism**, and/or multi-organ failure, including injury of the heart, liver or **kidneys**. People aged 60 years and over, and those with **underlying** medical problems such as high blood pressure, heart and **lung** problems, **diabetes**, **obesity** or cancer, are at higher risk of

developing serious illness. However, anyone can get **infected** with COVID-19 and become seriously ill or die. Some people who had COVID-19, whether they have needed **hospitalization** or not, continue to experience **symptoms**, including **fatigue**, **respiratory** and **neurological symptoms**.
導致死亡的**併發症**可能包括**呼吸衰竭**、**十分嚴重的**（急性）**呼吸的**窘迫**症候群**（ARDS）、**敗血症**和**感染性休克**、**血栓栓塞**，和/或多器官衰竭，包括心臟、肝臟或**腎臟**的損傷 。60 歲及以上的人，以及患有**潛在的**醫療問題（如高血壓、心**肺**問題、**糖尿病**、**肥胖**或癌症）的人，發展為重症的風險較高 。然而，任何人都可能**感染** COVID-19 並變成重症或死亡 。一些曾患有 COVID-19 的人，無論是否需要**送…住院治療**，都會繼續出現**癥狀**，包括**疲勞**、**呼吸的**和**神經系統的 癥狀** 。

Viruses constantly change through **mutation** and sometimes these **mutations** result in a new **variant** of the virus. Some **variants emerge** and disappear while others persist. New **variants** will continue to **emerge**. Five SARS-CoV-2 **variants** have been **designated** as **variants** of concern by the WHO: the Alpha, Beta, Gamma, Delta, and Omicron **variants**.
病毒透過**突變**不斷變化，有時這些**突變**會導致病毒的新**變體** 。一些**變體 出現**然後消失，而其他則持續存在 。新的**變體**將繼續**出現** 。WHO 已**指定**五種 SARS-CoV-2 **變體**為高關注變異株：Alpha、Beta、Gamma、Delta 和 Omicron **變體** 。

The Omicron **variants** are the most recent one **announced** at the end of 2021. They are believed to be most **transmissible** even though the World Health Organization (WHO), at the beginning of 2022, **claimed** that "It is not yet clear whether Omicron is more **transmissible** compared to other **variants**, including Delta" and "It is not yet clear whether **infection** with Omicron causes more **severe** disease compared to **infections** with other **variants**, including Delta."
Omicron **變體**是 2021 年底**宣佈**的最新變種 。人們認為它們最具**可傳播的**特性，儘管世界衛生組織（WHO）在 2022 年初**聲稱**「目前尚不清楚 Omicron 是否比其他**變體**（包括 Delta）更具**可傳播的**特性」，以及「目前尚不清楚 Omicron **感染**是否會比其他**變體**（包括 Delta）的**感染**導致更**非常嚴重的**疾病」 。

WHO is working with researchers and patient groups around the world to design and conduct studies on patients beyond the initial **acute** course of illness to understand the **proportion** of patients who have long term effects, how long they persist, and why they occur. These studies will be used to develop further guidance for patient care.
WHO 正與世界各地的研究人員和患者團體合作，設計並對度過最初**十分嚴重的**（急性）病程的患者進行研究，以了解產生長期影響的患者**比例**、這些影響持續多久以及發生原因 。這些研究將用於制定進一步的患者護理指南 。

Stay safe by taking some simple **precautions**, such as physical **distancing**, wearing a **mask**, especially when **distancing** cannot be maintained. Keeping rooms well **ventilated**, avoiding crowds and close **contact**, regularly cleaning your hands, and **coughing** into a **bent elbow** or **tissue**. Check local advice

where you live and work.
採取一些簡單的**預防措施**以保持安全，例如保持物理**距離**、戴**面罩**，尤其是在無法**保持距離**時 。保持房間良好**使通風**、避免人群和密切**聯絡**（接觸）、定期清潔雙手，並在**咳嗽**時用**彎曲的 肘**部或**面紙**遮擋 。查看您居住和工作地點的當地建議 。

In most situations, a **molecular** test is used to **detect** SARS-CoV-2 and **confirm infection**. Polymerase chain reaction (**PCR**) is the most commonly used **molecular** test. Medical personnel may collect **samples** from your nose or throat with a **swab**. **Molecular** tests **detect** virus in the **sample** by **amplifying viral** genetic material to detectable levels. For this reason, a **molecular** test is used to confirm an active **infection**, usually within a few days of **exposure** and around the time that **symptoms** may begin.
在大多數情況下，**分子的**檢測用於**檢測出** SARS-CoV-2 並**確認感染** 。**聚合酶連鎖反應**（**PCR**）是最常用的**分子的**檢測 。醫務人員可能會用**棉花棒**從您的鼻子或喉嚨收集**樣本** 。**分子的**檢測透過將**病毒的**遺傳物質**擴大**到可檢測水平來**檢測出 樣本**中的病毒 。因此，**分子的**檢測用於確認活動性**感染**，通常在**暴露**（接觸）後幾天內和**癥狀**可能開始時 。

Rapid **antigen** test (sometimes known as a rapid diagnostic test, or RDT) is an alternative in screening people **infected** by COVID-19. It **detects viral proteins** (known as **antigens**). In this test, **samples** are also collected from the nose or throat using a **swab**. This test is cheaper than **PCR** and will offer results more quickly, although they are normally less **reliable**. Rapid **antigen** tests perform best when there is more virus **circulating** in the community and when **sampled** from an individual during the time they are most infectious.
快速**抗原**檢測（有時稱為快速診斷檢測或 RDT）是篩查 COVID-19 **感染**者的替代方法 。它**檢測出 病毒的 蛋白質**（稱為**抗原**） 。在該測試中，也會使用**棉花棒**從鼻子或喉嚨收集**樣本** 。這項測試比 **PCR** 便宜，而且可以更快提供結果，儘管它們通常不太**可靠的** 。當社區中**傳遞**的病毒較多，且在個體最具傳染性期間進行**抽樣**時，快速**抗原**檢測的效果最佳 。

Both **isolation** and **quarantine** are implemented in **preventing** the spread of COVID-19. **Quarantine** is required for anyone who has **contacted** someone **infected** with the SARS-CoV-2 virus, whether the **infected** person has **symptoms** or not. **Quarantine** means that you remain separated from others because you have been exposed to the virus and you may be **infected**. **Quarantine** can be implemented in a designated facility or at home for a period of time (check the announcement of local government).
隔離和**隔離檢疫期**都是為了**預防** COVID-19 傳播而實施的 。任何**聯絡**（接觸）過 SARS-CoV-2 病毒**感染**者的人都需要**隔離檢疫期**，無論**感染**者是否有**癥狀** 。**隔離檢疫期**意味著您因為暴露於病毒並可能被**感染**而與他人分開 。**隔離檢疫期**可以在指定設施或家中實施一段時間（請查看當地政府的公告） 。

Isolation is adopted for people with COVID-19 **symptoms** or who have tested positive for the virus. Being in **isolation** means being separated from other people, ideally in a medically facility where you can receive clinical care. If **isolation** in a medical facility is not possible and you are not in a high risk group of

developing **severe** disease, **isolation** may also take place at home. If you are **infected** with or without **symptoms**, you should report to the local government and follow their instructions. You may need medical care if you do have **symptoms**.
隔離是針對出現 COVID-19 **癥狀**或病毒檢測呈陽性的人 。被**隔離**意味著與他人分開，理想情況下是在可以接受臨床護理的醫療機構中 。如果無法在醫療機構**隔離**，且您不屬於發展成**非常嚴重的**疾病的高危人群，也可以在家中進行**隔離** 。如果您被**感染**（無論是否有**癥狀**），都應向當地政府報告並遵循他們的指示 。如果您確實有**癥狀**，您可能需要醫療護理 。

Vaccines provide **protection** to people against virus. They save millions of lives each year. Vaccines work by training and preparing the body's natural defenses, the **immune** system, to **recognize** and fight against the viruses and **bacteria** they target. After **vaccination**, if the body is later exposed to those virus or germs, the body is immediately ready to destroy them, **preventing** illness.
疫苗為人們提供抵禦病毒的**保護** 。它們每年拯救數百萬人的生命 。疫苗的作用是訓練和準備身體的自然防禦（**免疫的**系統），以**認出**並對抗它們針對的病毒和**細菌** 。**接種疫苗**後，如果身體後來接觸到這些病毒或細菌，身體就會立即準備好消滅它們，從而**預防**疾病 。

There are several safe and effective vaccines that **prevent** people from getting seriously ill or dying from COVID-19. This is one part of managing COVID-19, in addition to the main **preventive** measures of staying at least 1 m away from others, covering a **cough** or **sneeze** in your **elbow**, frequently cleaning your hands, wearing a **mask** and avoiding poorly **ventilated** rooms or opening a window.
有幾種安全有效的疫苗可以**預防**人們因 COVID-19 而患重病或死亡 。這是管理 COVID-19 的一部分，除了主要的**防止的**（預防的）措施外，如與他人保持至少 1 公尺的距離、**咳嗽**或**打噴嚏**時用**肘**部遮擋、經常清潔雙手、戴**面罩**，並避免**使通風**不良的房間或打開窗戶 。

As of 15 November 2021, WHO has **announced** that the following vaccines against COVID-19 have met the necessary criteria for safety and **efficacy**: AstraZeneca/Oxford, Johnson and Johnson, Moderna, Pfizer/BioNTech, Sinopharm, Sinovac, COVAXIN.
截至 2021 年 11 月 15 日，WHO 已**宣佈**以下針對 COVID-19 的疫苗已達到安全性和**功效**的必要標準：AstraZeneca/Oxford、Johnson and Johnson、Moderna、Pfizer/BioNTech、Sinopharm、Sinovac、COVAXIN 。

Take whatever vaccine is available to you first, even if you have already had COVID-19. It is important to get **vaccinated** as soon as possible once it's your turn and do not wait. Approved COVID-19 vaccines provide a high degree of **protection** against getting seriously ill and dying from the disease, although no vaccine is 100% protective.
首先接種任何您可獲得的疫苗，即使您已經得過 COVID-19 。輪到您時，盡快**給…接種疫苗**非常重要，不要等待 。經批准的 COVID-19 疫苗可提供高度**保護**，防止染上該疾病成為重症和死亡，儘管沒有疫苗是 100% 具有保護力的 。

COVID-19 [1]pandemic is the disease caused by a new coronavirus called SARS-CoV-2. This new virus was known to the public on 31 December 2019, following a report of a cluster of cases of '[2]viral [3]pneumonia' in Wuhan, China.

The most common symptoms of COVID-19 include fever, dry cough, and fatigue. Other symptoms that are less common and may affect some people include:

- loss of taste or smell,
- nasal congestion,
- [4]conjunctivitis (also known as red eyes)
- sore throat,
- headache,
- muscle or joint pain,
- [5]diarrhea,
- different types of skin rash,
- chills or dizziness,
- [6]nausea or vomiting.

Symptoms of severe COVID-19 disease include shortness of breath, loss of appetite, confusion, persistent pain or pressure in the chest, and high temperature. Other less common symptoms include irritability, confusion,

reduced consciousness, anxiety, depression, and sleep disorders. More severe and rare [7]neurological complications include strokes, brain inflammation, delirium and nerve damage.

People of all ages who experience fever and/or cough associated with difficulty breathing or shortness of breath, chest pain or pressure, or loss of speech or movement should seek medical care immediately. Among those who develop symptoms, about 80% recover from the disease without needing hospital treatment. About 15% become seriously ill and require oxygen and 5% become critically ill and need intensive care.

Complications leading to death may include respiratory failure, [8][a]acute respiratory distress syndrome (ARDS), [9]sepsis and [10]septic shock, [11]thromboembolism, and/or [b]multi-organ failure, including injury of the heart, liver or kidneys. People aged 60 years and over, and those with underlying medical problems such as high blood pressure, heart and lung problems, diabetes, obesity or cancer, are at higher risk of developing serious illness. However, anyone can get infected with COVID-19 and become seriously ill or die. Some people who had COVID-19, whether they have needed hospitalization or not, continue to experience symptoms, including fatigue, respiratory and neurological symptoms.

Viruses constantly change through [12]mutation and sometimes these mutations result in a new [13]variant of the virus. Some variants emerge and disappear while others persist. New variants will continue to emerge. Five SARS-CoV-2 variants have been designated as variants of concern by the WHO: the

Alpha, Beta, Gamma, Delta, and Omicron variants. The Omicron variants are the most recent one announced at the end of 2021. They are believed to be most [14]transmissible even though the World Health Organization (WHO), at the beginning of 2022, claimed that "It is not yet clear whether Omicron is more transmissible compared to other variants, including Delta" and "It is not yet clear whether infection with Omicron causes more severe disease compared to infections with other variants, including Delta."

WHO is working with researchers and patient groups around the world to design and conduct studies on patients beyond the initial acute course of illness to understand the proportion of patients who have long term effects, how long they persist, and why they occur. These studies will be used to develop further guidance for patient care.

Stay safe by taking some simple [15]precautions, such as physical distancing, wearing a mask, especially when distancing cannot be maintained. Keeping rooms well ventilated, avoiding crowds and close contact, regularly cleaning your hands, and coughing into a bent elbow or tissue. Check local advice where you live and work.

In most situations, a [16]molecular test is used to detect SARS-CoV-2 and [c]confirm infection. [17][d]Polymerase chain reaction (PCR) is the most commonly used molecular test. Medical personnel may collect samples from your nose or throat with a [18]swab. Molecular tests detect virus in the sample by amplifying viral genetic material to detectable levels. For this reason, a molecular test is used to confirm an active infection, usually within a few days of

exposure and around the time that symptoms may begin.

Rapid [19]antigen test (sometimes known as a rapid diagnostic test, or RDT) is an alternative in screening people infected by COVID-19. It detects viral [20]proteins (known as antigens). In this test, samples are also collected from the nose or throat using a swab. This test is cheaper than PCR and will offer results more quickly, although they are normally less reliable. Rapid antigen tests perform best when there is more virus circulating in the community and when sampled from an individual during the time they are most infectious.

Both [21]isolation and [22]quarantine are implemented in preventing the spread of COVID-19. Quarantine is required for anyone who has contacted someone infected with the SARS-CoV-2 virus, whether the infected person has symptoms or not. Quarantine means that you remain separated from others because you have been exposed to the virus and you may be infected. Quarantine can be implemented in a designated facility or at home for a period of time (check the announcement of local government). Isolation is adopted for people with COVID-19 symptoms or who have tested positive for the virus. Being in isolation means being separated from other people, ideally in a medically facility where you can receive clinical care. If isolation in a medical facility is not possible and you are not in a high risk group of developing severe disease, isolation may also take place at home. If you are infected with or without symptoms, you should report to the local government and follow their instructions. You may need medical care if you do have symptoms.

Vaccines provide protection to people to against virus. They save millions of lives each year. Vaccines work by training and preparing the body's natural defenses, the [23]immune system, to recognize and fight against the viruses and bacteria they target. After vaccination, if the body is later exposed to those virus or germs, the body is immediately ready to destroy them, preventing illness.

There are several safe and effective vaccines that prevent people from getting seriously ill or dying from COVID-19. This is one part of managing COVID-19, in addition to the main preventive measures of staying at least 1 m away from others, covering a cough or sneeze in your elbow, frequently cleaning your hands, wearing a mask and avoiding poorly ventilated rooms or opening a window.

As of 15 November 2021, WHO has announced that the following vaccines against COVID-19 have met the necessary criteria for safety and [24]efficacy:

- AstraZeneca/Oxford
- Johnson and Johnson
- Moderna
- Pfizer/BionTech
- Sinopharm
- Sinovac
- COVAXIN

Take whatever vaccine is available to you first, even if you have already had COVID-19. It is important to get vaccinated as soon as possible once it's your turn and do not wait. Approved COVID-19 vaccines provide a high degree of protection against getting seriously ill and dying from the disease, although no vaccine is 100% protective.

A
acute：十分嚴重的
age：年齡；衰老
amplifying：擴大；增強
announce：宣佈；通告
antigen：抗原
anxiety：焦慮
appetite：食慾；胃口
B
bacteria：細菌
bent：彎曲的；不直的
C
chest：胸部；胸腔
chill / chill guy：寒冷；冷靜隨和的人
circulate：循環；散佈
claim：聲稱；主張
complication：併發症；複雜化
confirm infection：確認感染
confusion：困惑；混淆
congestion：擁擠；阻塞
conjunctivitis：結膜炎
consciousness：意識；覺悟
contact：聯繫；聯絡
corona / coronavirus：冠狀部；冠狀病毒
cough：咳；咳嗽
curse：詛咒；禍根
D
delirium：神志不清；語無倫次
depression：憂鬱；沮喪
designate：指定；委派
detect：偵測；檢測出
diabetes：糖尿病
diarrhea：腹瀉
disorder：混亂；失調
distance：保持距離
dizziness：眩暈
dose：劑量
E
efficacy：功效；效力
elbow：肘部
emerge：出現；浮現
exposure：暴露；接觸
F
fatigue：疲勞
fever：發燒
H
headache：頭痛
hospitalization：住院治療
I
immune：免疫的
infect：傳染；感染
inflammation：炎症
irritability：易怒；暴脾氣
isolation：隔離；孤獨
J
joint：關節
K
kidney：腎臟
L
lung：肺部
M
mask：面罩；口罩
molecular：分子的
muscle：肌肉
mutation：突變；變異
N
nasal：鼻的
nausea：噁心；嘔吐感
neurological：神經的
O
obesity：肥胖
oxygen：氧氣
P
pandemic：流行病；流行的
PCR：聚合酶連鎖反應
pneumonia：肺炎
precaution：預防措施
prevent / preventive：預防；防止的
proportion：部分；份額
protection：保護；防護
protein：蛋白質
Q
quarantine：隔離檢疫期
R
rash：皮疹
recognize：認出；識別
recover：康復；恢復
reliable：可靠的
respiratory / respiratory failure：呼吸的；呼吸衰竭
S
sample / sampling：樣本；抽樣
sepsis / septic shock：敗血症；敗血性休克
severe：非常嚴重的
shortage：缺乏；缺少
shortness of breath：呼吸困難
smell：嗅覺；氣味
sneeze：打噴嚏
sore：疼痛的
stroke：中風
swab：棉花棒
symptom：癥狀
syndrome：症候群；併發症
T
thromboembolism：血栓栓塞
tissue：面紙；組織
transmissible：可傳播的
treatment：處理；治療
U
underlying：潛在的；深層的
V
vaccinate / vaccination：接種疫苗
variant：變種；變體
ventilate：使通風
viral：病毒的
vomiting：嘔吐