



Fluid – particle interaction force (normal and tangential components)

Let us get the forces in *any chosen two* directions



Munson et al., 2010
Van der Akker, 2009

Fluid – particle interaction force (look in more detail)




$$\int dF_x = \int p \cos \theta dA + \int \tau_w \sin \theta dA \quad \rightarrow \text{Drag}$$

$$\int dF_y = -\int p \sin \theta dA + \int \tau_w \cos \theta dA \quad \rightarrow \text{Lift}$$


Fluid – particle interaction force: interesting situations

Distribution of pressure – flow over *a stationary particle*




What happens in other cases (contours of pressure are shown)?

A particle **moving in a different direction** than the mean fluid flow



Collection of particles moving in a different direction than the mean fluid flow



Drag force

For highly viscous flows: viscous drag (Stokes law)
– only this one an analytical solution


$$\mathbf{F}_D = 3\pi \mu_f d_P (\mathbf{u}_f - \mathbf{u}_P)$$

- Form drag: $1/3 FD$
- Skin (friction) drag: $2/3 FD$

What else can affect the drag force? See next slides.

Drag force - Effect of particle shape?

C_D : the drag force coefficient (to be explained later)




Sphericity



Drag force - Effect of wall?

Drag force increases – introduction of correction coefficients in modelling

$$\frac{C_D}{C_{D,Stokes}} = 1 + \textit{const} \frac{d_p}{h}$$




Drag force – Oscillating and/or distorted particles




C_D : function of the form of a droplet

Drag force – what if the surrounding flow is not a continuum?




A particle “sees” individual molecules

Kn – particle
Knudsen number
(particle
diameter/mean
free path)




Transverse forces can come from:

if the flow is non uniform –
presence of a velocity gradient




if the particle is rotating
(velocity gradient, collisions)




if the particle moves in the
vicinity of a wall

Transverse forces

Saffmann force – due to presence of a velocity gradient (shear)




Why do we have a force if we have such a disturbance velocity field?




Something else is not symmetrical

Vorticity!



Transverse forces

Magnus force – due to perturbation of the mean flow caused by angular motion of the particle



Saffmann lift force – presence of a velocity gradient in the direction normal to the main flow

$$\mathbf{F}_{saff} = 1.61 d_P^2 \sqrt{\rho_f \mu_f} \frac{(\mathbf{u}_f - \mathbf{u}_P) \times \boldsymbol{\omega}_f}{\sqrt{|\boldsymbol{\omega}_f|}}; \quad \boldsymbol{\omega}_f = rot \mathbf{u}_f$$

Magnus lift force – particle rotating in a fluid flow (rotation introduced due to, e.g., collisions)

$$\mathbf{F}_{mag} = \frac{\pi}{8} d_P^3 \rho_f \left(\frac{1}{2} \nabla \times \mathbf{u}_f - \boldsymbol{\omega}_P \right) \times (\mathbf{u}_f - \mathbf{u}_P)$$