

Example i.
Sequential
learning

Example ii.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution

Sequential Bayes
theorem: A rule
for updating
probabilities

Example iii.
Simple linear
regression

Example iv.
Stochastic
volatility
model

Example v.
Multiple linear
regression

Bayesian Ingredients

Hedibert Freitas Lopes

The University of Chicago Booth School of Business
5807 South Woodlawn Avenue, Chicago, IL 60637
<http://faculty.chicagobooth.edu/hedibert.lopes>

hlopes@ChicagoBooth.edu

Outline

- 1 Example *i*. Sequential learning
- 2 Example *ii*. Normal model and normal prior
- 3 Turning the Bayesian crank
 - Posterior and Prior predictive distributions
 - Posterior predictive distribution
 - Sequential Bayes theorem: A rule for updating probabilities
- 4 Example *iii*. Simple linear regression
- 5 Example *iv*. Stochastic volatility model
- 6 Example *v*. Multiple linear regression

Example *i*.
Sequential
learning

Example *ii*.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution

Sequential Bayes
theorem: A rule
for updating
probabilities

Example *iii*.
Simple linear
regression

Example *iv*.
Stochastic
volatility
model

Example *v*.
Multiple linear
regression

Example *i*. Sequential learning

Example *i*.
Sequential
learning

Example *ii*.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution

Sequential Bayes
theorem: A rule
for updating
probabilities

Example *iii*.
Simple linear
regression

Example *iv*.
Stochastic
volatility
model

Example *v*.
Multiple linear
regression

- John claims some discomfort and goes to the doctor.
- The doctor believes John may have the disease A.
- $\theta = 1$: John has disease A; $\theta = 0$: he does not.

- The doctor claims, based on his expertise (H), that

$$P(\theta = 1|H) = 0.70$$

- Examination X is related to θ as follows

$$\begin{cases} P(X = 1|\theta = 0) = 0.40, & \text{positive test given no disease} \\ P(X = 1|\theta = 1) = 0.95, & \text{positive test given disease} \end{cases}$$

Observe $X = 1$

Example i.
Sequential
learning

Example ii.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution

Sequential Bayes
theorem: A rule
for updating
probabilities

Example iii.
Simple linear
regression

Example iv.
Stochastic
volatility
model

Example v.
Multiple linear
regression

Exam's result: $X = 1$

$$P(\theta = 1|X = 1) \propto P(X = 1|\theta = 1)P(\theta = 1)$$

$$\propto (0.95)(0.70) = 0.665$$

$$P(\theta = 0|X = 1) \propto P(X = 1|\theta = 0)P(\theta = 0)$$

$$\propto (0.40)(0.30) = 0.120$$

Consequently

$$P(\theta = 0|X = 1) = 0.120/0.785 = 0.1528662 \text{ and}$$

$$P(\theta = 1|X = 1) = 0.665/0.785 = 0.8471338$$

The information $X = 1$ increases, for the doctor, the probability that John has the disease A from **70%** to **84.71%**.

Posterior predictive

John undertakes the test Y , which relates to θ as follows¹

$$P(Y = 1|\theta = 1) = 0.99 \quad \text{and} \quad P(Y = 1|\theta = 0) = 0.04$$

Then, the predictive of $Y = 0$ given $X = 1$ is given by

$$\begin{aligned} P(Y = 0|X = 1) &= P(Y = 0|\mathbf{X} = \mathbf{1}, \theta = 0)P(\theta = 0|X = 1) \\ &+ P(Y = 0|\mathbf{X} = \mathbf{1}, \theta = 1)P(\theta = 1|X = 1) \\ &= P(Y = 0|\theta = 0)P(\theta = 0|X = 1) \\ &+ P(Y = 0|\theta = 1)P(\theta = 1|X = 1) \\ &= (0.96)(0.1528662) + (0.01)(0.8471338) \\ &= 15.52\% \end{aligned}$$

Key condition: X and Y are conditionally independent given θ .

¹Recall that $P(X = 1|\theta = 1) = 0.95$ and $P(X = 1|\theta = 0) = 0.40$.

Model criticism

Suppose the observed result was $Y = 0$. This is a reasonably unexpected result as the doctor only gave it roughly 15% chance.

He should at least consider rethinking the model based on this result. In particular, he might want to ask himself

- 1 Did 0.7 adequately reflect his $P(\theta = 1|H)$?
- 2 Is test X really so unreliable?
- 3 Is the sample distribution of X correct?
- 4 Is the test Y so powerful?
- 5 Have the tests been carried out properly?

Example i.
Sequential
learning

Example ii.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution
Sequential Bayes
theorem: A rule
for updating
probabilities

Example iii.
Simple linear
regression

Example iv.
Stochastic
volatility
model

Example v.
Multiple linear
regression

Observe $Y = 0$

Let $H_2 = \{X = 1, Y = 0\}$. Then, Bayes theorem leads to

$$\begin{aligned}P(\theta = 1|H_2) &\propto P(Y = 0|\theta = 1)P(\theta = 1|X = 1) \\&\propto (0.01)(0.8471338) = 0.008471338 \\P(\theta = 0|H_2) &\propto P(Y = 0|\theta = 0)P(\theta = 0|X = 1) \\&\propto (0.96)(0.1528662) = 0.1467516\end{aligned}$$

Therefore,

$$P(\theta = 1|X = 1, Y = 0) = \frac{P(Y = 0, \theta = 1|X = 1)}{P(Y = 0|X = 1)} = 0.0545753$$

$$P(\theta = 1|H_i) = \begin{cases} 0.7000 & , H_0: \text{before } X \text{ and } Y \\ 0.8446 & , H_1: \text{after } X=1 \text{ and before } Y \\ 0.0546 & , H_2: \text{after } X=1 \text{ and } Y=0 \end{cases}$$

Example i.
Sequential
learning

Example ii.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution
Sequential Bayes
theorem: A rule
for updating
probabilities

Example iii.
Simple linear
regression

Example iv.
Stochastic
volatility
model

Example v.
Multiple linear
regression

Example *ii*. Normal model and normal prior

Let us now consider a simple measurement error model with normal prior for the unobserved measurement.

$$\begin{aligned}X|\theta &\sim N(\theta, \sigma^2) \\ \theta &\sim N(\theta_0, \tau_0^2)\end{aligned}$$

with σ^2 , θ_0 and τ_0^2 known for now. It is easy to show that the posterior of θ given $X = x$ is also normal.

More precisely, $(\theta|X = x) \sim N(\theta_1, \tau_1^2)$ where

$$\begin{aligned}\theta_1 &= w\theta_0 + (1 - w)x \\ \tau_1^{-2} &= \tau_0^{-2} + \sigma^{-2} \\ w &= \tau_0^{-2}/(\tau_0^{-2} + \sigma^{-2})\end{aligned}$$

w measures the relative information contained in the prior distribution with respect to the total information (prior plus likelihood).

Example *i*.
Sequential
learning

Example *ii*.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution

Sequential Bayes
theorem: A rule
for updating
probabilities

Example *iii*.
Simple linear
regression

Example *iv*.
Stochastic
volatility
model

Example *v*.
Multiple linear
regression

Example from Box & Tiao (1973)

Example i.
Sequential
learning

Example ii.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution
Sequential Bayes
theorem: A rule
for updating
probabilities

Example iii.
Simple linear
regression

Example iv.
Stochastic
volatility
model

Example v.
Multiple linear
regression

Prior A: Physicist A (large experience): $\theta \sim N(900, (20)^2)$

Prior B: Physicist B (not so experienced): $\theta \sim N(800, (80)^2)$.

Model: $(X|\theta) \sim N(\theta, (40)^2)$.

Observation: $X = 850$

$$(\theta|X = 850, H_A) \sim N(890, (17.9)^2)$$

$$(\theta|X = 850, H_B) \sim N(840, (35.7)^2)$$

Information (precision)

Physicist A: from 0.002500 to 0.003120 (an increase of 25%)

Physicist B: from 0.000156 to 0.000781 (an increase of 400%)

Example i.
Sequential
learning

Example ii.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions


Posterior
predictive
distribution

Sequential Bayes
theorem: A rule
for updating
probabilities

Example iii.
Simple linear
regression

Example iv.
Stochastic
volatility
model

Example v.
Multiple linear
regression



Two additional examples

Observations \mathbf{x} , parameters θ and history H .

Likelihood functions/models - $p(\mathbf{x}|\theta, H)$

Example iii : $x_i|\theta, H \sim N(\theta z_i; \sigma^2) \quad i = 1, \dots, n$

Example iv : $x_t|\theta, H \sim N(0; e^{\theta t}) \quad t = 1, \dots, T$

Prior distributions - $p(\theta|H)$

Example iii : $\theta|H \sim N(\theta_0, \tau_0^2)$

Example iv : $\theta_t|H \sim N(\alpha + \beta\theta_{t-1}, \sigma^2) \quad t = 1, \dots, T$

Assume, for now, that $(z_1, \dots, z_n, \sigma^2, \nu_0, \theta_0, \tau_0^2)$ and $(\alpha, \beta, \sigma^2, \theta_0)$ are known and belong to H .

Example i.
Sequential
learning

Example ii.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution
Sequential Bayes
theorem: A rule
for updating
probabilities

Example iii.
Simple linear
regression

Example iv.
Stochastic
volatility
model

Example v.
Multiple linear
regression

Turning the Bayesian crank

Example i.
Sequential
learning

Example ii.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution

Sequential Bayes
theorem: A rule
for updating
probabilities

Example iii.
Simple linear
regression

Example iv.
Stochastic
volatility
model

Example v.
Multiple linear
regression

Posterior (Bayes' Theorem):

$$\begin{aligned} p(\theta|\mathbf{x}, H) &= \frac{p(\theta, \mathbf{x}|H)}{p(\mathbf{x}|H)} \\ &= \frac{p(\mathbf{x}|\theta, H)p(\theta|H)}{p(\mathbf{x}|H)} \\ &\propto p(\mathbf{x}|\theta, H)p(\theta|H) \end{aligned}$$

Prior predictive distribution:

$$p(\mathbf{x}|H) = \int_{\Theta} p(\mathbf{x}|\theta, H)p(\theta|H) d\theta = E_{\theta}[p(\mathbf{x}|\theta, H)]$$

Posterior predictive distribution

Example i.
Sequential
learning

Example ii.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution

Sequential Bayes
theorem: A rule
for updating
probabilities

Example iii.
Simple linear
regression

Example iv.
Stochastic
volatility
model

Example v.
Multiple linear
regression

Let \mathbf{y} be a new set of observations conditionally independent of \mathbf{x} given θ . Then,

$$\begin{aligned} p(\mathbf{y}|\mathbf{x}, H) &= \int_{\Theta} p(\mathbf{y}, \theta|\mathbf{x}, H) d\theta = \int_{\Theta} p(\mathbf{y}|\theta, \mathbf{x}, H) p(\theta|\mathbf{x}, H) d\theta \\ &= \int_{\Theta} p(\mathbf{y}|\theta, H) p(\theta|\mathbf{x}, H) d\theta = E_{\theta|\mathbf{x}} [p(\mathbf{y}|\theta, H)] \end{aligned}$$

Note 1: In general, but not always (time series, for example) \mathbf{x} and \mathbf{y} are independent given θ .

Note 2: It might be more useful to concentrate on prediction rather than on estimation since the former is *verifiable*. In other words, \mathbf{x} and \mathbf{y} can be observed; not θ .

Sequential Bayes theorem: A rule for updating probabilities

Example i.
Sequential learning

Experimental result: $\mathbf{x}_1 \sim p_1(\mathbf{x}_1 | \theta)$

$$p(\theta | \mathbf{x}_1) \propto l_1(\theta; \mathbf{x}_1)p(\theta)$$

Example ii.
Normal model and normal prior

Experimental result: $\mathbf{x}_2 \sim p_2(\mathbf{x}_2 | \theta)$

$$\begin{aligned} p(\theta | \mathbf{x}_2, \mathbf{x}_1) &\propto l_2(\theta; \mathbf{x}_2)p(\theta | \mathbf{x}_1) \\ &\propto l_2(\theta; \mathbf{x}_2)l_1(\theta; \mathbf{x}_1)p(\theta) \end{aligned}$$

Turning the Bayesian crank

Posterior and Prior predictive distributions

Posterior predictive distribution

Sequential Bayes theorem: A rule for updating probabilities

Experimental results: $\mathbf{x}_i \sim p_i(\mathbf{x}_i | \theta)$, for $i = 3, \dots, n$

$$\begin{aligned} p(\theta | \mathbf{x}_n, \dots, \mathbf{x}_1) &\propto l_n(\theta; \mathbf{x}_n)p(\theta | \mathbf{x}_{n-1}, \dots, \mathbf{x}_1) \\ &\propto \left[\prod_{i=1}^n l_i(\theta; \mathbf{x}_i) \right] p(\theta) \end{aligned}$$

Example iv.
Stochastic volatility model

Example v.
Multiple linear regression

Example iii. Simple linear regression

Example i.
Sequential
learning

Example ii.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution

Sequential Bayes
theorem: A rule
for updating
probabilities

Example iii.
Simple linear
regression

Example iv.
Stochastic
volatility
model

Example v.
Multiple linear
regression

Model, prior and posterior:

$$x_i | \theta, H \sim N(\theta z_i; \sigma^2) \quad i = 1, \dots, n$$

$$\theta | H \sim N(\theta_0, \tau_0^2)$$

$$\theta | \mathbf{x}, H \sim N(\theta_1, \tau_1^2)$$

where

$$\tau_1^{-2} = \tau_0^{-2} + \mathbf{z}'\mathbf{z}/\sigma^2 \quad \text{and} \quad \theta_1 = \tau_1^2 (\theta_0 \tau_0^{-2} + \mathbf{z}'\mathbf{x}/\sigma^2)$$

Note 1: As n increases, $\tau_1 \rightarrow \sigma^2(\mathbf{z}'\mathbf{z})^{-1}$ and $\theta_1 \rightarrow (\mathbf{z}'\mathbf{z})^{-1}\mathbf{z}'\mathbf{x}$.

Note 2: The same applies when $\tau_0^{-2} \rightarrow 0$, i.e. with 'little' prior knowledge about θ .

Example iv. SV model

Model, prior and posterior:

$$x_t | \theta_t, H \sim N(0; e^{\theta_t}) \quad t = 1, \dots, T$$

$$\theta_t | H \sim N(\alpha + \beta \theta_{t-1}, \sigma^2) \quad t = 1, \dots, T$$

$$p(\theta | \mathbf{x}, H) \propto \prod_{t=1}^T e^{-\theta_t/2} \exp \left\{ -\frac{1}{2} x_t e^{-\theta_t} \right\} \\ \times \prod_{t=1}^T \exp \left\{ -\frac{1}{2\sigma^2} (\theta_t - \alpha - \beta \theta_{t-1})^2 \right\}$$

Unfortunately, closed form solutions are rare.

- How to compute $E(\theta_{43} | \mathbf{x}, H)$ or $V(\theta_{11} | \mathbf{x}, H)$?
- How to obtain a 95% credible region for $(\theta_{35}, \theta_{36} | \mathbf{x}, H)$?
- How to sample from $p(\theta | \mathbf{x}, H)$?
- How to compute $p(\mathbf{x} | H)$ or $p(x_{T+1}, \dots, x_{T+k} | \mathbf{x}, H)$?

Example i.
Sequential
learning

Example ii.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution

Sequential Bayes
theorem: A rule
for updating
probabilities

Example iii.
Simple linear
regression

Example iv.
Stochastic
volatility
model

Example v.
Multiple linear
regression

Example v. Multiple linear regression

Example i.
Sequential
learning

Example ii.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution

Sequential Bayes
theorem: A rule
for updating
probabilities

Example iii.
Simple linear
regression

Example iv.
Stochastic
volatility
model

Example v.
Multiple linear
regression

The standard Bayesian approach to multiple linear regression is

$$(y|X, \beta, \sigma^2) \sim N(X\beta, \sigma^2 I_n)$$

where $y = (y_1, \dots, y_n)$, $X = (x_1, \dots, x_n)'$ is the $(n \times q)$, design matrix and $q = p + 1$.

The prior distribution of (β, σ^2) is $NIG(b_0, B_0, n_0, S_0)$, i.e.

$$\beta|\sigma^2 \sim N(b_0, \sigma^2 B_0)$$

$$\sigma^2 \sim IG(n_0/2, n_0 S_0/2)$$

for known hyperparameters b_0, B_0, n_0 and S_0 .

Example v. Conditionals and marginals

It is easy to show that (β, σ^2) is $NIG(b_1, B_1, n_1, S_1)$, i.e.

$$\begin{aligned}(\beta|y, X) &\sim N(b_1, \sigma^2 B_1) \\ (\sigma^2|y, X) &\sim IG(n_1/2, n_1 S_1/2)\end{aligned}$$

where

$$\begin{aligned}B_1^{-1} &= B_0^{-1} + X'X \\ B_1^{-1}b_1 &= B_0^{-1}b_0 + X'y \\ n_1 &= n_0 + n \\ n_1 S_1 &= n_0 S_0 + (y - Xb_1)'y + (b_0 - b_1)'B_0^{-1}b_0.\end{aligned}$$

It is also easy to derive the full conditional distributions, i.e.

$$\begin{aligned}(\beta|y, X) &\sim t_{n_1}(b_1, S_1 B_1) \\ (\sigma^2|\beta, y, X) &\sim IG(n_1/2, n_1 S_{11}/2)\end{aligned}$$

where

$$n_1 S_{11} = n_0 S_0 + (y - X\beta)'(y - X\beta).$$

Example i.
Sequential
learning

Example ii.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution

Sequential Bayes
theorem: A rule
for updating
probabilities

Example iii.
Simple linear
regression

Example iv.
Stochastic
volatility
model

Example v.
Multiple linear
regression

Example v. Ordinary least squares

It is well known that

$$\begin{aligned}\hat{\beta} &= (X'X)^{-1}X'y \\ \hat{\sigma}^2 &= \frac{S_e}{n-q} = \frac{(y - X\hat{\beta})'(y - X\hat{\beta})}{n-q}\end{aligned}$$

are the OLS estimates of β and σ^2 , respectively.

The conditional and unconditional sampling distributions of $\hat{\beta}$ are

$$\begin{aligned}(\hat{\beta}|\sigma^2, y, X) &\sim N(\beta, \sigma^2(X'X)^{-1}) \\ (\hat{\beta}|y, X) &\sim t_{n-q}(\beta, S_e)\end{aligned}$$

respectively, with

$$(\hat{\sigma}^2|\sigma^2) \sim IG((n-q)/2, ((n-q)\sigma^2/2)).$$

Example i.
Sequential
learning

Example ii.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution
Sequential Bayes
theorem: A rule
for updating
probabilities

Example iii.
Simple linear
regression

Example iv.
Stochastic
volatility
model

Example v.
Multiple linear
regression

Example v. Sufficient statistics

Example i.
Sequential
learning

Example ii.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution

Sequential Bayes
theorem: A rule
for updating
probabilities

Example iii.
Simple linear
regression

Example iv.
Stochastic
volatility
model

Example v.
Multiple linear
regression

Recall $(y_t|x_t, \beta, \sigma^2) \sim N(x_t'\beta, \sigma^2)$ for $t = 1, \dots, n$, with prior $\beta|\sigma^2 \sim N(b_0, \sigma^2 B_0)$ and $\sigma^2 \sim IG(n_0/2, n_0 S_0/2)$.

Then, for $y^t = (y_1, \dots, y_t)$ and $X^t = (x_1, \dots, x_t)'$, it follows:

$$\begin{aligned}(\beta|\sigma^2, y^t, X^t) &\sim N(b_t, \sigma^2 B_t) \\ (\sigma^2|y^t, X^t) &\sim IG(n_t/2, n_t S_t/2)\end{aligned}$$

where $n_t = n_{t-1} + 1$, $B_t^{-1} = B_{t-1}^{-1} + x_t x_t'$, $B_t^{-1} b_t = B_{t-1}^{-1} b_{t-1} + y_t x_t$
and $n_t S_t = n_{t-1} S_{t-1} + (y_t - b_{t-1}' x_t) y_t + (b_{t-1} - b_t)' B_{t-1}^{-1} b_{t-1}$.

The only ingredients needed are: $x_t x_t'$, $y_t x_t$ and y_t^2 .

These recursions will play an important role later on when deriving **sequential Monte Carlo** methods for conditionally Gaussian dynamic linear models, like many stochastic volatility models.

Example v. Predictive

The predictive density can be seen as the *marginal likelihood*, i.e.

$$p(y|X) = \int p(y|X, \beta, \sigma^2) p(\beta|\sigma^2) p(\sigma^2) d\beta d\sigma^2$$

or, by Bayes' theorem, as the *normalizing constant*, i.e.

$$p(y|X) = \frac{p(y|X, \beta, \sigma^2) p(\beta|\sigma^2) p(\sigma^2)}{p(\beta|\sigma^2, y, X) p(\sigma^2|y, X)}$$

which is valid for all (β, σ^2) .

Closed form solution is available for the multiple normal linear regression:

$$(y|X) \sim t_{n_0}(Xb_0, S_0(I_n + XB_0X')).$$

Example i.
Sequential
learning

Example ii.
Normal model
and normal
prior

Turning the
Bayesian
crank

Posterior and
Prior predictive
distributions

Posterior
predictive
distribution

Sequential Bayes
theorem: A rule
for updating
probabilities

Example iii.
Simple linear
regression

Example iv.
Stochastic
volatility
model

Example v.
Multiple linear
regression