

# Insights into the treatment burden of hereditary angioedema in the evolving treatment landscape

Aleena Banerji, M.D.,<sup>1</sup> Marc A. Riedl, M.D., M.S.,<sup>2</sup> Timothy J. Craig, D.O.,<sup>3</sup> and Cristine Radojicic, M.D.<sup>4</sup>

(Allergy Asthma Proc 42:S1–S3, 2021; doi: 10.2500/aap.2021.42.210018)

**H**ereditary angioedema (HAE) is a rare genetic disease characterized by unpredictable, recurrent, and potentially life-threatening episodes of cutaneous and submucosal swelling in various parts of the body.<sup>1,2</sup> The estimated prevalence of HAE is ~1:50,000.<sup>3,4</sup> HAE is commonly caused by an inherited (~75% of cases) or *de novo* (~25% of cases) alteration in the *SERPINC1* gene, which codes for the serine protease inhibitor C1 esterase inhibitor (C1-INH).<sup>1,3,4</sup> Pathogenic mutations in the *SERPINC1* gene cause C1-INH quantitative deficiency (type I HAE) or dysfunction (type II HAE), which results in dysregulation of the kallikrein–kinin cascade, causing uncontrolled vascular permeability and subsequent swelling.<sup>1,2</sup>

HAE symptoms typically manifest during childhood or early adolescence, with the development of recurrent episodes of angioedema in the extremities, face, gastrointestinal tract, or upper airway.<sup>5–7</sup> Untreated HAE attacks usually resolve within 2–5 days but can cause localized pain and gastrointestinal discomfort and can carry a risk of life-threatening asphyxiation if the larynx is involved.<sup>5,6</sup> Clinical presentation is highly variable among patients

with HAE, with some experiencing swelling episodes multiple times a week and others experiencing swelling attacks less than once per year.<sup>5,6,8</sup> There are some identifiable triggers for HAE attacks (such as physical trauma or stress), but many episodes occur spontaneously, with varying frequency and severity.<sup>5</sup> HAE symptoms can be debilitating if not well controlled and have a substantial impact on patients' quality of life by interfering with their productivity and daily activities.<sup>9,10</sup> In addition, levels of depression and anxiety are elevated in patients with HAE compared with the average population, potentially because of the unpredictable and life-threatening nature of their disease.<sup>7,11,12</sup>

Although there is no cure for HAE, on-demand therapies can be used to reduce the severity and duration of acute attacks, and prophylactic treatments can be used to reduce the severity and frequency of attacks.<sup>2,13</sup> Historically, prophylactic HAE medications were limited by efficacy concerns (antifibrinolytics) or numerous adverse effects (androgens); however, the HAE prophylactic armamentarium has expanded in recent years to include several treatments with improved safety and efficacy.<sup>2,14,15</sup> Current standard-of-care prophylactic treatment options include two self-administered subcutaneous (SC) treatments: a plasma-derived C1-INH concentrate, Haegarda (CSL Behring, King of Prussia, PA) (SC-C1-INH), and a plasma kallikrein inhibitor, Takhzyro (Takeda, Cambridge, MA) (lanadelumab); also, there is a newly approved oral plasma kallikrein inhibitor, Orladeyo (BioCryst, Durham, NC) (berotralstat).<sup>16–19</sup> These therapies have benefited patients greatly; however, patients can experience substantial burdens related to the management of their disease. For example, learning to self-administer an intravenous or SC therapy often requires training, and the self-administration process itself can be cumbersome.<sup>20,21</sup> In addition, repeated injections can be time-consuming and are associated with injection-site pain/reactions and can cause anxiety for patients who are averse to injections.<sup>22–25</sup> Caregivers and family members can also experience tremendous socioeconomic burdens associated with caring for the patient with HAE, including missing work or leisure

---

From the <sup>1</sup>Division of Rheumatology, Allergy and Immunology, Department of Medicine, Massachusetts General Hospital, Boston, Massachusetts, <sup>2</sup>Division of Rheumatology, Allergy, and Immunology, Department of Medicine, University of California, San Diego, La Jolla, California, <sup>3</sup>Department of Medicine and Pediatrics, Penn State University, Hershey Medical Center, Hershey, Pennsylvania, and <sup>4</sup>Department of Medicine, Duke University School of Medicine, Durham, North Carolina

A. Banerji has received research funding from Takeda and BioCryst, and consulting fees from Takeda, BioCryst, CSL Behring, Pharming, KalVista, and Pharoaris. M. Riedl has been a consultant and/or speaker for and/or has received grants from Adverum, Attune, BioCryst, BioMarin, CSL Behring, Ionis, KalVista, Pfizer, Pharming, Pharoaris, REGENXBIO, and Shire/Takeda. T. Craig has received research funding from BioCryst, CSL Behring, Takeda, and Ionis; speaker fees from CSL Behring and Takeda; and consulting fees from BioCryst, CSL Behring, Takeda, and Pharming. C. Radojicic has received consulting fees from BioCryst and CSL Behring, and speaker fees from CSL Behring

This study was deemed exempt by the Western Institutional Review Board and was funded by BioCryst Pharmaceuticals, Inc. Writing support was provided by Ashly Pavlovsky, Ph.D., from Porterhouse Medical Group

Supplemental data available at [www.IngentaConnect.com](http://www.IngentaConnect.com)

Address correspondence to Aleena Banerji, M.D., Allergy and Immunology, Department of Medicine, Massachusetts General Hospital, 100 Blossom Street, Boston, MA 02114

E-mail address: [abanerji@mgh.harvard.edu](mailto:abanerji@mgh.harvard.edu)

Copyright © 2021, OceanSide Publications, Inc., U.S.A.

---

days to help the patient they care for and absorbing the financial costs of chronic disease management (such as long-term treatment and hospitalizations).<sup>26–28</sup>

As the HAE treatment landscape continues to evolve, it is important to understand how modern prophylactic treatments are used and how they affect patients' and caregivers' quality of life. Personal characteristics, lifestyle, and disease course are unique to each patient; therefore, personalized medicine facilitated by shared decision-making should be the goal of HAE management as more therapies become available. We conducted three independent, corresponding surveys to better understand the patient, caregiver, and physician experiences with managing HAE and to identify factors that impact prophylaxis-associated treatment burden, overall well-being of patients and caregivers, and treatment preferences. The three studies provide valuable insights into the patient (Radojicic *et al.*),<sup>29</sup> caregiver (Craig *et al.*),<sup>30</sup> and physician (Riedl *et al.*)<sup>31</sup> perspectives.

The survey questions covered a range of topics, including respondent characteristics and current experience with prophylactic treatments. Fully anchored, 4-point Likert scales were used to assess agreement (strongly agree, somewhat agree, somewhat disagree, strongly disagree). Responses of "somewhat agree" and "strongly agree" were reported as "agree" unless otherwise noted. Other Likert scales, such as 10-point scales, were used to assess survey responses that measured impact, influence, likelihood, and satisfaction. A table available in the supplementary material (Supplemental Table 1) provides a list of questions and responses included in the reports, although this is not a comprehensive list of all the questions included in the surveys.

Survey-based studies such as those reported in this supplement are commonly conducted to understand various stakeholder perspectives; however, we recognize that studies of this nature have limitations that should be considered. For example, the respondents included in the study were identified in collaboration with an online panel company to which the respondents had provided permission to be contacted for research purposes. This recruitment method may have introduced selection bias to the study results. In addition, the collected data were from self-reported surveys, which may have introduced responder bias.

## REFERENCES

- Banday AZ, Kaur A, Jindal AK, et al. An update on the genetics and pathogenesis of hereditary angioedema. *Genes Dis.* 2020; 7:75–83.
- Longhurst HJ, Bork K. Hereditary angioedema: an update on causes, manifestations and treatment. *Br J Hosp Med (Lond).* 2019; 80:391–398.
- Lumry WR, Settipane RA. Hereditary angioedema: epidemiology and burden of disease. *Allergy Asthma Proc.* 2020; 41(suppl 1):S08–S13.
- Busse PJ, Christiansen SC, Riedl MA, et al. US HAEA Medical Advisory Board 2020 guidelines for the management of hereditary angioedema. *J Allergy Clin Immunol Pract.* 2021;9:132–150. e3. (DOI: <http://dx.doi.org/10.1016/j.jaip.2020.08.046>).
- Johnston DT. Diagnosis and management of hereditary angioedema. *J Am Osteopath Assoc.* 2011; 111:28–36.
- Bork K, Meng G, Staubach P, et al. Hereditary angioedema: new findings concerning symptoms, affected organs, and course. *Am J Med.* 2006; 119:267–274.
- Banerji A, Davis KH, Brown TM, et al. Patient-reported burden of hereditary angioedema: findings from a patient survey in the United States. *Ann Allergy Asthma Immunol.* 2020; 124:600–607.
- Banerji A, Busse P, Christiansen SC, et al. Current state of hereditary angioedema management: a patient survey. *Allergy Asthma Proc.* 2015; 36:213–217.
- Lumry WR, Castaldo AJ, Vernon MK, et al. The humanistic burden of hereditary angioedema: impact on health-related quality of life, productivity, and depression. *Allergy Asthma Proc.* 2010; 31:407–414.
- Aygören-Pürsün E, Bygum A, Beusterien K, et al. Estimation of EuroQol 5-Dimensions health status utility values in hereditary angioedema. *Patient Prefer Adherence.* 2016; 10:1699–1707.
- Fouche AS, Saunders EFH, Craig T. Depression and anxiety in patients with hereditary angioedema. *Ann Allergy Asthma Immunol.* 2014; 112:371–375.
- Caballero T, Aygören-Pürsün E, Bygum A, et al. The humanistic burden of hereditary angioedema: results from the Burden of Illness Study in Europe. *Allergy Asthma Proc.* 2014; 35:47–53.
- Maurer M, Magerl M, Ansotegui I, et al. The international WAO/EAACI guideline for the management of hereditary angioedema – the 2017 revision and update. *Allergy.* 2018; 73:1575–1596. <https://pubmed.ncbi.nlm.nih.gov/29318628/>.
- Bork K, Bygum A, Hardt J. Benefits and risks of danazol in hereditary angioedema: a long-term survey of 118 patients. *Ann Allergy Asthma Immunol.* 2008; 100:153–161.
- Zanichelli A, Vacchini R, Badini M, et al. Standard care impact on angioedema because of hereditary C1 inhibitor deficiency: a 21-month prospective study in a cohort of 103 patients. *Allergy.* 2011; 66:192–196.
- TAKHZYRO (lanadelumab-flyo) – prescribing information; 2018. Available online at [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/761090s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/761090s000lbl.pdf); accessed December 15, 2020.
- HAEGARDA (C1 esterase inhibitor subcutaneous [human]) – prescribing information; September 2020. Available online at <https://www.fda.gov/media/105611/download>; accessed December 15, 2020.
- ORLADEYO (berotralstat) capsules, for oral use – prescribing information; December 2020. Available online at [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/214094s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/214094s000lbl.pdf); accessed December 16, 2020.
- US Hereditary Angioedema Association. FDA approved treatments. Available online at <https://www.haea.org/pages/p/ApprovedTreatments>; accessed December 16, 2020.
- Caballero T, Sala-Cunill A, Cancian M, et al. Current status of implementation of self-administration training in various regions of Europe, Canada and the USA in the management of hereditary angioedema. *Int Arch Allergy Immunol.* 2013; 161(suppl 1):10–16.
- Murphy E, Donahue C, Omert L, et al. Training patients for self-administration of a new subcutaneous C1-inhibitor concentrate for hereditary angioedema. *Nurs Open.* 2018; 6:126–135.
- Banerji A, Riedl MA, Bernstein JA, et al. Effect of lanadelumab compared with placebo on prevention of hereditary angioedema attacks: a randomized clinical trial. *JAMA.* 2018; 320:2108–2121.

23. Longhurst H, Cicardi M, Craig T, et al. Prevention of hereditary angioedema attacks with a subcutaneous C1 inhibitor. *N Engl J Med*. 2017; 376:1131–1140.
24. Wright S, Yelland M, Heathcote K, et al. Fear of needles: nature and prevalence in general practice. *Aust Fam Physician*. 2009; 38:172–176.
25. Tuong L-AC, Olivieri K, Craig TJ. Barriers to self-administered therapy for hereditary angioedema. *Allergy Asthma Proc*. 2014; 35:250–254.
26. Bygum A, Aygören-Pürsün E, Beusterien K, et al. Burden of illness in hereditary angioedema: a conceptual model. *Acta Derm Venereol*. 2015; 95:706–710.
27. Aygören-Pürsün E, Bygum A, Beusterien K, et al. Socioeconomic burden of hereditary angioedema: results from the hereditary angioedema burden of illness study in Europe. *Orphanet J Rare Dis*. 2014; 9:99.
28. Lumry WR. Hereditary angioedema: the economics of treatment of an orphan disease. *Front Med (Lausanne)*. 2018; 5:22.
29. Radojicic C, Riedl MA, Craig TJ, et al. Patient perspectives on the treatment burden of injectable medication for hereditary angioedema. *Allergy Asthma Proc*. 2021; 42(Suppl 1):S4–S10.
30. Riedl MA, Craig TJ, Banerji A, et al. Physician and patient perspectives on the management of hereditary angioedema: a survey on treatment burden and needs. *Allergy Asthma Proc*. 2021; 42(Suppl 1):S17–S25.
31. Craig TJ, Banerji A, Riedl MA, et al. Caregivers' role in managing hereditary angioedema and perceptions of treatment-related burden. *Allergy Asthma Proc*. 2021; 42(Suppl 1):S11–S16. □