
Smooth dogfish s(Surface temp.) s(Bottom temp.) s(Surface salinity) 0 --5 **-**-10 Component smooth _1 <u>-2</u> <u>-1</u> Ö 2 Ö 2 3 -6 **-**3 Ö 1 s(Bottom salinity) s(Depth) 0 --5 **-**-10 **-**-15 - -6 -3 0 0 2

Covariate


Chain dogfish s(Surface temp.) s(Bottom temp.) s(Surface salinity) 0 -4 _8 **-**Component smooth <u>-2</u> <u>-1</u> Ö 2 <u>-</u>1 Ö 2 3 -6 <u>-</u>3 Ö s(Bottom salinity) s(Depth) 0 --4 -8 **-**-12 **-**-3 0 0 2 -6 6 Covariate

Spiny dogfish s(Surface temp.) s(Bottom temp.) s(Surface salinity) 5.0 2.5 0.0 -2.5 -5.0 Component smooth -2 <u>-1</u> Ö 2 <u>-1</u> Ö 2 3 -6 -3 Ö 1 s(Bottom salinity) s(Depth) 2.5 0.0 -2.5-5.0 -3 0 0 -6 Covariate


Barndoor skate




Winter skate




Clearnose skate




Rosette skate




Little Skate




Smooth skate




Thorny skate




Silver hake



Atlantic cod s(Surface temp.) s(Bottom temp.) s(Surface salinity) 2 -2 <u>-</u>1 Ö 2 <u>-</u>1 Ö 1 3 -6 -3 Ö s(Bottom salinity) s(Depth)




Haddock




Pollock s(Surface temp.) s(Bottom temp.) s(Surface salinity) 5.0 2.5 0.0 -2.5-5.0 Component smooth -3 -2 <u>-1</u> Ö 2 <u>-</u>1 Ö 1 2 3 -6 Ö 1 s(Bottom salinity) s(Depth) 5.0 2.5 0.0 -2.5 -5.0 -6 -3 0 0 Covariate


White hake




Red hake



Spotted hake



Fourbeard rockling



American plaice s(Surface temp.) s(Surface salinity) s(Bottom temp.) 5.0 2.5 0.0 -2.5Component smooth -3 -2 <u>-1</u> Ö 2 <u>-</u>1 Ö i 2 3 -6 Ö 1 s(Bottom salinity) s(Depth) 5.0 2.5 0.0 -2.5

-5.0

-6


-3

0


0

Covariate


Summer flounder



Fourspot flounder



Yellowtail flounder




Winter flounder s(Surface temp.) s(Bottom temp.) s(Surface salinity) 0 --20 **-**-40 **-**Component smooth <u>-2</u> <u>-1</u> Ö 2 <u>-1</u> Ö 2 3 -6 -3 Ö s(Bottom salinity) s(Depth) -20 **-**-40 **-**-60 **-**-3 0 0


Covariate

-6

Witch flounder



Windowpane s(Surface temp.) s(Surface salinity) s(Bottom temp.) 2.5 0.0 -2.5 -5.0Component smooth -2 <u>-1</u> Ö 2 <u>-1</u> Ö 2 3 -6 -3 Ö 1 s(Bottom salinity) s(Depth) 2.5 0.0 -2.5 -5.0-7.5 -3 0 0 -6 Covariate



Atlantic croaker s(Surface temp.) s(Bottom temp.) s(Surface salinity) 2.5 0.0 -2.5 Component smooth -3 -2 <u>-1</u> Ö 2 <u>-</u>1 Ö i 2 3 -6 Ö s(Bottom salinity) s(Depth) 2.5 0.0 -2.5

-5.0

-6


-3

0

0


Covariate

Black sea.bass



Scup s(Surface temp.) s(Bottom temp.) s(Surface salinity) 2.5 0.0 -2.5 Component smooth <u>-</u>2 <u>-1</u> Ö 2 <u>-1</u> Ö 1 2 3 -6 -3 Ö 1 s(Bottom salinity) s(Depth) 2.5 0.0 -2.5 -3 0 0 -6 Covariate

Weakfish



Acadian redfish s(Surface temp.) s(Bottom temp.) s(Surface salinity) 0 --5 **-**-10 **-**Component smooth –15 **-**-2 <u>-1</u> Ö 2 <u>-</u>1 Ö i 2 3 -6 **-**3 Ö 1 s(Bottom salinity) s(Depth) 0 --5 **-**-10 **-**–15 **-**

-3

-6

0


0


2

Covariate


6

Blackbelly rosefish





Sea Raven



Northern searobin s(Surface temp.) s(Bottom temp.) s(Surface salinity) 0 --4 -8 **-**Component smooth _1 -3 <u>-2</u> <u>-1</u> Ö 2 Ö 2 3 -6 Ö 1 s(Bottom salinity) s(Depth) 0 --4

-12 **-**


-6


-3

0


0

2 Covariate Striped searobin





Fawn cusk.eel



Goosefish

