

Classification of quantum graphs over complex 3×3 matrices

(Klasyfikacja grafów kwantowych nad
zespolonymi macierzami 3×3)

Filip Rabięga

Master's thesis

Supervisor: Dr. Mariusz Tobolski

Uniwersytet Wrocławski
Wydział Matematyki i Informatyki
Instytut Matematyczny

February 31st, 2026

Abstract

According to all known laws of aviation, there is no way a bee should be able to fly. Its wings are too small to get its fat little body off the ground. The bee, of course, flies anyway because bees don't care what humans think is impossible.

Według wszystkich znanych praw lotnictwa, pszczoła nie powinna w ogóle latać. Jej skrzydła są zbyt małe, by unieść jej pulchne ciało z ziemi. Pszczoła, oczywiście, lata mimo wszystko, ponieważ pszczoły nie przejmują się tym, co ludzie uważają za niemożliwe.

Contents

1	Introduction	7
1.1	Motivation	7
1.2	Scope	7
1.3	Acknowledgements	7
2	Preliminaries	9
2.1	Group theory	9
2.2	C^* -algebras	10
2.2.1	Crossed-product C^* -algebra	11
3	Quantum sets and quantum graphs	13
3.1	Quantum sets	13
3.1.1	Crossed product quantum sets	17
3.2	Quantum graphs	17
3.3	Voltage and Derived Quantum Graphs	18
3.3.1	Classical Voltage Graphs	18
3.3.2	Generalisation to quantum voltage and derived quantum graphs	20
3.4	Isomorphisms and Quantum Isomorphisms of Quantum Sets	21
4	Quantum graphs over 3×3 matrices	23
4.1	Gromada's theorem	23

Chapter 1

Introduction

1.1 Motivation

1.2 Scope

1.3 Acknowledgements

Chapter 2

Preliminaries

2.1 Group theory

In the following section we will introduce some group theoretic notions that be used throughout this thesis. We assume that the Reader is familiar with the notion of a group.

Definition 2.1.1 (Dual of a group). Let G be a group. The *dual group* $\hat{G}$ is the set of all homomorphisms from G into the multiplicative group of roots of unity in $\mathbb{C}$ with the group operation defined as

$$(\varphi\psi)(g) = \varphi(g)\psi(g)$$

for all $\varphi, \psi \in \hat{G}$ and $g \in G$, where the latter multiplication takes place in $\mathbb{C}$.

Remark 2.1.1. We will often call the elements of the dual group *characters*. If a group is isomorphic to its complement it will be called *self-dual*.

The following proposition will be used throughout the thesis, as almost all of the groups we will be dealing with will be finite and abelian.

Proposition 2.1.1. *If G is finite and abelian, then it is self-dual.*

Proof. First we will prove the statement for finite cyclic groups. Let $G = \langle g \rangle$, $\text{ord}(g) = n$. We will prove that the dual is cyclic of order n . Let $\chi: G \rightarrow \mathbb{C}$ be a character. Since the group is cyclic, the character is entirely determined by the value it takes on the generator g . Since $g^n = e$, obviously $\chi(g)^n = 1$, so $\chi(g)^n \in \mu_n = \{\exp(2k\pi i/n) : k = 0, 1, \dots, n-1\}$. For each possible choice of $\chi(g) \in \mu_n$ we get a different character. It is then easy to see that $\chi: G \rightarrow \mathbb{C}$ such that $\chi(g) = \exp(2\pi i/n)$ generates this group, i.e. $\hat{G} = \{\chi, \chi^2, \dots, \chi^{n-1}, \chi^n = e_{\hat{G}}\}$.

Now since every finite abelian group can be written as a finite product of finite cycles, all we need to do is check whether the statement holds for a product of two

groups A, B that are themselves self-dual, i.e. $\widehat{A \times B} \cong \widehat{A} \times \widehat{B}$. This is self-evident as every character $\chi \in \widehat{A \times B}$ is determined by its restrictions to $A \times \{e_B\}$ and $\{e_A\} \times B$, since

$$\chi(a, b) = \chi((a, e_B)(e_A, b)) = \chi(a, e_B)\chi(e_A, b).$$

Thus $\chi \leftrightarrow (\chi|_A, \chi|_B)$ is an isomorphism. $\square$

2.2 C^* -algebras

In the following section we will introduce some notions stemming from the theory of C^* -algebras. We assume the Reader is familiar with the notion of a C^* -algebra. For reference, see Blackadar [1].

Multiplication in a C^* -algebra A is a bilinear map

$$m: A \times A \rightarrow A.$$

By the universal property of the tensor product, m extends uniquely to a linear map

$$m^*: A \otimes A \rightarrow A.$$

Throughout this thesis, we will use the same notation (m) for both the bilinear multiplication and its unique linear extension, relying on the context to distinguish between the two.

Definition 2.2.1 (Linear functional). A *linear functional* on a C^* -algebra A is a linear function $\psi: A \rightarrow \mathbb{C}$.

Remark 2.2.1. We will call linear functionals simply as *functionals*.

The following notions will be useful for defining an inner product on C^* -algebras. Positiveness and faithfulness are all we need to define a norm on a finite-dimensional C^* -algebra equipped with a suitable functional.

Definition 2.2.2 (Positive functional). A functional ψ on a C^* -algebra A is called *positive*, if

$$\psi(a^*a) \geq 0$$

for any $a \in A$.

Definition 2.2.3 (Faithful functional). A positive functional ψ on a C^* -algebra A is called *faithful*, if

$$\psi(a^*a) = 0 \Leftrightarrow a = 0$$

for any $a \in A$.

Remark 2.2.2. In the literature, the functional ψ is often called a *state*.

The following notion is akin to the notion of a group action on a set.

Definition 2.2.4 (Group action on a C^* -algebra). By *group action* of a group G on a C^* -algebra A we mean a group homomorphism

$$\alpha: G \rightarrow \text{Aut}(A).$$

We will write $g \cdot a$ instead of $\alpha(g)(a)$ then the action in question is clear.

2.2.1 Crossed-product C^* -algebra

In this section we will introduce the notion of a crossed-product C^* -algebra. We will do so in a limited manner, only to the extent to which the theory of such products is necessary for our purposes. For a more general reference we refer the Reader to Blackadar [1]. $\alpha: G \rightarrow \text{Aut}(A)$ is a group action on A . For a C^* dynamical system (A, G, α) with G finite we define

$$C_c(G, A, \alpha) = \{f: G \rightarrow A: \text{supp}(f) \text{ is finite}\},$$

i.e. the space of all finitely supported functions on G with values in A .

$C_c(G, A, \alpha)$ is equipped with both a product and a $*$ -operation in such a way that the action becomes "inner", i.e. $g \cdot a = u_g a u_g^*$ for all $g \in G, a \in A$. More precisely we define those two operations in the following way:

$$ab = \sum_{g, h \in G}$$

Definition 2.2.5 (Covariant representation). Fix a dynamical system (A, G, α) . A *covariant representation* (π, u, H) of (A, G, α) consist of a unitary representation $u: G \rightarrow B(\mathcal{H})$ of G and a representation $\pi: A \rightarrow B(\mathcal{H})$ of A such that $u(g)\pi(a)u(g)^* = \pi(g \cdot a)$ for every $g \in G$ and $a \in A$.

Definition 2.2.6 (Full C^* -algebra norm on $C_c(G, A, \alpha)$). The full C^* -algebra norm on $C_c(G, A, \alpha)$ is given by

$$\|\cdot\| = \sup \|(\pi \times u)(\cdot)\|,$$

where the supremum is taken over all covariant representations (π, u, H) of (A, G, α) .

Definition 2.2.7 (Crossed product C^* -algebra). The crossed product of a C^* -algebra and a discrete group G is the completion of $C_c(G, A, \alpha)$ with respect to the full norm. We denote the crossed product by $\tilde{B} \rtimes_{\hat{\alpha}} \hat{G}$.

Chapter 3

Quantum sets and quantum graphs

In this chapter we will introduce the notions of quantum sets and quantum graphs. They were first considered by...

3.1 Quantum sets

Let B be a C^* -algebra equipped with a faithful positive functional ψ . By the Gelfand-Naimark theorem, B can be realized concretely as a closed $*$ -subalgebra of the bounded linear operators $B(\mathcal{H})$ on some Hilbert space $\mathcal{H}$. When B is finite-dimensional, this abstract framework simplifies considerably: we can identify B with its own Hilbert space completion $L^2(B, \psi)$ by defining the inner product directly via the functional:

$$\langle x, y \rangle = \psi(x^*y) \quad \forall x, y \in B.$$

The following definition introduces one of the central notions used in this thesis. It is taken directly from Schäfer i Tobolski [5].

Definition 3.1.1. (Quantum set, Schäfer i Tobolski [5, Definition 2.1]) A quantum set (B, ψ) is a pair consisting of a finite-dimensional C^* -algebra B and a faithful positive functional $\psi: B \rightarrow \mathbb{C}$ such that

$$mm^* = \text{Id}_B$$

holds for the multiplication map $m: B \otimes B \rightarrow B$ and its adjoint m^* given by

$$\langle m(a \otimes b), c \rangle = \langle a \otimes b, m^*(c) \rangle_{\psi \otimes \psi}$$

for all $a, b, c \in B$.

Remark 3.1.1. We will call any functional with the property given in the definition above a *quantum set functional* on B .

Remark 3.1.2. Recall that in general, a *comultiplication* on a C^* -algebra is a $*$ -homomorphism

$$\Delta \rightarrow A \otimes A$$

such that the following diagram commutes:

$$\begin{array}{ccc} A & \xrightarrow{\Delta} & A \otimes A \\ \Delta \downarrow & & \downarrow \Delta \otimes \text{id} \\ A \otimes A & \xrightarrow{\text{id} \otimes \Delta} & A \otimes A \otimes A \end{array}$$

Note that in general a comultiplication is *not* unique. In our setting it is uniquely determined by the additional structure and the Riesz representation theorem for Hilbert spaces.

Remark 3.1.3. Since in our setup the comultiplication is given by the scalar product and regular multiplication m , we will denote it by m^* .

Example 3.1.1. Every finite set X gives rise to a quantum set $(C(X), \psi_X)$ with $\psi_X: C(X) \rightarrow \mathbb{C}$, $\psi_X(f) = \sum_{x \in X} f(x)$ for any $f \in C(X)$. Let X be a finite set and $C(X)$ be the $*$ -algebra of complex-valued functions on X , where the involution is given by pointwise complex conjugation: $f^*(x) = \overline{f(x)}$.

To show that ψ_X is positive, we must verify that $\psi_X(f^*f) \geq 0$ for all $f \in C(X)$. For any function f , the product f^*f evaluates pointwise to:

$$(f^*f)(x) = \overline{f(x)}f(x) = |f(x)|^2.$$

Applying the functional yields:

$$\psi_X(f^*f) = \sum_{x \in X} |f(x)|^2.$$

Thus the functional is positive.

To prove faithfulness, suppose $\psi_X(f^*f) = 0$ for some $f \in C(X)$. This implies:

$$\sum_{x \in X} |f(x)|^2 = 0.$$

Because each term $|f(x)|^2$ is strictly non-negative, a sum of such terms can only equal zero if every individual term is zero. Thus, we must have:

$$|f(x)|^2 = 0 \quad \forall x \in X,$$

which directly implies $f(x) = 0$ for all $x \in X$. Therefore, f is the zero function ($f = 0$), establishing that ψ_X is faithful.

Let $\{e_x\}_{x \in X}$ be the standard orthonormal basis of characteristic functions on X . Recall the definitions of the multiplication map m and its adjoint m^* :

$$\begin{aligned} m(f \otimes g) &= fg, \\ m^*(e_x) &= e_x \otimes e_x. \end{aligned}$$

We wish to show that the composition $mm^*: C(X) \rightarrow C(X)$ is the identity operator (Id). Let us evaluate the action of mm^* on an arbitrary basis element e_x :

$$(mm^*)(e_x) = m(m^*(e_x)) = m(e_x \otimes e_x) = e_x e_x = e_x.$$

Because the linear operators mm^* and Id agree on every element of the basis $\{e_x\}_{x \in X}$, by linearity they must be equal on the entire space $C(X)$. Therefore:

$$mm^* = \text{Id}.$$

Example 3.1.2. In particular, $(\mathbb{C}^n, \psi_n)$ with $\psi_n(e_i) = 1$ is a quantum set. The functional naturally extends to:

$$\psi_n(x) = \sum_{i=1}^n x_i.$$

To show that ψ_n is positive, we evaluate it on the element x^*x . Using the orthogonality of the basis, we have:

$$x^*x = \left(\sum_{i=1}^n \overline{x_i} e_i \right) \left(\sum_{j=1}^n x_j e_j \right) = \sum_{i,j=1}^n \overline{x_i} x_j e_i e_j = \sum_{i=1}^n \overline{x_i} x_i e_i = \sum_{i=1}^n |x_i|^2 e_i.$$

Applying the functional yields:

$$\psi_n(x^*x) = \psi_n \left(\sum_{i=1}^n |x_i|^2 e_i \right) = \sum_{i=1}^n |x_i|^2 \psi_n(e_i) = \sum_{i=1}^n |x_i|^2.$$

Since $|x_i|^2 \geq 0$ for all i , the sum of these non-negative real numbers is also non-negative, meaning $\psi_n(x^*x) \geq 0$. Thus, the functional is positive.

To prove faithfulness, suppose $\psi_n(x^*x) = 0$ for some $x \in \mathbb{C}^n$. This implies:

$$\sum_{i=1}^n |x_i|^2 = 0.$$

Because each term $|x_i|^2$ is strictly non-negative, the sum can only be zero if every individual component satisfies $|x_i|^2 = 0$ for all $i \in \{1, \dots, n\}$. This directly implies $x_i = 0$ for all i . Therefore, $x = \sum_{i=1}^n 0 \cdot e_i = 0$, establishing that ψ_n is faithful.

The adjoint map $m^*: \mathbb{C}^n \rightarrow \mathbb{C}^n \otimes \mathbb{C}^n$ acts on a basis element e_i by:

$$m^*(e_i) = e_i \otimes e_i.$$

We wish to show that the composition $mm^*: \mathbb{C}^n \rightarrow \mathbb{C}^n$ is the identity operator. We check:

$$(mm^*)(e_i) = m(m^*(e_i)) = m(e_i \otimes e_i) = e_i.$$

Because the linear operators mm^* and Id agree on every element of the basis $\{e_i\}_{i=1}^n$, by linearity they must be equal on the entire vector space $\mathbb{C}^n$. Therefore:

$$mm^* = \text{Id}.$$

Example 3.1.3. For any $n \in \mathbb{N}_+$, $(M_n, n \text{ Tr})$ is a quantum set, with Tr being the trace. We will later show that this structure can be obtained as the crossed product $(\mathbb{C}^n, \psi_n)$ and $\hat{\mathbb{Z}}_n$ with the standard action, i.e.

$$(M_n, n \text{ Tr}) \cong (\mathbb{C}^n \rtimes \hat{\mathbb{Z}}_n, \psi_{\rtimes}).$$

This is proven in 3.1.2.

Lemma 3.1.1 (Comultiplication for matrix algebras). *In the context of $(M_n, n \text{ Tr})$, the comultiplication is given by the formula*

$$m^*(e_{ij}) = \sum_{k=1}^n e_{ik} \otimes e_{kj}$$

with e_{ij} being the matrix with a single 1 in the i -th row and j -th column, and extended linearly.

Proof. This is simply a computational exercise. We will use the following well-known formulas:

1. $e_{ab}e_{cd} = \delta_{bc}e_{ad}$,
2. $\langle e_{ij}, e_{kl} \rangle = \text{Tr}(e_{ji}e_{kl}) = \delta_{ik}\delta_{jl}$,

with

$$\delta_{ij} = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{otherwise.} \end{cases}$$

In particular $\{e_{ij}\}_{i,j=1,\dots,n}$ form an orthonormal basis.

Take arbitrary basis vectors e_{ab}, e_{cd} . Then

$$m(e_{ab} \otimes e_{cd}) = e_{ab}e_{cd} = \delta_{bc}e_{ad},$$

hence

$$\begin{aligned} \langle m(e_{ab} \otimes e_{cd}), e_{ij} \rangle &= \delta_{bc} \langle e_{ad}, e_{ij} \rangle \\ &= \delta_{bc} \delta_{ai} \delta_{dj}. \end{aligned}$$

Now compute the inner product with the proposed adjoint:

$$\begin{aligned} \left\langle e_{ab} \otimes e_{cd}, \sum_{k=1}^n e_{ik} \otimes e_{kj} \right\rangle &= \sum_{k=1}^n \langle e_{ab}, e_{ik} \rangle \langle e_{cd}, e_{kj} \rangle \\ &= \sum_{k=1}^n \delta_{ai} \delta_{bk} \delta_{ck} \delta_{dj} \\ &= \delta_{ai} \delta_{bc} \delta_{dj}. \end{aligned}$$

The two expressions agree. Since the matrix units form an orthonormal basis, this linearly extends to all matrices. $\square$

3.1.1 Crossed product quantum sets

Schäfer i Tobolski [5] have introduced the notion of crossed product quantum sets. Suppose we are given a finite-dimensional C^* -algebra $\tilde{B}$ along with a dual group action $\hat{\alpha}: \hat{G} \rightarrow \text{Aut}(\tilde{B})$ where G is finite and abelian. One can then consider the crossed product

$$B := \tilde{B} \rtimes_{\hat{\alpha}} \hat{G}.$$

If $\tilde{B}$ is equipped with a faithful functional $\tilde{\psi}$ in such a way as to make $(\tilde{B}, \tilde{\psi})$ a quantum set, one can define a new faithful functional on B to make it a quantum set, as described below.

Definition 3.1.2. (Crossed product quantum set, Schäfer i Tobolski [5, Definition 3.1]) Let $(\tilde{B}, \tilde{\psi})$ be a quantum set and let $\hat{\alpha}: \hat{G} \rightarrow \text{Aut}(\tilde{B}, \tilde{\psi})$ be an action of the dual of a finite abelian group G . The *crossed product quantum set* is defined as

$$(\tilde{B}, \tilde{\psi}) \rtimes_{\hat{\alpha}} \hat{G} := \left(\tilde{B} \rtimes_{\hat{\alpha}} \hat{G}, \psi_{\rtimes} \right),$$

where $\tilde{B} \rtimes_{\hat{\alpha}} \hat{G}$ is the crossed product C^* -algebra and $\psi_{\rtimes}$ is the functional given by

$$\psi_{\rtimes} \left(\sum_{\chi \in \hat{G}} b_{\chi} u_{\chi} \right) = n \tilde{\psi}(b_1)$$

with $n = |\hat{G}|$.

The proof that this definition is sensible, i.e. the newly defined object is indeed a quantum set, can be seen in Schäfer i Tobolski [5], Proposition 3.4.

Fact 3.1.1. (tobolksi26, Example 3.5) If $\hat{\alpha} = \text{triv}$, i.e. it is the trivial action, then

$$\tilde{B} \rtimes_{\hat{\alpha}} \hat{G} \cong \tilde{B} \otimes C(G),$$

where the isomorphism is given by $b_{\chi} u_{\chi} \mapsto b_{\chi} \otimes \tilde{u}_{\chi}$, where $\tilde{u}_{\chi} \in C(G)$ is the function that evaluates to $\chi(g)$ on $g \in G$.

Lemma 3.1.2. Let $\hat{\alpha}: \hat{\mathbb{Z}}_n \rightarrow A$ be the action of cyclically shifting elements. Then

$$(M_n, n \text{ Tr}) \cong (\mathbb{C}^n \rtimes_{\alpha} \hat{\mathbb{Z}}_n, \psi_{\rtimes}).$$

3.2 Quantum graphs

A simple graph is classically defined as a pair (V, E) with V serving as the *set of vertices* and $E \subseteq [V]^2$ serving as the *set of edges*. Such a graph could be interpreted as a matrix $A = [a_{ij}] \in \{0, 1\}^{n \times n}$ with $\{v_i, v_j\} \in E \Leftrightarrow a_{ij} = 1$; such a representation is often called the *adjacency matrix*. The key property of such matrices is that they are invariant with respect to entrywise multiplication. In turn we can use this property to generalise graphs on sets to graphs on quantum sets.

The following definitions are due to Schäfer i Tobolski [5].

Definition 3.2.1. (Quantum adjacency matrix, Schäfer i Tobolski [5, Definition 2.5]) Let (B, ψ) be a quantum set. A *quantum adjacency matrix* on this quantum set is a linear map $A: B \rightarrow B$ which is Schur-idempotent and $*$ -preserving, by which we mean that it satisfies

$$m(A \otimes A)m^* = A, \quad A(b^*) = A(b)^* \quad \text{for all } b \in B.$$

Definition 3.2.2. (Quantum graph, Schäfer i Tobolski [5, Definition 2.5]) A *quantum graph* on (B, ψ) is given by a quantum adjacency matrix on this quantum set.

For all intents and purposes we will identify both notions and simply call an appropriate $A: B \rightarrow B$ both a quantum adjacency matrix and a quantum graph.

Remark 3.2.1. There are other equivalent definitions of a quantum graph, namely those based on

- an *edge projection*, i.e. a projection $\tilde{A} \in C(X) \otimes C(X)^{\text{op}}$, see Musto, Reutter i Verdon [4],
-

Definition 3.2.3. (Schäfer i Tobolski [5, Definition 4.6]) Let A be a quantum graph on a quantum set (B, ψ) . We call the quantum graph

- *loopfree* if $m(A \otimes \text{Id})m^* = 0$,
- *undirected* if $A = A^*$, where the adjoint is taken with respect to the inner product induced by ψ .

We say that a voltage quantum graph $\tilde{A} = \{\tilde{A}_g\}_{g \in G}$ has these properties if the same equations hold for $\sum_{g \in G} \tilde{A}_g$, respectively.

3.3 Voltage and Derived Quantum Graphs

3.3.1 Classical Voltage Graphs

Definition 3.3.1 (Classical voltage graph). A *voltage graph* is a pair (Γ, λ) consisting of (not necessarily simple) finite directed graph Γ with vertex set V and edge set E , together with a labelling $\lambda: E \rightarrow G$ of the edges of Γ by elements of a finite group G .

Definition 3.3.2 (Classical derived graph). The *derived graph* Γ^λ of (Γ, λ) is a graph with vertex set $V \times G$ and edge set $E \times G$. Whenever $e \in E$ is an edge from u to v in Γ and $g \in G$ is a group element, then the edge (e, g) goes from (u, g) to $(v, g\lambda(e))$ in Γ^λ .




Figure 3.1: Base voltage graph Γ over $\mathbb{Z}_2$




Figure 3.2: Derived graph Γ^λ (Directed 4-cycle)




Figure 3.3: Base voltage graph Γ over $\mathbb{Z}_3$




Figure 3.4: Derived graph Γ^λ (Two 3-cycles)




Figure 3.5: Base Theta graph Γ with edge labels




Figure 3.6: Derived graph Γ^λ ($K_{3,3}$ colored by edge class)

3.3.2 Generalisation to quantum voltage and derived quantum graphs

In the following definition we will introduce some key notions. The motivation behind the definition of X_g will be given later in the thesis.

Definition 3.3.3. (Schäfer i Tobolski [5], Definition 4.1) Let $(\tilde{B}, \tilde{\psi})$, $\hat{\alpha} : \hat{G} \rightarrow \text{Aut}(\tilde{B}, \tilde{\psi})$, $(B, \psi_{\rtimes})$ and $(u_{\chi})_{\chi \in \hat{G}} \subset B$ be as above.

1. For every $g \in G$ let $X_g := \frac{1}{n} \sum_{\chi \in \hat{G}} \overline{\chi(g)} u_{\chi} u_{\chi}^{\dagger}$, i.e.

$$X_g : B \rightarrow B, \quad b \mapsto \frac{1}{n} \sum_{\chi \in \hat{G}} \overline{\chi(g)} \langle u_{\chi}, b \rangle_{\psi_{\rtimes}} u_{\chi}.$$

2. A *voltage quantum graph* on the quantum set $(\tilde{B}, \tilde{\psi})$ with respect to the group action $\hat{\alpha} : \hat{G} \rightarrow \text{Aut}(\tilde{B}, \tilde{\psi})$ is a family $\tilde{A} = (\tilde{A}_g)_{g \in G}$ of quantum adjacency matrices that is indexed by G and satisfies

$$\tilde{A}_g \hat{\alpha}_{\chi} = \hat{\alpha}_{\chi} \tilde{A}_g$$

for all $g \in G$ and $\chi \in \hat{G}$.

3. The *derived quantum graph* $\tilde{A} \rtimes_{\hat{\alpha}} \hat{G} : B \rightarrow B$ on $(B, \psi_{\rtimes})$ is then given by

$$\tilde{A} \rtimes_{\hat{\alpha}} \hat{G} := \sum_{g \in G} m(X_g \otimes E_{\rtimes}^* \tilde{A}_g E_{\rtimes}) m^*,$$

where we write (here and in the rest of the section) m for the multiplication map on B , i.e. $m = m_{\rtimes}$.

One can easily check that X_g is a adjacency matrix for any $g \in G$.

Lemma 3.3.1. *If $A, B \in M_n(\{0, 1\})$ and $m^* : \mathbb{C}^n \rightarrow \mathbb{C}^n \otimes \mathbb{C}^n$ is the adjoint of the multiplication map m , then $m(A \otimes B) m^* = A \odot B$, the entrywise (Hadamard) product of A and B .*

Proof. Let $\{e_1, \dots, e_n\}$ be the standard basis of $\mathbb{C}^n$. The componentwise multiplication map m acts as:

$$m(e_i \otimes e_j) = \delta_{ij} e_i$$

By definition of the adjoint via the standard inner product, $\langle m(e_i \otimes e_j), e_k \rangle = \langle e_i \otimes e_j, m^*(e_k) \rangle$. Since the left-hand side is $\delta_{ij} \delta_{ik}$, it follows that:

$$m^*(e_k) = e_k \otimes e_k$$

We expand the action of the matrices A and B on a basis vector e_k via their entries A_{ik}, B_{jk} :

$$Ae_k = \sum_{i=1}^n A_{ik} e_i, \quad Be_k = \sum_{j=1}^n B_{jk} e_j$$

Applying the composite operator $m(A \otimes B)m^*$ to e_k yields:

$$\begin{aligned}
m(A \otimes B)m^*(e_k) &= m(Ae_k \otimes Be_k) \\
&= m\left(\sum_{i=1}^n \sum_{j=1}^n A_{ik}B_{jk}(e_i \otimes e_j)\right) \\
&= \sum_{i=1}^n \sum_{j=1}^n A_{ik}B_{jk}\delta_{ij}e_i \\
&= \sum_{i=1}^n (A_{ik}B_{ik})e_i
\end{aligned}$$

Thus, the (i, k) -th entry of the matrix $m(A \otimes B)m^*$ is exactly $A_{ik}B_{ik}$. Because A and B are binary, this entry evaluates to 1 if and only if $A_{ik} = 1$ and $B_{ik} = 1$, and 0 otherwise. $\square$

Remark 3.3.1. $m(A \otimes B)m^*$ can be thought of as taking the intersecion of two quantum matrices blabla...

3.4 Isomorphisms and Quantum Isomorphisms of Quantum Sets

Definition 3.4.1. (Quantum isomorphisms, Schäfer i Tobolski [5, Definition 2.8]) Let (B_1, ψ_1) and (B_2, ψ_2) be quantum sets and let $A_1: B_1 \rightarrow B_1$ and $A_2: B_2 \rightarrow B_2$ be quantum graphs on (B_1, ψ_1) , (B_2, ψ_2) , respectively.

- The two quantum sets are *isomorphic* if there is a $*$ -isomorphism $\phi: B_1 \rightarrow B_2$ such that $\psi_2\phi = \psi_1$. We denote by $\text{Aut}(B, \psi)$ the group of quantum set automorphisms of a quantum set (B, ψ) .
- The two quantum graphs are *isomorphic* if there is an isomorphism $\phi: B_1 \rightarrow B_2$ of the underlying quantum sets such that $\phi A_1 = A_2 \phi$. We denote by $\text{Aut}(A)$ the group of quantum graph automorphisms of a quantum graph A .
- The two quantum sets are *quantum isomorphic*, written $(B_1, \psi_1) \cong (B_2, \psi_2)$, if there is a finite-dimensional Hilbert space $\mathcal{H}$ and a unital $*$ -homomorphism

$$\rho: B_1 \rightarrow B_2 \otimes B(\mathcal{H}),$$

such that the map

$$p: L^2(B_1) \otimes \mathcal{H} \rightarrow L^2(B_2) \otimes \mathcal{H}, \quad a \otimes \xi \mapsto \rho(a)(1 \otimes \xi)$$

is unitary as an operator between Hilbert spaces.

- The two quantum graphs are *quantum isomorphic*, written $A_1 \cong_q A_2$, if there is a quantum isomorphism $\rho: B_1 \rightarrow B_2 \otimes B(\mathcal{H})$ between the underlying quantum sets such that

$$\rho A_1 = (A_2 \otimes \text{Id}_{B(\mathcal{H})})\rho$$

holds.

Chapter 4

Quantum graphs over 3×3 matrices

The classification on 2×2 matrices was done by Matsuda [3].

4.1 Gromada's theorem

In the following section we will state...

The following theorem is due to Gromada [2].

Theorem 4.1.1 (Gromada [2], Theorem 3.7). *Consider $m \in \{0, 1, \dots, n^2\}$. Quantum graphs on M_n with m quantum edges are in a one-to-one correspondence with m -dimensional subspaces $V \subseteq \mathfrak{gl}_n$. The adjacency matrix is given by $A_V = \sum_{s=1}^m \xi_s \otimes \xi_s^*$, where $(\xi_1, \xi_2, \dots, \xi_m)$ is some orthonormal basis of V .*

The quantum graph has no loops if and only if $V \subseteq \mathfrak{sl}_n$. It is undirected if and only if $V = V^$.*

Proof. See Gromada's paper for reference. □

Lemma 4.1.1 (Gromada [2], Lemma 3.8). *Any vector subspace $V \subseteq \mathfrak{gl}_n$ with $V = V^*$ has a self-adjoint basis $\{\xi_s\}$, $\xi_s = \xi_s^*$.*

Proof. We construct the basis by induction. Suppose we already have a basis $\{\xi_s\}$ of a subspace $V_1 \subseteq V$ such that $\xi_s = \xi_s^*$. This means that $V_1 = V_1^*$. Let V_2 be complement to V_1 . Take any $\xi \in V_2$ and consider $(\xi + \xi^*)/2$ and $i(\xi^* - \xi)/2$. Both elements are self-adjoint and elements of V_2 . They might or might not be linearly independent, but definitely at least one of them is non-zero. So, include both (if they are linearly independent) or one of them (if they are not) to our basis $\{\xi_s\}$ and repeat. □

Remark 4.1.1. The following lemma tells us that any undirected graph on M_n can be thought of as being made from symmetric quantum edges only.

Remark 4.1.2. A complex vector space $V = V^* \subseteq \mathfrak{gl}_n$ uniquely determines a real vector space spanned by the basis $\{\xi_s\}$ of self-adjoint elements $\xi_s = \xi_s^*$. Indeed, it is easy to see that any other self-adjoint element $\xi = \xi^*$ must be a real linear combination of $\{\xi_s\}$.

Bibliografia

- [1] B. Blackadar. *Operator Algebras: Theory of C^* -Algebras and von Neumann Algebras*. Encyclopaedia of Mathematical Sciences. Springer Berlin Heidelberg, 2006. ISBN: 9783540285175. URL: <https://books.google.pl/books?id=7-6MZLdRfdAC>.
- [2] Daniel Gromada. “Some examples of quantum graphs”. W: *Letters in Mathematical Physics* 112.6 (list. 2022). ISSN: 1573-0530. DOI: 10.1007/s11005-022-01603-5. URL: <http://dx.doi.org/10.1007/s11005-022-01603-5>.
- [3] Junichiro Matsuda. “Classification of quantum graphs on $\mathbb{N}^2$ and their quantum automorphism groups”. W: *Journal of Mathematical Physics* 63.9 (2022). ISSN: 1089-7658. DOI: 10.1063/5.0081059. URL: <http://dx.doi.org/10.1063/5.0081059>.
- [4] Benjamin Musto, David Reutter i Dominic Verdon. “A compositional approach to quantum functions”. W: *Journal of Mathematical Physics* 59.8 (sierp. 2018). ISSN: 1089-7658. DOI: 10.1063/1.5020566. URL: <http://dx.doi.org/10.1063/1.5020566>.
- [5] Björn Schäfer i Mariusz Tobolski. “Voltage Quantum Graphs and a Gross–Tucker Theorem for Quantum Graphs”. W: (2026). arXiv: 2602.20996 [math.OA].