

Jet Measurements at 8 TeV with the ALICE Experiment.

A Dissertation Presented for the

Doctor of Philosophy

Degree

The University of Tennessee, Knoxville

Andrew John Castro

September 2019

© by Andrew John Castro, 2019

All Rights Reserved.

“Most people are not looking for provable truths. As you said, truth is often accompanied by intense pain, and almost no one is looking for painful truths.”

— Haruki Murakami

Acknowledgments

My time in Tennessee has been some of the most enriching, fun, and frustrating time in my life. Through my time here I have had one steady fast pillar that supported me through it all, Megan. I could never thank you nor love you enough.

To my family. I would like to thank my father, Jaime. For shaping me into the man I have become and for the long hikes and backpacking trips that we've done together. He helped curate my love and passion for being in the outdoors. I would like to give a very special thanks to my sister Ali, her husband Adam, to my amazing nieces Amara, Amelia, and my equally amazing nephew Alyas. I would also like to thank Nancy, Jim, and Lindsey along with all the members of the Carr and Stobie families for making me feel like a part of them.

Thank you to my friend Saul. For all the great memories we've had growing up together, for the crazy trips to India and Mexico, and for taking a Grey Hound from San Francisco to Knoxville to visit me.

I would like to deeply thank the heavy ion group at the University of Tennessee for the support they gave me and the guidance. I would like to Soren Sorensen and Christine

Nattrass for their guidance. To Rebecca and Kyle for becoming life long friends. To Pat, Redmer, and Adam for giving me input and trouble shooting on the analysis.

I would also like to thank the members of my committee; Christine Nattrass, Ken Read, Soren Sorensen, and Ivan Maldonado.

Abstract

In nuclear collisions a deconfined stated of partons, quarks and gluons, interact creating a near ‘perfect’ fluid called the Quark–gluon plasma(QGP). As this state of matter expands and cools the quarks once again recombine into the hadrons measured in particle detectors. Understanding the energy loss mechanisms of this phase of matter is one of the major goals of the nuclear physics program at the Large Hadron Collider(LHC). The first principle processes that govern hadronization are not understood from a theoretical Quantum Chromodynamic(QCD) framework but are well described phenomenologically using Monte Carlo simulations. As partons interact with one another they will fragment into collimated sparys of particles known as jets. The topologies and properties of hadronic jets measured in collider experiment can be correlated to the hadrionzation phase in nuclear collisons. Measuring jet cross sections as a function of the jet radius in high energy experiments can constrain different hadronization models from one another and allow for more precise measurements of the QGP. Inclusive jet cross sections and ratios of jet cross sections are measured in this thesis using the 2012 proton–proton data collected at $\sqrt{s} = 8$ TeV and an integrated luminosity of 9.1 pb^{-1} with the ALICE detector at the LHC. This thesis presents results of jets with radii from 0.1 to 0.5 and over a wide kinematic range between

10 GeV and 150 GeV using both minimum bias and single shower triggered data from the ALICE Electromagnetic Calorimeter. The cross sections are corrected for detector effects by unfolding and results compared to Monte Carlo simulations using PYTHIA and PHOJET tuned with different hadronization models.

Table of Contents

1	Introduction	1
2	Quantum Chromodynamics	4
2.1	The QCD Lagrangian	4
2.2	Jets	5
2.2.1	Jet Production and the Factorization Theorem	7
2.3	Jet Finding Algorithms	16
2.3.1	Cone Algorithms	17
2.3.2	Sequential/Recombination Algorithms	18
2.3.3	FastJet	21
2.4	Monte-Carlo Generators	22
2.5	The Quark-Gluon Plasma	25
2.5.1	Nuclear Collisions	25
2.5.2	Jets as a Probe of the QGP	27
2.6	QGP in Proton-Proton Collisions?	29
3	The LHC and ALICE	31

3.1	Overview of the LHC	31
3.1.1	LHC Operations	32
3.1.2	LHC Accelerator Complex	33
3.2	The ALICE Experiment	34
3.2.1	TZERO	36
3.2.2	VZERO	36
3.2.3	Inner Tracking System	37
3.2.4	Time Projection Chamber	38
3.2.5	Electromagnetic Calorimeter	41
4	The ALICE TPC Upgrade	46
4.0.1	Physics Motivation	46
4.0.2	Gaseous Electron Multiplier Foils	49
4.0.3	Research and Development	51
4.0.4	Production of the Inner Readout Chambers in the U.S.	55
4.0.5	GEM and Chamber Quality Assurance	57
4.0.6	Outlook	61
5	Analysis Methods and Corrections	63
5.1	Raw Jet Spectra	65
5.2	8 TeV Data Quality	65
5.2.1	EMCal Cluster Selection	69
5.2.2	TPC Track Selection	73
5.3	Jet Selection	75

5.3.1	$z_{leading}$ cut	76
5.3.2	Jet Area Cut	77
5.3.3	NEF cut	77
5.4	EMCal Triggered Data	78
5.5	Acceptance Correction	81
5.6	Unfolding	82
5.6.1	Response Matrix	83
5.6.2	Corrections to Particle Level	84
5.6.3	Unfolded MB Spectra	85
5.6.4	Unfolded EMCal Triggered Spectra	86
5.6.5	Jet Reconstruction and Matching Efficiency	88
6	Conclusion and Outlook	91
6.1	Systematic Uncertainties	91
6.1.1	Systematic Uncertainty to Jet Energy Scale	92
6.1.2	Systematic Uncertainty to Jet Yield	95
6.2	8 TeV Inclusive Jet Results from CMS and ATLAS	98
6.3	Inclusive Jet Spectra and Cross Section Ratios at 2.76 TeV	99
Bibliography		103
Appendices		118
Vita		119

List of Tables

5.1	2012 8 TeV data taking period.	67
5.2	EMCal jet acceptance for radii 0.1 - 0.5.	81

List of Figures

1.1	The fundamental particles of the Standard Model[1].	2
2.1	Strong coupling constant (α_s) as a function of the momentum transfer (Q)[2].	6
2.2	Diagram showing a jet created by two partons under going a hard scattering, forming into hadrons, and detected in a calorimeter[3].	6
2.3	Timeline of a proton-proton collision. Starting from the bottom, two partons confined within the colliding protons have a hard interaction. The outgoing partons will induce partonic showers by radiating quarks and gluons. The partonic showers will eventually form into final state hadrons due to confinement which are measured in high energy experiments[4].	8
2.4	Proton PDF at $Q^2 = 10$ GeV (left) and $Q^2 = 10$ TeV (right) from the NNPDF Collaboration[5].	11
2.5	Lowest order quark-antiquark annihilation to top-antitop pair[6].	12
2.6	Parton cascade in a hadronic collision[7].	13
2.7	$u\bar{d}$ generating a $d\bar{d}$ pair via string breaking which will form color neutral hadrons, black lines show the string like equipotentials.[8].	14
2.8	Fragmentation Functions from e^+e^- and DIS experiments with fits[9].	15

2.9	Cartoon showing Collinear and Infrared safe jet candidates[10].	16
2.10	Lego plot of all four jet finders used on a single event with $R = 1$ jet radius[11].	20
2.11	The QCD phase diagram[12].	26
2.12	Comparison between a proton proton collision with no medium and a heavy ion collision with a colored medium stage[13].	27
2.13	Jet energy loss in a QCD medium[12].	28
2.14	Jet R_{AA} at 5.02 TeV with the ATLAS experiment[14].	29
3.1	LHC accelerator complex. The four main experiments are shown in their relative locations[15].	32
3.2	The ALICE Detector at CERN[16].	35
3.3	ALICE tracker, multiplicity, timing, and vertex detectors located near the interaction point[16].	35
3.4	Multiplicity measured in the V0 detector with Glauber fits corresponding to centrality[17].	37
3.5	The ALICE Time Projection Chamber[18].	38
3.6	The TPC readout region[19].	39
3.7	TPC momentum and tracking resolution[20].	41
3.8	ALICE EMCal along with super modules, tower strips, and towers[21].	42
3.9	Energy resolution in the EMCal measured in a 2007 test beam at CERN(blue) compared to GEANT3 simulations of the EMCal(orange), fits for the parameters A, B, and C are also shown[22].	43

3.10 Cluster Spectra from the ALICE EMCal. MinBias is shown in black while the red and blue points show the spectra using the gamma trigger at two energy thresholds[23].	45
4.1 Simulation of the invariant mass spectra for dileptons in a typical heavy-ion run with current ALICE performance (left) and after upgrade of ALICE for Run-3 (right) in PbPb at $\sqrt{s_{NN}} = 5.5$ TeV. The dilepton yields originating from the QGP are shown (red and orange), along with background contributions from light-hadrons (blue), and charm (magenta)[24].	48
4.2 Scanning electron microscope image of a GEM foil from top (left) and profile (right)[25].	49
4.3 Profile of GEM with electric-field lines and gradients (left). Simulation of an ionization electron (yellow line) entering a GEM from a drift volume, amplification electrons (green dots, yellow lines) and back flow ions (red lines) are created (right)[26].	50
4.4 ITS-TPC matching (<i>left</i>) and inverse momentum resolution (<i>right</i>) for a 4-GEM stack simulated in Garfield++ [27].	52
4.5 Final design of the upgraded readout chambers with a stack of 4 GEMS [28].	52
4.6 dE/dx resolution of the 4-GEM IROC prototype(<i>left</i>) and the separation power between electrons and pions as a function of gain (<i>right</i>)[29].	54
4.7 Energy resolution of the iron peak as measured from the prototype IROC with varying GEM voltages as a function of IBF%[29].	54

4.8	Simulation of the four GEM (blue) layers after test beam. The configuration is such that the two GEMs closest to the drift volume, right side, absorb the amplification ions created by the two GEMs closest to the readout (right) [29].	55
4.9	Production flow of the IROCs (red), OROCs (blue), and GEM foils (green)[27].	56
4.10	The author assembling an Inner Readout Chamber at the University of Tennessee.	57
4.11	Schematic for the setup of the GEM foil spark test (<i>left</i>)[30] and the GEM mounted in the HV gas box (<i>right</i>).	58
4.12	Schematic of the gas tightness testing setup at the University of Tennessee (<i>Courtesy of Joseph Rasson</i>).	59
4.13	Leak rate of the 47 chambers built at Tennessee with the maximum failure rate at 0.25 ml/hr shown (<i>Courtesy of Charles Hughes</i>).	60
4.14	The author testing spark testing chambers next to the LHC beam line (<i>left</i>) and real time output from the spark test during a live beam (<i>right</i>).	61
5.1	LHC state during the 8 TeV run.	63
5.2	Raw inclusive $R = 0.2$ jet spectra from the 8 TeV Min Bias and EMCAL triggered data	65
5.3	Raw inclusive $R = 0.3$ jet spectra from the 8 TeV Min Bias and EMCAL triggered data	66
5.4	Raw inclusive $R = 0.4$ jet spectra from the 8 TeV Min Bias and EMCAL triggered data	67

5.5	Min Bias event rejection summary.	69
5.6	Vertex displacement from primary interaction point for accepted Min Bias events.	70
5.7	EMCal cell occupancy after bad channels removed.	71
5.8	Matched track-cluster distance.	71
5.9	EMCal cluster time distribution before cuts.	72
5.10	Corrected EMCal cluster yield.	73
5.11	Hybrid Track η and ϕ yields.	74
5.12	Accepted hybrid track resolution.	74
5.13	Accepted track p_T yield.	75
5.14	$R = 0.2$ leading track p_T per jet p_T .	75
5.15	$R = 0.2$ number of constituents in a jet per jet p_T .	76
5.16	$R = 0.2$ $z_{leading}$ from the Min Bias data sample.	76
5.17	$R = 0.2$ NEF per jet P_T .	78
5.18	Distance to closest reconstructed EJE patch to $R = 0.2$ jet with the Min Bias Pythia Monte Carlo.	79
5.19	Distance to closest reconstructed EGA patch to $R = 0.2$ jet with the Min Bias Pythia Monte Carlo.	80
5.20	Response Matrices for $R = 0.2$, $R=0.3$, and $R = 0.4$ jets.	84
5.21	Unfolded jet spectra with fine binning for $R = 0.3$	85
5.22	Unfolded Min Bias $R = 0.2$ jet spectra with corrections factors using a variable binning.	86

5.23 Unfolded Min Bias $R = 0.3$ jet spectra with corrections factors using a variable binning.	87
5.24 Unfolded Min Bias $R = 0.4$ jet spectra with corrections factors using a variable binning.	88
5.25 Unfolded EMCAL triggered $R = 0.2$ jet spectra with corrections factors using a variable binning.	89
5.26 Unfolded EMCAL triggered $R = 0.3$ jet spectra with corrections factors using a variable binning.	89
5.27 Unfolded EMCAL triggered $R = 0.4$ jet spectra with corrections factors using a variable binning.	90
6.1 Systematic due to TPC tracking efficiency.	93
6.2 Systematic due to Hadronic correction.	93
6.3 Systematic due to clusterization algorithm.	94
6.4 Inclusive track resolution, Min Bias 8 TeV.	95
6.5 Systematic due to P_T resolution.	96
6.6 Systematic due to energy resolution.	96
6.7 8 TeV CMS inclusive jet cross sections with radii of $R = 0.7$ and binned by jet rapidity compared to NLO calculations with non-perturbative corrections[31].	99
6.8 $R = 0.4$ inclusive jet cross section at 8 TeV from ATLAS in binned by jet rapidity compared to NLO QCD predictions[32].	100
6.9 The 8 TeV ATLAS jet cross sections rescaled to better show comparisons with NLO and non-perturbative calculations at low p_T [32].	100

6.10 Inclusive differential cross section from the 2.76 TeV proton proton run with ALICE	101
6.11 LHC state during the 8 TeV run.	101
6.12 Mimimum Bias Event Rejection	102
6.13 Emcal Triggered Event Rejection	102

Chapter 1

Introduction

From the Vedas to the ancient Greeks, generations have described the constituents of nature in terms of indivisible ‘elements’. It wasn’t until the beginning of the 20th century that the ancient elements of earth, wind, fire, water, and the aether were abandoned for the atomic theory of nature. By the 1960’s, what would become known as the Standard Model of Particle Physics was taking shape. The five ancient elements were replaced by the fundamental particles: quarks and leptons comprising the spin 1/2 fermions and the force carrying bosons with spin 1 as seen in Fig. 1.1

The Standard Model is the unification of the three symmetry groups, $SU(3) \times SU(2) \times U(1)$, representing the strong, weak, and electromagnetic forces[33]. In terms of scientific accomplishments the Standard Model is one of the most tested theories of nature with an agreement between the theory and observed results up to ten digits[34].

Even though the Standard Model gives us a deep understanding for many natural phenomena and has a wide range of uses; understanding the evolution of the Big Bang, how atoms and molecules bond, the nature of light, to cancer treatments and nuclear security it




Figure 1.1: The fundamental particles of the Standard Model[1].

is fundamentally an incomplete theory of nature. The fact that Gravity has yet to be unified into a quantum theory tells us that the Standard Model is incomplete. High energy experiments give us some of the most extreme conditions possible to test the Standard Model and to look for phenomena outside of the theory. Are their new symmetries and laws that manifest at high energies? Can we create dark matter or dark energy in a laboratory? Are quarks and leptons fundamental or finite in size? Do the four fundamental forces emerge from some yet unknown unified force? And why is antimatter absent in the Universe? All of these open questions are of great interest and currently form large areas of active research.

The theory of strong interactions, Quantum Chromodynamics(QCD), is described by the SU(3) group and similar in analogy to the electric charge of Quantum Electrodynamics (QED) carries color charge, red, green, and blue. Quarks and gluons are colloquially known as partons and are particles that interact via the strong force. At low energies and over large length scales partons are confined to a color neutral state and these particles must clump together into color neutral hadrons. As two colored partons began to separate, at some point it becomes energetically favorable to create a quark–antiquark pair out of the vacuum

rather than expanding the distance between neighboring partons. Due to confinement, quark interactions at high energy collider experiments manifest themselves as a spray of hadrons known as a 'jet'. The other main attribute of QCD is asymptotic freedom, as the interactions between partons becomes more energetic and the length scale decreases the strong coupling constant becomes exceedingly small, $\alpha_{strong} \ll 1$, and the partons freely interact with one another. Due to asymptotic freedom nuclear matter undergoes a phase transition called the Quark–Gluon Plasma (QGP) at high energies and densities

This thesis will present an overview of Quantum Chromodynamics in Chapter 2, with an emphasis on jet physics and heavy ion collisions. Chapter 3 will give a brief overview of the Large Hadron Collider and the ALICE experiment, including the relevant subsystems for this jet analysis. Jet results from the ALICE detector along with comparisons to QCD simulations with systematic uncertainty calculations and unfolding for detector effects will be given in Chapter 4. The ALICE time projection chamber will be upgraded to a continuous readout mode, the physics motivation behind this and author's contributions will be discussed in Chapter 5. Chapter 6 will serve as a final discussion, outlook, and conclusion.

Chapter 2

Quantum Chromodynamics

In 1968 deep inelastic scatterings performed at the Stanford Linear Accelerator Center showed that the proton had internal structure[35] called partons at the time. Within a decade of this discovery the partons were broken into two categories: the mass carrying fermions were known as the quarks and the gauge boson force carriers were called gluons. The interactions of these two types of particles were described by the quantum field theory known as Quantum Chromodynamics (QCD) and by the SU(3) symmetry group. SU(3) guarantees that color charge is conserved and this results in quarks grouping together into ‘colorless’ hadrons.

2.1 The QCD Lagrangian

QCD is the strongest of the known fundamental forces. It is a gauge field theory described by the Lagrangian density

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}^\alpha F_\alpha^{\mu\nu} - \alpha_s(\bar{q}_j \gamma^\mu T_\alpha q_j) G_\alpha^\mu + \bar{q}_j(i\gamma^\mu \partial_\mu - m)q_j \quad (2.1)$$

where q and \bar{q} represent the color anti-color fields summed over color j , α_s is the strong coupling strength, γ^μ is the Dirac gamma matrix, G_α^μ is the gauge field for color α , is similar in analogy to the \mathbf{W} matrix from the electroweak theory. $F_{\mu\nu}^\alpha$ is the field strength tensor and it describes the gluon interactions. The first term of the Lagrangian is the gluon contribution and carries no mass term. The second term of the Lagrangian describes how quarks and gluons interact with each other. The final term describes quark interactions and the coupling between them and will be explored further in this thesis.

At short distances, less than 0.2 fm , the strong coupling constant becomes exceedingly small and second term of the Lagrangian displays an important property known as asymptotic freedom[36].

$$\alpha_s = \frac{1}{\beta_0 \ln(Q^2/\Lambda^2)} \quad (2.2)$$

where α_s is the strong coupling constant, Q^2 is the momentum transfer between two interacting partons, and Λ^2 is a cutoff below which QCD phenomena are strongly suppressed and β_0 is a correction factor.

2.2 Jets

Hard probes (large Q^2 interactions), are produced in the earliest stages of a high energy collision when the largest momentum transfer processes occur. As two highly energetic




Figure 2.1: Strong coupling constant (α_s) as a function of the momentum transfer (Q)[2].

partons propagate away from one another, in a back-to-back fashion, they will instigate a shower of daughter partons via gluon radiation and the generation of low-mass $q\bar{q}$ pairs. These daughter partons will go on to form hadrons and the clustering of these hadrons is colloquially known as a ‘jet’. If the jet was created in a high energy experiment the final state hadrons will be recorded as tracks in a tracking detector or energy deposits in a calorimeter.

This process is shown in Figure 2.2.




Figure 2.2: Diagram showing a jet created by two partons undergoing a hard scattering, forming into hadrons, and detected in a calorimeter[3].

The physicist James Daniel Bjorken postulated that a correlation could be surmised by summing over the final state transverse momentum of the hadrons that form a jet to the parton that initiated the hard scattering[37][38]. This has lead to jets becoming the work-horse for both experimentalists and theorists over the past 30 years in probing QCD phenomena. This thesis makes use of jets as an important probe of QCD and the following sections are devoted to developing a background for both the theoretical and experimental treatment of jet physics. The following sections will be devoted to how jets are produced from a physics point-of-view and the latest results which this thesis compares to.

2.2.1 Jet Production and the Factorization Theorem

Due to confinement bare quarks are unobserved, therefore experimentalists must probe QCD interactions by detecting the color neutral final state hadrons measured in collider experiments. Fortunately, the factorization theorem (Equation 2.3) allows for the final state jet cross section to be broken into a number of steps that can either be calculated perturbatively using pQCD or modeled phenomenologically. Using the factorization theorem the jet cross section in a pp collision is,

$$d\sigma^{pp \rightarrow jet} \sim f_{a/A}(x_1, Q^2) \otimes f_{b/B}(x_2, Q^2) \otimes d\sigma_{ab \rightarrow c+X}(x_1, x_2) \otimes D_{c \rightarrow h/jet}(z, Q^2) \quad (2.3)$$

- $f_{a/A}(x_1, Q^2)$ and $f_{b/B}(x_2, Q^2)$ are the parton distribution functions (PDF) that describe the probability of finding parton, a or b , within nuclei, A and B , with a given momentum fraction, $x = p_{parton}/p_{hadron}$ as a function of Q^2 .

- $d\sigma_{ab \rightarrow c+X}(x_1, x_2)$ is the pQCD parton-parton cross section due to the hard scattering of the two partons, a and b , to an intermediate parton (c).
- $D_{c \rightarrow h/jet}(z, Q^2)$ is the fragmentation function (FF) that describes the probability the an outgoing parton, c , fragments and hadronizes into a final state hadron, h , within a jet with momentum fraction, $z \equiv p_{hadron}/p_{parton}$.




Figure 2.3: Timeline of a proton-proton collision. Starting from the bottom, two partons confined within the colliding protons have a hard interaction. The outgoing partons will induce partonic showers by radiating quarks and gluons. The partonic showers will eventually form into final state hadrons due to confinement which are measured in high energy experiments[4].

Figure 2.3 shows a timeline of a pp collision broken into the relevant steps in accordance to the factorization theorem. One of the best places to fundamentally test QCD phenomena using hard probes, i.e. jets, are with high energy hadron colliders such as those found at

CERN¹, Fermilab, and BNL. The time scale that a hard probe is created in a high energy collision is on the order of $\tau \approx 1/p_T \approx 0.1 \text{ fm}/c$ which probes the initial state these interactions. The factorization theorem allows for a high level of agreement between the QCD theory of nature and experimental observables but to ascertain this connection we should discuss each term of the factorization theorem in more depth.

Parton Distribution Functions

The PDF occurs twice in Equation 2.3 due to the two partons that will undergo the hard scattering being confined in two different protons. PDFs may be thought of as conveying the structure of a nucleon in terms of the number of flavored quarks or gluons ($u(x)$, $d(x)$, $s(x)$, $\bar{u}(x)$, $\bar{d}(x)$, $\bar{s}(x)$, $g(x)$) and must obey certain constraints and summation rules. In the case of a proton, with electric charge ($e = +1$),

$$+1 = \frac{2}{3} \int_0^1 [u(x) - \bar{u}(x)] dx - \frac{1}{3} \int_0^1 [d(x) - \bar{d}(x)] dx \quad (2.4)$$

and isospin ($I = 1/2$),

$$\frac{1}{2} = \frac{1}{2} \int_0^1 [u(x) - \bar{u}(x)] dx - \frac{1}{2} \int_0^1 [d(x) - \bar{d}(x)] dx \quad (2.5)$$

have a solution,

$$\int_0^1 [u(x) - \bar{u}(x)] = 2 \quad (2.6)$$

¹Discussed in detail in Chapter 3

$$\int_0^1 [d(x) - \bar{d}(x)] dx = 1 \quad (2.7)$$

This corresponds to the classical partonic view that protons contained two up quarks and a down quark, similarly the neutron, with charge $e = 0$ and isospin $I = -1/2$, can be shown that it compromises two down quarks and a up quark. Naively, we could assume that the three quarks composing a proton would each carry a momentum fraction of approximately $1/3$ the total momentum of a proton. However, high energy deep inelastic scattering experiments conducted at the Stanford Linear Collider in the 1960's[39] measured the momentum carried by the three quarks only accounting for about $1/2$ the total proton momentum. This lead to a more complex and dynamic model of the proton structure with the other half of the proton momentum being occupied by neutral partons which would eventually become known as gluons.

Measuring the structure of the partons making up a nucleon is a major endeavor by both theorists and experimentalists. Two of the most popular PDFs available to physicists are the CTEQ[40] (Coordinated Theoretical-Experimental Project on QCD) and the NNPDF[41] (Neural Network Parton Distribution Function) sets. Figure 2.4 shows the proton PDF as a function of the momentum fraction for two energy ranges, at high values of x the two up quarks account for about $2/3$ of the momentum fraction while down quark accounts for about $1/3$ the total momentum, these quarks are collectively called the valence quarks. At high energies, low values of x , we see that the proton has non negligible contributions from gluons, anti-quarks, strange, and even charm quarks, these are collectively known as the sea partons. Today, the modern picture of a protons structure is that it is mostly composed of




Figure 2.4: Proton PDF at $Q^2 = 10 \text{ GeV}$ (left) and $Q^2 = 10 \text{ TeV}$ (right) from the NNPDF Collaboration[5].

gluons and sea quarks at low values of x and this domination only increases as a function of Q^2 [42].

Parton-Parton Cross-Section

The parton-parton cross section can be calculated using perturbation theory. To the zeroth order in α_s this cross-section would be a simple quark-antiquark annihilation and would be calculable using Feynman diagrams as seen in Figure 2.5[43]. Higher ordered contributions, such as the creation of virtual gluons, require the hard cross-section to be expanded as a series in terms of α_s . Calculations of the hard cross section that incorporate these higher order terms are known as *next-to-leading order* (NLO) with N denoting the number of terms after the leading order that have been included in the cross-section calculation. Various

calculations of the hard cross-section of different QCD processes have been performed over the years typically using either power series or logarithmic expansions of α_s [44] and corrections for LO, NLO, and even NNLO constitutes a very active field in high energy physics.




Figure 2.5: Lowest order quark-antiquark annihilation to top-antitop pair[6].

Perturbative techniques of the hard cross-section have been extremely successfully in describing jet features in hadronic collisions[42].

Hadronization

Hadronization is the process by which the colored pQCD partons of form into colorless non-pQCD hadrons and represents a significant barrier in progressing jet physics. This is due to the fact that hadronization encompass several smaller processes which in themselves are hard to characterize. Thus, like PDFs, an accurate description of hadronization requires a phenomenological approach by which experimental results help complement theoretical calculations. Jet production via hadronization[7] follows two distinct stages. First, the partons that underwent a hard scattering start to emit radiation via gluon bremsstrahlung up until time, $t < Q^2$, this is known as the parton cascade. The parton cascade is the precursor of what will become a jet as most of the radiation generated will travel in the same direction as the initial hard scattered parton. However, this immediately poses an

issue in jet physics as radiation generated at a wide angle away from the momentum axis of the initial hard scattered parton will not be associated with the jet.




Figure 2.6: Parton cascade in a hadronic collision[7].

After the cascade has ended the partons must fragment into color neutral hadrons. There are two main phenomenological models used to describe the hadron forming process, the Lund String Model and the Cluster Hadronization Model. The QCD potential is,

$$V(r) = -\frac{\alpha_s}{r} + \sigma r \quad (2.8)$$

where the first term of Equation 2.8 goes as the Coulomb potential with a $1/r$ dependence and is the dominate term at short distance and the second term has a string-like potential with σ referring to a string-like tension. The Lund String Model ignores gluon radiation and has fragmentation occur via breaking the string tension with the production of $q\bar{q}$ sea quarks. The created sea quarks will carry some momentum fraction, z , of the initial parton until z falls below some cutoff. Figure 2.7 shows a two quarks undergoing a string breaking,

each of the quarks initiating the string breaking will combine with a sea quark in an iterative manner to form hadrons.




Figure 2.7: $u\bar{d}$ generating a $d\bar{d}$ pair via string breaking which will form color neutral hadrons, black lines show the string like equipotentials.[8].

The Cluster Hadronization Model has gluons splitting after the parton cascade phase into $q\bar{q}$ pairs. These pairs will form color-singlet clusters with other neighboring quarks in phase-space. These color-singlets will typically be a few GeV/c^2 in mass and are treated as excited meson resonances. These pseudo-resonances will decay via their normal branching ratios into the stable hadrons[45].

Fragmentation

Similar to how a PDF quantitatively describes the structure of a nucleon the FF quantitatively describes the hadronization process. The FF is also similar to the PDF in that it is also a probability distribution, thus it follows the probabilistic rule that,

$$\sum \int z D_{c \rightarrow h/\text{jet}}(z, Q^2) dz = 1 \quad (2.9)$$

Ideally, the fractional momentum of the hadrons created from the fragmenting parton, $z \equiv p_{\text{hadron}}/p_{\text{parton}}$. The Parton-Hadron Duality[46] states that the leading hadron should encompass the quantum numbers and kinematic properties associated with the hard

scattered quark that initiated the jet. Thus we can measure the fragmentation function as $z = p_{hadron}/p_{jet}$. The formulation of the FF as a fractional energy carried by the hadrons in a jet was a breakthrough in pQCD techniques and is analogous to how an electron passing through an absorber creates photon showers with these photons generating conversion electrons until the total energy has been dissipated into the material.




Figure 2.8: Fragmentation Functions from e^+e^- and DIS experiments with fits[9].

Figure 2.8 is the FF in terms of the Gaussian equation, $z dN/dz = -dN/d\xi$ with $\xi = -\ln 1/z$. The observation that the Gaussian peaks of Figure 2.8 along with the suppression of the FF at low z values due to gluon coherence were predicted by pQCD.

2.3 Jet Finding Algorithms

A jet arises from the fragmentation of a hard parton to final state hadrons. However, grouping the hadrons together into a jet is a non-trivial task and jet finding algorithms are deployed in order to achieve this objective. Early on in jet physics, both theorists and experimentalists used a wide variety of jet finders that made comparisons between experiments or to theoretical calculations nearly impossible[11]. For example, a radiated gluon that splits into a quark anti-quark pair may become one or two jets depending on the angular separation and the algorithm used. Early jet finders tended to be sensitive to soft particles or could give widely varying yields to the number of jets in an event. In 1990, the Snowmass Accord[47] reached a standardized definition of a jet between experimental and theorists. The agreement maintained that any algorithm that clusters particles into a jet must be both infrared and collinear safe (IRC).




Figure 2.9: Cartoon showing Collinear and Infrared safe jet candidates[10].

Collinear safety ensures that a high- p_T particle split into two or more particles should not influence the kinematics of a hard jet, this makes the jet finders insensitive to how hadrons are grouped together. Infrared safety in turn requires that the emission of soft radiation should not affect the properties of a jet, this makes jets returned by the algorithm

independent of soft physics and a true signature of a hard process. Both of these processes are shown in Figure 2.9. After the adoption of these standards from the Snowmass Accord, old algorithms that violated these rules were patched and new jet finders were developed to comply with IRC safety. The most prevalent jet finding algorithms today fall into two categories: cone algorithms and sequential recombination/clustering algorithms.

2.3.1 Cone Algorithms

Cone algorithms made up the bulk of early jet finders. The only IRC safe cone algorithm still in use today is the seedless infra-red safe cone algorithm (SIScone). SIScone defines a cone of radius, R , around the highest momentum particle in the coordinates of (η, ϕ) ², this is the proto-jet. SIScone then proceeds through an iterative process of finding all the particles within the jet radius such that $R \leq \sqrt{\phi^2 + \eta^2}$ and calculates a new jet center based on these particles momenta and a new weighted jet axis(η, ϕ). If the new center matches the proto-jet center, the proto-jet is tagged as a stable jet, all the particles in that jet are removed, and SIScone moves onto the next highest p_T particle. Cone algorithms tend to be unpopular due to being computationally expensive, they are hard to implement theoretically, and can give results not calculable in perturbation theory.

²It is possible to use a Cartesian coordinate system in particle colliders, with the z-component referring to points along the beam axis while the xy-plane is perpendicular to the beam axis. However, this system is not invariant under a Lorentz boost. Therefore it is more useful to use the cylindrical-like coordinates of pseudorapidity (η) and the azimuth angle (ϕ). Pseudorapidity may be thought of as the polar angle in a cylindrical coordinate system with $\eta = 0$ when the polar angle is perpendicular to the beam axis and $\eta = \inf$ along the beam axis. ϕ is the azimuth angle that rotates around the beam axis. Both, η and ϕ are invariant for Lorentz boosts along the beamline and allow for easy comparisons between the center-of-mass frame and the laboratory frame of a high energy collision.

2.3.2 Sequential/Recombination Algorithms

The other class of jet finders are the sequential/recombination algorithms, which are favored by experimentalists and theorists, and are IRC safe. There are three types of sequential/recombination algorithms: k_T , Anti- k_T , and the Cambridge/Aachen jet finders, with k_T referring to the component of a jet constituents momentum perpendicular to the jet axis. All of the algorithms use a similar method, first they find the distance between every pair of particles, $d_{i,j}$, such that

$$d_{i,j} = \min[p_{T,i}^a, p_{T,j}^a] \frac{\Delta_{ij}^2}{R^2} \quad (2.10)$$

where $p_{T,i}^a$ is the transverse momentum of particle i , a is free parameter that is set based on which algorithm is used, $\Delta_{ij}^2 = (\eta_i - \eta_j)^2 + (\phi_i + \phi_j)^2$ is the distance between the particles, and R is the radius of the jet. A second distance is defined in the sequential/recombination algorithm scheme,

$$d_{i,B} = p_{T,i}^a \quad (2.11)$$

this is the distance between a given particle i and the beam axis. Sequential/Recombination algorithms find the set of all particles, $d_{i,j}, d_{i,B}$, such that if $d_{i,B}$ is the minimum for particle i it is tagged as a jet and removed from the list. If $d_{i,j}$ are a minimum for particles i and j these two particles are merged together into a new particle (ij) and a new minimum is found between (ij) and particle k until all the particles are either merged into jets or the minimization function is no longer satisfied.

k_T Algorithm

The k_T algorithm sets the value a to 2, this results in a minimization function,

$$d_{i,j} = \min[p_{T,i}^2, p_{T,j}^2] \frac{\Delta_{ij}^2}{R^2} \quad (2.12)$$

which clusters low momentum particles first, making this algorithm susceptible to the underlying event (UE) or pile-up (PU). Thus the k_T algorithm is good at estimating any background present in a high energy collision.

Anti- k_T Algorithm

The Anti- k_T algorithm sets the value a to -2, resulting in a minimization function,

$$d_{i,j} = \min\left[\frac{1}{p_{T,i}^2}, \frac{1}{p_{T,j}^2}\right] \frac{\Delta_{ij}^2}{R^2}. \quad (2.13)$$

The minimization function is dominated by high- p_T particles, thus the area and axis of a jet is only slightly perturbed by soft particles. This makes the Anti- k_T algorithm robust in jet finding with events having a UE and PU. The Anti- k_T algorithm is the default jet finding algorithm used at the Large Hadron Collider and is the one used in this thesis.

Cambridge/Aachen Algorithm

The Cambridge/Aachen algorithm sets a to 0 and this results in a minimization function of,

$$d_{i,j} = \frac{\Delta_{ij}^2}{R^2} \quad (2.14)$$

which makes it independent of particle momentum and sensitive to PU and UE. Due to the fact that the Cambridge/Aachen algorithm is only dependent on the particle coordinate it is most useful in studying jet structure.




Figure 2.10: Lego plot of all four jet finders used on a single event with $R = 1$ jet radius[11].

Figure 2.10 shows the jets found in a single event using all four jet finding algorithms. It should be noted that the Cambridge/Aachen and k_T algorithms have highly irregular and large shapes, making them both susceptible to the presence of a UE, while SIScone finds an additional jet due to splitting. The Anti- k_T algorithm finds circular jets which demonstrates its robustness to hard radiation.

Once a stable jet is found, a recombination scheme is deployed in order to garner the jet kinematics. By adding the 4-vector, $\mathbf{p}^\mu = (E, \mathbf{p}_x, \mathbf{p}_y, \mathbf{p}_Z)$, for all of the associated particles composing a jet, we may obtain the jet momentum, energy, coordinates, etc³. In a particle collider with the tracks from a tracking detector measuring particle momentum and the towers of a calorimeter measuring particle energy we obtain the following relationships

$$p_T^{jet} = \sum_{particles} p_T = \sum_{tracks} p_T \quad (2.15)$$

$$E^{jet} = \sum_{particles} E = \sum_{towers} E \quad (2.16)$$

$$\eta^{jet} = \frac{1}{2} \ln \left(\frac{|\mathbf{p}^{jet}| + p_L^{jet}}{|\mathbf{p}^{jet}| - p_L^{jet}} \right) \quad (2.17)$$

$$\tan \phi^{jet} = \frac{p_y^{jet}}{p_x^{jet}} \quad (2.18)$$

where p_L refers to the longitudinal momentum which is the momentum component parallel to the beam axis. This method of adding the 4-vector of the particles composing the jet together in order to gain the jet kinematics is known as the E-scheme[48].

2.3.3 FastJet

FastJet[48] is a C++ software package that performs jet finding. Due to the computational efficiency, ease of use, and straight forward implementation, FastJet is the de-facto preferred

³For a review of relativistic kinematic see Appendix ...

jet finder used by theoreticians and all current high energy experiments. It implements the four previously discussed jet finders along with both the E-scheme and a boost invariant p_T scheme (BIpt-scheme) for recombination. The BIpt-scheme obtains the jet momentum and energy in the same manner as the E-scheme but uses a weighted average to find the jet coordinates,

$$\eta^{jet} = \sum_{particle} \frac{p_T^{particle}}{P_T^{jet}} \eta^{particle} \quad (2.19)$$

$$\phi^{jet} = \sum_{particle} \frac{p_T^{particle}}{P_T^{jet}} \phi^{particle} \quad (2.20)$$

In addition to basic jet measurements, FastJet contains a number of advance features, which allows it to be used to study jet area, jet substructure, and jet background subtraction[49].

2.4 Monte-Carlo Generators

Monte-Carlos (MC) allow for the simulation of high energy events on a statistical basis. Particle level generators use different phenomenological models of the factorization theorem in order to simulate the energy, momentum, particle species, multiplicity, and direction of travel expected in a high energy collision. In order to validate an analysis the particle level simulations are further propagated through a detector level simulation of an experiment, such as Geant3[50], in order to negate detector effects on the output observables from the MC simulation. In this section only the particle level simulations used in the thesis are discussed.

PYTHIA

PYTHIA[51], is a C++ Monte Carlo software tool-kit used to model proton-proton collisions. The package uses pre-defined parton distribution functions as input afterwards it simulates the partonic showers and radiation due to a hard scattering by generating the LO scattering matrix elements. Hadronization is performed in PYTHIA using the Lund String Model. After which relative branching ratios are used to statistically throw the decay modes of the hadrons until they are stabilized.

PYTHIA underestimates jet production due to the limitations of using LO calculations. Therefore, it uses an arbitrary value (K-factor) to make NLO corrections to the LO cross section. The K-factor is defined as,

$$K = \frac{\sigma_{NLO}}{\sigma_{LO}}. \quad (2.21)$$

NLO corrections to the cross-section will not match experimental results, PYTHIA implements additional phenomenological adjustments used to better match data. PYTHIA encompass these corrections into sets known as ‘tunes’, with PYTHIA 6.4 Perugia-2010 tune being used for this analysis[52].

PHOJET

PHOJET is a FORTRAN 77 Monte Carlo simulator used to model proton-proton collisions. It is an alternative to PYTHIA and is better at modeling soft physics processes present in high energy collisions. PHOJET implements the Dual Parton model[53][54] and multiple parton interactions[55] to model soft physics. Hard interactions are implemented in PHOJET

using LO scattering elements and it uses PYTHIA for the fragmentation and hadronization phase. Due to its ability to model soft physics, PHOJET is better at comparing to minimum bias⁴ data and understanding jet results in a low kinematic range. PHOJET also acts as a benchmark in understanding any bias due to using other MC generators, such as PYTHIA. PHOJET v1.2 is used in this thesis.

HERWIG

The Herwig[56] Monte Carlo generator is a FORTRAN 77 package used to generate proton proton events. It is similar to PYTHIA in that it does the LO hard scattering of partons, however it uses the cluster model of hadronization to produce jets based on gluon splitting. It is also similar to PHOJET in regards to the evolution of final state jets with soft gluon angular ordering. By comparing HERWIG, PYTHIA, and PHOJET it is possible to test for sensitivities to jet production in high energy events due to different types of hadronization models and soft radiation.

AliROOT

AliROOT is the official framework of C++ classes developed for the ALICE Experiment. The software is broken into two packages AliRoot, which contains code mostly used for the sub-detectors calibrations and reconstruction of particles, and AliPhysics which contains user defined classes to perform physics related analysis. This thesis developed a number of classes in both packages and are part of the offical repository available on GitHub.

⁴Events with a low total transverse momentum and high cross section

2.5 The Quark-Gluon Plasma

At the temperatures and pressures typical to the Universe today nuclear matter is confined to a colorless hadronic gas. However, it was theorized that at extreme temperatures, such as those experienced in the early Universe, partons would have undergone a phase transition where by they were no longer bound to a color neutral state. This state of matter would have been analogous to a conventional plasma where by the electrons are no longer bound to a nucleus, thus the state was dubbed the Quark-Gluon Plasma (QGP) The nuclear phase diagram is shown in Figure 2.11 as a function of temperature and the net baryon density. Normal nuclear matter is confined to the bottom left while increasing temperatures and/or densities correspond to the QGP. Modern particle colliders , such as RHIC and the LHC, are able to obtain the densities and temperatures necessary to create a QGP and are likewise shown in the figure. The reason for particle colliders being located at low baryon density is due to the fact that at high Q^2 energies more quark antiquark pairs are produce. This dilutes the total baryon density in the initial system and is more akin to what the early Universe was like.

2.5.1 Nuclear Collisions

By colliding heavy nuclei together in high energy collisions it is possible to obtain the energy densities and temperatures associated with the QGP state. The first signatures for the QGP were measured via a J/ψ suppression at the Super Proton Synchrotron located at CERN in 2000[57]. In 2005, the four experiments on the RHIC collider: BRAHMS[58], PHENIX[59], PHOBOS[60], and STAR[61], co-announced the observation of a new state of




Figure 2.11: The QCD phase diagram[12].

matter consistent with the hot and dense QGP. Surprisingly, the results from RHIC pointed at the QGP behaving more like a perfect fluid over a plasma-like state[62]. The QGP offers a rich and dynamic environment to test QCD predictions under some of the most extreme circumstances. Figure 2.12 shows the difference between a proton-proton collision and a heavy-ion collision. The heavy-ion collision mirrors the processes in a proton-proton collision (left) described in depth in Section 2.2.1. After the initial hard scattering the phase transition to a QGP occurs. The QGP undergoes a hydrodynamical evolution and expansion until it cools to a colorless hadronic gas. After the phase transition occurs unstable hadrons will decay until they are stable at which point the final state particle composition is set and the chemical freeze out occurs. The hadron gas expands and cools until all soft interactions cease, this is the kinetic freeze out, after which the final momentum spectra is set.

2.5.2 Jets as a Probe of the QGP

Jets are an excellent probe of the thermodynamic properties of the QGP. This is due to them being produced in the earliest stages, before the formation of the QGP, and surviving the full evolution of a heavy-ion collision. As a jet propagates through the QGP it will lose energy to the medium through a combination of gluon radiation to the colored medium and inelastic scatterings. These energy loss mechanisms are dependent on the distance a parton travels through the QGP and on the species of the parton.




Figure 2.12: Comparison between a proton-proton collision with no medium and a heavy ion collision with a colored medium stage[13].

Figure 2.13 shows two back-to-back partons undergoing a hard scattering. Both will fragment into jets, but the first parton with transverse energy, E_{T1} , will be subjected to much less energy loss over the second parton due to the first parton only traveling through the outer edge of the QGP. The species dependent parton energy loss arises from kinematic




Figure 2.13: Jet energy loss in a QCD medium[12].

constraints to gluon emission from the heaviest of quarks. This radiation is suppressed at angles smaller than the ratio of the quark mass to its energy and has been dubbed the *Dead-Cone Effect*. Tagging the flavor dependence of jets, either via measuring electrons from semi-leptonic decays or reconstructing the secondary vertex of heavy flavor mesons, has recently shown that energy loss via the Dead-Cone effect is strongly suppressed with jets containing a charm quark[63].

One way of quantifying the energy loss in a heavy-ion collision is via measurements of the nuclear modification factor, R_{AA} ,

$$R_{AA} = \frac{1}{N_{binary}} \frac{d^2 N_{AA}/dp_T d\eta}{d^2 N_{pp}/dp_T d\eta} \quad (2.22)$$

where N_{binary} is the number of binary collisions and is estimated using a Glauber model[64] of a nucleus while $d^2 N_{AA}/dp_T d\eta$ and $d^2 N_{pp}/dp_T d\eta$ are the spectra measured in nucleus nucleus and proton proton collision respectively. R_{AA} may be thought of as asking the question: Does a heavy-ion collision scale as a superposition of N_{binary} proton-proton collisions? If a heavy-ion collision simply looks like some N_{binary} factor of

Figure 2.13 shows the nuclear modification factor with $R = 0.4$ jets in the ATLAS experiment at 5.02 TeV[14]. The different colored bands in the figure are broken into centrality⁵




Figure 2.14: Jet R_{AA} at 5.02 TeV with the ATLAS experiment[14].

2.6 QGP in Proton-Proton Collisions?

As previously stated a QGP is believed to be absent in proton-proton collisions, thus any signature of a QGP should likewise be absent. However, one way of quantifying the presence of the QGP is via the Bjorken energy density.

$$\varepsilon = \frac{1}{\tau A} \frac{dE_T}{d\eta} \quad (2.23)$$

⁵The purple 10 - 20% band denotes the most central events (i.e. the two colliding nuclei have a low impact parameter and collide nearly head-on), while the 70 - 80% red band denotes the least central events (i.e. the two colliding nuclei have a high impact parameter and barely graze one another). An in depth discussion of centrality may be found here[65].

where A is the transverse area of the nuclei, τ is the proper time, and $dE_T/d\eta$ is the transverse energy per unit psuedorapidity. It can be shown that the 150 MeV critical temperature need for the phase transition to the QGP corresponds to $1 - 3 \text{ GeV}/fm^3$ energy density. The quantity $dE_T/d\eta$ can be related to the mean transverse momentum $\langle p_T \rangle$ and particle multiplicity⁶ per unity psuedorapidity as:

$$\frac{dE_t}{d\eta} \approx \langle p_T \rangle \frac{dN}{d\eta} \quad (2.24)$$

where $\langle p_T \rangle$ is the mean transverse momentum and $dN/d\eta$ is the particle multiplicity per unit psuedorapidity. This suggest that in very high multiplicity proton-proton events signatures of the QGP may be present. Although suppression has never been observed in high multiplicity proton-proton collisions physicists have recently measured collective flow in such systems[66]. CMS presented collective flow results in proton-proton collisions at 13 TeV using soft-particles, $p_T \leq 2 \text{ GeV}/c$, consistent with hydrodynamical predictions[67]. These results have opened new debates and questions into the very nature of the QGP and whether jets can be used to enlighten our understanding in such systems will make up a very active and interesting segment of high energy physics research in the coming years.

⁶Multiplicity is defined as the number of particles per event

Chapter 3

The LHC and ALICE

3.1 Overview of the LHC

The Large Hadron Collider (LHC)[\[68\]](#) is a circular particle accelerator located on the Franco-Swiss border near the city of Geneva. It is operated by the European Organization for Nuclear Research (CERN) and has carried out proton-proton (pp), lead-proton (pPb), and lead-lead (PbPb) collisions at center of mass energies of 0.9-14 TeV, 5.0 TeV, and 2.76-5.5 TeV, respectively. The LHC is approximately 17 miles in circumference and located 200 meters underground. It is located inside the old accelerator tunnel used by the Large Electron-Positron[\[69\]](#) collider of the 1980's. There are over 8000 physicists and engineers making up the four main experiments at the LHC: ATLAS[\[70\]](#), CMS[\[71\]](#), LHCb[\[72\]](#), and ALICE[\[73\]](#). Numerous physics results have been published, with the most famous being the discovery of the Higgs boson in 2012[\[74\]](#)[\[75\]](#).

Figure 3.1 shows a schematic of the LHC along with the pre-accelerators that help to accelerate protons and ions to their final energies before a collision at one of the four




Figure 3.1: LHC accelerator complex. The four main experiments are shown in their relative locations[15].

experimental interaction points(IP). Protons are injected into the LHC together in groups called ‘bunches’. Every bunch is comprised of about 120 billion protons with about 50 nanoseconds between the arrival of the next bunch. The bunch scheme during the heavy-ion run is reduced to 200 nanoseconds due to the high multiplicity of the events and additional computational resources needed.

3.1.1 LHC Operations

The LHC first attempted particle collisions in September of 2008. The initial ramping up of the super conducting magnets led to mechanical failure of the helium pipes inside of the LHC beam line. This fault caused the LHC to remain shut down for over a year while the accelerator was repaired and new safety procedures were implemented. The first successful collisions occurred in 2009 with proton collisions at a reduced energy of 0.9 TeV. 2010 marked the beginning of a new era in the high energy frontier with proton collisions at a

record setting 7 TeV. The only other major fault that has occurred was in the summer of 2016. A stone marten chewed through a high voltage line in a power transformer on a ground level building at the LHC. The LHC went offline for about a week while repairs occurred and quickly resumed the physics program. Unfortunately, the marten did not survive.

The typical operating year at the LHC allows for any repairs or upgrades on any of the experiments to be performed during the offline period for the first few months. After the offline period the proton physics program begins and lasts until approximately mid-November. The heavy ion program lasts until the first week of December, after which the LHC shuts down for the remainder of the year. From 2014 until early 2015 the LHC was shutdown for major renovations and upgrades fto the accelerator and a number of sub-detectors on each experiment,. This was known as long shutdown 1 (LS1). After 2018, the LHC will go through another shutdown (LS2) during which the accelerator will have a high luminosity (Hi-Lumi) upgrade. This will be discussed in detail along with the upgrades that will be done to ALICE in chapter [4](#).

3.1.2 LHC Accelerator Complex

The LHC accelerator complex is a succession of particle accelerators that increase the energy of particles before they are injected into the next accelerator. Hydrogen atoms are first passed through a high voltage environment that strips any electrons around the nuclei. Once the nuclei are stripped of their electrons they are injected into the linear accelerator(LINAC). The LINAC uses radio frequency cavities to accelerate particles to 50 MeV before they enter the first circular accelerator the Proton Synchrotron(PS). The PS begins to focus the protons

into bunches and further accelerates them to 1.4 GeV before the beam enters the Super Proton Synchrotron(SPS). The SPS will further accelerate the particles to 450 GeV. The beam is then injected into the LHC. Once the beams are injected into the LHC they are accelerated to the final collision energy. Afterwards the beams get ‘squeezed’, or tightly focused, with a series of quadrupole magnets. The final step is to ‘adjust’ the two beams to overlap with one another at each experiments IP. Once the adjust phase is completed, collisions will occur at each experiment and data begins to get collected. This entire process from stripping the electrons to having collisions in each IP takes 20 minutes.

In order for the beam parameters to be maintained in the LHC, numerous dipole and quadrupole magnets are deployed to accelerate, focus, and bend the particle beams. The magnets use a superconducting niobium-titanium alloy that are maintained at an operating temperature of 1.9 K using helium-4. Upgrading these magnets is one of the major goals during LS2 as part of the Hi-Lumi upgrade of the LHC[76].

3.2 The ALICE Experiment

A Large Ion Collider Experiment(ALICE) is a general purpose detector that covers a solid angle of 4π around the IP. It is 26 m long, 16 m high, 16 m wide, and weighs approximately 10,000 tons[76]. Like many other large scale detectors ALICE is made up of a number of sub-detectors¹ that perform tracking, particle identification(PID), timing, vertex reconstruction, and calorimetry.

¹In total ALICE has 18 sub-detectors




Figure 3.2: The ALICE Detector at CERN[16].




Figure 3.3: ALICE tracker, multiplicity, timing, and vertex detectors located near the interaction point[16].

Figure 3.2 shows the ALICE detector with human figures to set the scale. Figure 3.3 shows the area closest to the IP with the TZERO, VZERO, Inner Tracking System and Forward Multiplicity Detector; these detectors give basic information on the collision such as vertex location, centrality, timing. Further out from the central region are a number of tracking detectors like the Time Projection Chamber and Time-of-Flight detectors that focus on measuring charged particle momentum and PID. Next are the calorimeters that measure particle and jet energies, such as the Electromagnetic Calorimeter, Photon Spectrometer,

and the Dijet Calorimeter. All of these sub-detectors are housed in the L3 magnet seen as the red octagon in Figure 3.2. The L3 magnet provides a uniform magnetic field over the central area of ALICE and is responsible for the high PID performance ALICE has over a wide kinematic region[77]. At high rapidity, there is a muon tracker and trigger for muon identification. The following sections give a more detailed discussion of the sub-detectors used for this analysis

3.2.1 TZERO

The TZERO(T0)[78] detector is a double layer Cherenkov counter located at 70 cm(T0A) and 370 cm(T0B) from the IP. T0 functions as a trigger and timing detector that determines the precise moment in time at which an event ‘starts’ in the ALICE detector. The timing information from the T0 is fed to other sub-detectors, like the Time-of-Flight and Time Projection Chamber detector, for track reconstruction. The T0 also gives feedback on the target luminosity of the ALICE experiment to the LHC operations center.

3.2.2 VZERO

The VZERO(V0)[79] detector is a double layer scintillator array and similar to the T0 is asymmetrically placed at a distance of 86 cm(V0A) and 329 cm(V0C) away from the primary IP. It provides the ‘minimum bias’(MinBias)² trigger information for events and centrality information during the heavy-ion run. Centrality³ is determined by measuring the

²A MinBias event is unsurprisingly defined as an event with the least amount of bias possible. Events recorded with a MinBias trigger attempt to not artificially prefer either diffractive or non-diffractive processes over one another[80].

³See Section ??

multiplicity amplitude from the V0 and fitting these results to a Glauber⁴ distribution, as seen in Figure 3.4. The V0 is also capable of precision measurements of the target luminosity in the ALICE detector.




Figure 3.4: Multiplicity measured in the V0 detector with Glauber fits corresponding to centrality[17].

3.2.3 Inner Tracking System

The Inner Tracking System(ITS)[82] is six layers of solid state silicon detectors. Closest to the beam line are two layers of Silicon Pixel Detectors. The next two layers are Silicon Drift Detectors and furthest from the beam line are two layers of Silicon Strip Detectors. The main purpose of the ITS is to perform momentum measurements, PID, and vertex reconstruction of charged tracks. PID⁵ is performed by measuring the ionization energy, $\frac{dE}{dx}$, of charged particles as they traverse the detector[83]. The ITS has a spatial resolution of $100 \mu\text{m}$. This

⁴A Glauber model treats the nucleons composing a nucleus as hard shells, more can be found here [81].

⁵See Appendix ??

allows for measurements of short lived hadrons by reconstructing secondary vertices, which is the distance of closest approach a track has to the primary vertex.

3.2.4 Time Projection Chamber




Figure 3.5: The ALICE Time Projection Chamber[18].

The Time Projection Chamber(TPC)[18] is a gaseous charged particle tracker and the largest of its kind in the world. The TPC has full azimuthal coverage, a pseudorapidity acceptance of $|\eta| \leq 0.7$, and a volume of $93 m^3$. Figure 3.5 shows a schematic of the TPC. As charged particles traverse the drift volume of the TPC, they ionize the gas inside⁶. A central cathode in the TPC with a voltage of 100 kV induces a uniform electric field of 400 V/m along the beam axis throughout the drift volume. Ionized electrons will drift down to the cylindrical endcaps of the TPC where the read-out chambers(ROC) are located. There are 18 ROCs on each side of the TPC which are further broken into a Inner Read Out Chamber(IROC) and Outer Read Out Chamber(OROC).

⁶The TPC has operated with $Ne - CO_2$ (90-10) and $Ar - CO_2$ (90-10) gas mixtures in the past

The TPC readout




Figure 3.6: The TPC readout region[19].

The TPC incorporates a Multi-Wire Proportional Chamber(MWPC) design for amplification and copper pads for readout⁷. Ionized electrons created from charged particles take approximately $100 \mu\text{s}$ to move from the drift volume to the readout region. Once these electrons enter the readout region they will undergo an amplification process with the MWPC, seen as the sensing grid wires in Figure 3.6. This amplification process will turn the few dozen ionization electrons generated from a charged particle into thousands of amplification electrons that are easily sensed by the cooper pads and read from the front-end electronics(FEE). Amplification using MWPCs has the disadvantage of creating thousands

⁷Their are 72 MWPCs and 500K copper pads in the ALICE TPC

of ions known as ‘backflow ions’ that can move back into the drift volume of the TPC. The presence of backflow ions in the drift volume of the TPC will cause distortions in the uniform electric field of the TPC. These distortions are known as ‘space-charge’ distortions and will compromise the physics performance of the TPC. In order to minimize the space-charge distortions the TPC incorporates a gating grid[?]. Once an event is detected in the readout electronics of the TPC a high voltage is induced on the gating grid. This will capture any backflow ions moving from the amplification region to the drift volume. When engaged the gating grid introduces a dead time as any ionization electrons from new events occurring in ALICE will also get captured. The current configuration of the gating grid is designed to engage it for $300 \mu s$ after an event is first detected. This has been shown to absorb approximately 99% of the backflow ions created while preserving the TPC physics performance. The deadtime due to the gating grid along with the drift time for charged particles in the TPC limits the readout to 3.5 kHz. Upgrading the triggered operation of the current TPC to a continuous readout for the Hi-Lumi upgrade of the LHC will be discussed in detail in Chapter 4.

TPC Performance

In order to reconstruct the trajectory of a particle, an iterative process known as the Kalman filter approach is deployed. The x-y coordinate which is perpendicular to the beamline are determined via the signal induced on the copper pads. The z component which is parallel to the beamline is reconstructed using the timing information from the T0.

The TPC has excellent momentum resolution between $150 \text{ MeV}/c$ to $100 \text{ GeV}/c$ [84]. Figure 3.7a shows the inverse momentum resolution as being below 10% in black. The momentum




Figure 3.7: TPC momentum and tracking resolution[20].

resolution was further improved to almost 5% over the full kinematic range by matching TPC tracks to ITS tracks and constraining those tracks to originate from the primary vertex region, red and green respectively. The matching efficiency between ITS tracks to TPC tracks is stable at about 80%, Figure 3.7b.

3.2.5 Electromagnetic Calorimeter

The Electromagnetic Calorimeter(EMCal)[85] is a lead based sampling calorimeter located at a radius of 4.5 meters from the beam pipe. It covers a pseudorapidity of $|\eta| \leq 0.7$ and has azimuthal coverage of $\Delta\phi = 107$ deg.

Figure 3.8 shows the layout of the EMCal. The smallest element of the EMCal is the ‘tower’⁸. The tower serves as the readout and is made up of several layers of alternating scintillator and Pb-absorber. Particles that interact via the electromagnetic force initiate a shower in the absorber material in the tower. This electromagnetic shower induces light in the scintillator to accumulate in the avalanche photodiodes(APD) in proportion to the particle’s energy. A

⁸There are 12K towers in the EMCal




Figure 3.8: ALICE EMCal along with super modules, tower strips, and towers[21].

‘module’ is an array of four towers that share readout electronics. Twelve modules will be placed in a single strip that provides support to the structure. The largest component of the EMCal is the super-module, consisting of 1,100 towers, which serves as the mounting structure to the ALICE detector. In 2015 Dijet Calorimeter(DCAL) was installed in the ALICE detector to perform back-to-back jet measurements.

EMCal Performance

As particles enter the EMCal they initiate an electromagnetic shower. The shower of electromagnetic particles spans several neighboring towers, these towers are grouped together into ‘clusters’ and the Analog-To-Digital Conversion(ADC) signal from the clusters corresponds to the energy deposited by the particle. The EMCal was designed so that photons and electrons will fully shower inside of the tower region and thus fully deposit their energy. Hadrons on the other hand will punch through the EMCal and only deposit a small

fraction of their energy. PID can be performed using the EMCAL via track-cluster matching from the TPC. TPC tracks are geometrically matched to the centroid of a cluster and if no track is matched the cluster originated from a photon. If a track is matched, then the ratio of E/P, the energy of a matched cluster to the momentum of a TPC track, can be used to separate electrons from hadrons.

The energy resolution of the EMCAL follows the form seen in equation 3.1

$$\frac{\sigma}{E} = \sqrt{A^2 + \frac{B^2}{E} + \frac{C^2}{E^2}} \quad (3.1)$$

where E is the cluster energy, A characterizes stochastic fluctuations such as photon collection efficiency, B is a function of the systematic effects such as detector non-uniformity, and C is a function of the noise in the Front-end electronics.




Figure 3.9: Energy resolution in the EMCAL measured in a 2007 test beam at CERN(blue) compared to GEANT3 simulations of the EMCAL(orange), fits for the parameters A, B, and C are also shown[22].

As seen in figure 3.1 excellent agreement exists between the measured performance of the EMCal compared to simulations in a kinematic range between $10\text{GeV} - 100\text{GeV}$. The stochastic term, A, is the largest source of uncertainty in the energy resolution due to the EMCal being a sampling calorimeter. Unlike the TPC the resolution improves at high- p_t making the EMCal ideal for measuring high energy particles and jets

EMCal Trigger

Due to the high luminosity in the LHC only a small fraction of events may be recorded to disk for later analysis. ALICE employs a variety of triggers to record events that have the highest value for performing quality physics analysis. The EMCal can trigger on events in order to increase the effective sample size for high- p_T jets, photons, and electrons. The two main triggers[86][87] for the EMCal are a jet trigger and a gamma trigger. The gamma trigger is comprised of a 4x4 patch of EMCal towers, while the jet trigger is a 16x16 patch of towers. Once the gamma trigger has surpassed a minimum energy threshold of 5 GeV[88] the event is tagged as a gamma event and the patch location is recorded.

EMCal jet triggered events have an energy threshold of 20 GeV and are similarly tagged and recorded. Figure 3.10 shows the spectra from clusters measured in the EMCal using MinBias data in black and the gamma trigger from the EMCal set to two thresholds, 5 GeV and 7 GeV. Recording the events that satisfy the EMCal triggers introduces a bias towards high- p_T processes, however by using events that had an EMCal trigger we can extend the kinematic range of an inclusive jet measurement as seen in Figure 3.10. In order to account for this bias it is necessary to calculate a trigger efficiency by comparing spectra from inclusive




Figure 3.10: Cluster Spectra from the ALICE EMCal. MinBias is shown in black while the red and blue points show the spectra using the gamma trigger at two energy thresholds[23].

jets recorded using the MinBias trigger to the spectra generated from the EMCal triggers.

The calculation of the trigger efficiency will be discussed in depth in section ??

Chapter 4

The ALICE TPC Upgrade

From 2014 until 2018, I worked on the upgrade of the TPC to a continuous readout. This includes working on a test beam in 2014 for a prototype Read-Out Chamber (ROCs) using Micropattern Gaseous Detectors (MPGD) over the current Multi-Wire Proportional Chamber (MWPC) design. Once a final design for the new ROCs was chosen the production of the new Inner Read-Out Chambers (IROCs) using a stack of four Gaseous Electron Multiplier (GEM) foils took place at the University of Tennessee. During this period I assembled 49 new IROCs. In 2018, I traveled to Yale University and CERN to help with quality assurance of the IROCs built at Tennessee.

4.0.1 Physics Motivation

After the 2018 heavy-ion run ends, the LHC will begin the second long shutdown (LS2) and upgrade to the High Luminosity LHC (Hi-Lumi LHC)[[89](#)] which was briefly mentioned in section [3.1](#). The goal of the LHC upgrade is to move into an era of high precision measurements of rare QCD processes and increase the production of soft probes. Once the

upgrade is complete the expected event rate in ALICE will be 50 kHz in MinBias PbPb collisions. This will lead to a factor of x100 increase in MinBias data and a factor of x10 increase in high- p_T triggered data recorded by ALICE.

The ALICE experiment has focused on probing the thermodynamic properties of the QGP in the past. The increase in event rate expected from the Hi-Lumi LHC would open new ways to probe the QGP including[24]:

- Increasing the production of jets and allowing separation of jet measurements based on the flavor of the initial parton.
- Studying the production mechanisms of light-nuclei, antihyper-nuclei, and other exotic hadrons.
- Probing the initial state temperature and equation of state of the QGP by measuring low-mass dileptons that originate from the earliest stages of a heavy-ion collision.
- Increase the yields of heavy-flavor mesons reconstructed via the secondary-vertices.

In order to do this ALICE will implement a number of upgrades [90] during the shutdown that include:

- Replacing the V0 and T0 detectors with a combined detector, called the Forward Interaction Trigger (FIT)[91].
- Improving the ITS and Muon Tracker spatial resolution by using Monolithic Active Pixel Sensors (MAPS)[92][93].

- Removing the 400 nanosecond dead time associated with the TPC and upgrade the FEE cards to handle the increase in data bandwidth[24].
- New Online/Offline (O^2) data processing architecture[94].

Leptons weakly couple to the QGP[95] and by studying these the initial states in heavy ion collisions may be probed[96]. Figure 4.1 shows a simulation of the mass spectra for dileptons between $0 \text{ GeV}/c - 1 \text{ GeV}/c$ using the current ALICE central detectors with tight kinematic cuts (left). The yields that are quantifiable do not allow for the separation of leptons originating from the QGP from those originating from background sources. After the increase in event rate and the upgraded ITS with improved tracking capabilities, measuring the low-mass dileptons that interacted with the QGP will be possible as shown on the right of Figure 4.1.




Figure 4.1: Simulation of the invariant mass spectra for dileptons in a typical heavy-ion run with current ALICE performance (left) and after upgrade of ALICE for Run-3 (right) in PbPb at $\sqrt{s_{NN}} = 5.5 \text{ TeV}$. The dilepton yields originating from the QGP are shown (red and orange), along with background contributions from light-hadrons (blue), and charm (magenta)[24].

The TPC used in the ALICE experiment along with the readout was discussed in section [3.2.4](#). The ALICE collaboration first proposed an upgrade of the TPC in 2012 with a Letter of Intent (LoI)[\[24\]](#). A Technical Design Report(TDR)[\[28\]](#) was published in 2013 with an initial design overview. An addendum to the TDR[\[29\]](#) was published in 2015 after the performance of prototype ROCs was measured on a test beam at CERN in 2014. The overall goals of the TPC upgrade are to continuously readout events in the high luminosity environment expected after LS2, to minimize the accumulation of space-charge distortions in the drift field, and to maintain the PID and tracking performance of the current TPC.

4.0.2 Gaseous Electron Multiplier Foils

Gaseous Electron Multiplier (GEM) Foils were first proposed by CERN physicist Fabio Sauli in 1997[\[97\]](#). GEMs belong to a new form of detector technology known as Micropattern Gaseous Detectors[\[98\]](#) that use microelectronic or chemical etching techniques to print a readout structure onto a material surface.




Figure 4.2: Scanning electron microscope image of a GEM foil from top (left) and profile (right)[\[25\]](#).

GEM foils are a Kapton sheet, typically $50 \mu\text{m}$ thick, with a thin copper coat on both sides of the surface. A weak acid and stencil are used to chemically etch holes throughout the foil.

Typically, the holes are between $10 \mu m$ - $100 \mu m$ in diameter and between $100 \mu m$ - $300 \mu m$ in pitch as seen in Figure 4.2. A few hundred volts is applied to each of the copper surfaces of the GEM foil causing a strong electric field to be induced in the GEM holes¹.

When an ionization electron from a charged track enters a GEM hole it will cause an avalanche of electrons and ions to be produced, amplifying the signal, to MWPCs. Due to the high voltage and strong electric fields around the GEM foil, any back flow ions created from the avalanche will get absorbed by the copper surfaces on the GEM foil, as seen in Figure 4.3.




Figure 4.3: Profile of GEM with electric-field lines and gradients (left). Simulation of an ionization electron (yellow line) entering a GEM from a drift volume, amplification electrons (green dots, yellow lines) and back flow ions (red lines) are created (right)[26].

The configuration of the pitch and size holes in a GEM foil is directly correlated to the amplification properties and ability to capture back flow ions. GEMS with larger holes or shorter pitch between the holes will have more amplification but will also be more ineffective at capturing ions. Likewise, GEMs with shorter holes or larger pitch will have better ion

¹The electric field is on the order of 10 kV/cm

capture abilities yet worse amplification properties. By stacking multiple GEMs on top of one another it is possible to maximize the amplification properties and minimize the backflow.

Creating GEMs with a uniform hole distribution and stacking them in a consistent manner in order to minimize the overlap of holes from one layer to another restricted them to small prototypes and impeded their use on large experiments. Around 2010 two methods were developed, the single-mask technique[99] and the NS2 assembly[100], which allowed for a systematic way of etching holes and stretching GEMs so they could be properly aligned. These methods allowed for GEMs to become a viable amplification device for large experiments. As of 2018 the TOTEM, COMPASS, ATLAS, and LHCb experiments have all incorporated GEMs in their trackers. Future colliders, such as the Electron-Ion Collider (EIC), plan on using them[101].

4.0.3 Research and Development

Initially, it was decided that the new ROCs would incorporate a stack of three GEMs with a hole geometry similar to the one incorporated on the LHCb experiments[102] tracking detector. This was a starting point to bench mark performance for the ALICE TPC upgrade as well as an opportunity to use experts in GEM technology already present at CERN. The first phase of the R&D involved simulating the performance of the LHCb 3-GEM stack in the high event rate environment expected after the Hi-Lumi LHC upgrade using a the software package called Garfield++[?], which is a GEANT4[103] add-on package built to simulate MPGDs. It was quickly observed from the simulations that a 3-GEM stack would be insufficient in maintaining the performance needed for the TPC while minimizing the ion

back flow. An additional layer of GEM was added to the 3-GEM stack in simulation and it was observed that this configuration would preserve the TPC performance.




Figure 4.4: ITS-TPC matching (*left*) and inverse momentum resolution (*right*) for a 4-GEM stack simulated in Garfield++ [27].

Figure 4.4 shows simulations of the track matching efficiency from the ITS to the TPC and the inverse momentum resolution for a 4-GEM stack as a function of several collisions. The efficiency and resolution only start to diminish at an event rate above 100 kHz, which is double the expected rate after the LHC upgrade, and are well within the range of the current TPC operating at 3.5 kHz.




Figure 4.5: Final design of the upgraded readout chambers with a stack of 4 GEMS [28].

The optimal hole configuration was also explored with simulations. Having a first and forth layer with a pitch of $140 \mu m$ and the second and third layer with a pitch of $280 \mu m$, allowed for the the ROCs to maintain minimal ion backflow while amplifying the signal from tracks. Figure 4.5 shows the design of the ROCs with a 4-GEM stack. The copper pad plane is glued to a reinforced fiberglass sheet, known as the strongback, to add material strength.

2014 Test Beam

During the 2014 test beam, I mounted the prototypes on the beam line, recorded data from the test beam, and monitored the performance in real time. To quantify the performances of the prototype, it is useful to define the effective gain (G_{eff}) and ionic backflow (IBF%)

$$G_{eff} = \frac{I_{anode}}{e N_{ion} R} \quad (4.1)$$

where e is the fundamental electric charge of an electron, N_{ion} is the numbered of captured ions, and R is the illumination rate from an active source.

$$IBF\% = \frac{I_{cathode}}{I_{anode}} = \frac{1 + \epsilon}{G_{eff}} \quad (4.2)$$

where $I_{cathode}/I_{anode}$ is the ratio of the currents measured in the cathode and anode portion of a readout as seen in Figure 3.6. The IBF% can also be defined in terms of the number of backflowing ions (ϵ) created at an effective gain (G_{eff}). The effective gain is a measure of the amplification properties and in a gaseous detector is defined as

The test involved using the beam from both the SPS and PS² at the LHC cast onto an iron target. This iron target created secondary showers and tracks were measured by the prototype for both energy resolution and PID performance. The nominal TPC gas of $CO_2 - N_2$ (90-10) was used during the test beam.




Figure 4.6: dE/dx resolution of the 4-GEM IROC prototype (left) and the separation power between electrons and pions as a function of gain (right)[29].

Figure 4.6 shows the PID performance for separating electron and pion tracks created by the iron target. The separation power between the two Gaussian peaks increases until a gain of 2000 so this was chosen as the target effective gain for the new ROCs.




Figure 4.7: Energy resolution of the iron peak as measured from the prototype IROC with varying GEM voltages as a function of IBF%[29].

²See Section 3.1.2.

Figure 4.7 shows the energy resolution of the iron peak as a function of the relative voltages between GEM 2 (U_{GEM2}) and the ratio of voltages between GEM 3 (U_{GEM3}) and GEM 4 (U_{GEM4}). This shows that the IBF% remains below 1% at a energy resolution of approximately 10% which is consistent with the current TPC performance. The voltages were chosen such that GEM 1 and GEM2, which are closest to the drift volume, would focus mostly on capturing amplification ions while GEM 3 and GEM 4, which are closest to the copper pad readout, would primarily perform the amplification as shown in Figure 4.8.




Figure 4.8: Simulation of the four GEM (blue) layers after test beam. The configuration is such that the two GEMs closest to the drift volume, right side, absorb the amplification ions created by the two GEMs closest to the readout (right) [29].

4.0.4 Production of the Inner Readout Chambers in the U.S.

Once a final design was settled on from the simulation and prototype R&D, the project entered the production phase. A minimum of 36 new ROCs would need to be built with the 4-GEM stack to replace the old ROCs in the TPC while 4 additional ROCs would be constructed as spares. Due to the size and weight of the ROCs, production of the IROCs would take place in the United States and the OROC production mainly in Germany.




Figure 4.9: Production flow of the IROCs (red), OROCs (blue), and GEM foils (green)[27].

Figure 4.13 shows the production flow for the construction of the 4-GEM ROCs. After GEMs are received from the manufacturer, they go through a number of quality assurance tests before they are shipped to Wayne State University, where they are stretched and mounted for the IROCs. The aluminum bodies are manufactured at the University of Texas Austin and shipped to Tennessee. At Tennessee we glue the pad plane, aluminum body and strong back together before it is shipped to Yale, where the chambers are fit with the GEMs from Wayne State. The production steps for the OROC mirror those performed by the IROC except that it was performed mainly German institutes.

Figure 4.10 shows the procedure followed at Tennessee in order to build an IROC. Furthest on the left is a picture of a copper pipe glued into the aluminum body. This copper pipe is flushed with a coolant that maintains the temperature of the Front End Electronics (FEE). The next two photos show the pad plane being mounted on a vacuum table while glue is applied to it and the aluminum body. Once all the components are placed




Figure 4.10: The author assembling an Inner Readout Chamber at the University of Tennessee.

glued and mounted on the table the IROC is loaded with lead bricks and allowed to cure for 24 hours. The right most picture shows the final step of the HV feedthroughs mounted and glued for the GEM foils. Before a completed IROC leaves Tennessee we performed a gas tightness test that will be discussed more in the next section. Full production of the IROCs ended in November of 2018 with a total of 47 chambers being built at Tennessee. A surplus of chambers were built with excess materials to provide spares.

4.0.5 GEM and Chamber Quality Assurance

A stringent set of quality assurance (QA)[30][25] protocols were implemented to monitor any damage sustained during transport and to quickly identify any flaws in the production procedures. The QA can be broken into two major categories: QA performed on the GEM foils as received from the manufacturer and QA performed on complete/semi-complete chambers as they were going through the different production steps. I will discuss only the QA tasks that I was directly involved with.

GEM QA

In order to evaluate the performance of a GEM foil, a spark test was performed on every foil. Sparks are caused by the discharge of the foil and may be due to the presence of dust on a foil, imperfections in the hole geometry, or a short between the two copper layers of the GEM. The final design of the GEM foils had each foil segmented into twelve sectors with a $10 M\Omega$ bias resistor across every segment. Any sparks that occur on the GEM will only discharge a given segment and not the entire foil. Because of the delicate nature of the GEM foil, sparks have the potential to permanently damage a foil or seriously affect the performance of a ROC. Thus GEMs with a high spark rate should be excluded as soon as possible in the production flow.




Figure 4.11: Schematic for the setup of the GEM foil spark test (*left*)^[30] and the GEM mounted in the HV gas box (*right*).

The GEM foil spark test involves mounting each foil in a High Voltage (HV) box, which is flushed with N_2 until it reaches a relative humidity of $\approx 10\%$. Once this humidity is reached 500 V is placed across each segment of the GEM and monitored with a LabView interface. A spark is defined as the LabView recording a current above 10 nA across the bias resistor

through any GEM segment and read by a multi-channel picoammeter. The criteria for a GEM to fail this QA was to have more than five sparks over a 20 minute recording window.

Chamber QA

Most of the QA I helped with was on ROC chambers as they went through the production steps. Before completed IROC chambers were sent to Yale, I performed a gas leak test. The leak test consisted of mounting an individual chamber into an aluminum test vessel and flushing the inside of the vessel with N_2 gas. By using a flowmeter to measure the rate that N_2 gas enters the test vessel and an oxygen sensor to measure the amount of oxygen present at the output of the test vessel we can infer the leak rate of each IROC chamber. Figure 4.12 shows a schematic of the setup (left) and what the typical output from the LabView interface (right). By flowing at two different rates and measuring the oxygen content at each rate we could confirm the leak rate.




Figure 4.12: Schematic of the gas tightness testing setup at the University of Tennessee (Courtesy of Joseph Rasson).

The leak rate of a chamber, f_L , can be quantified as

$$f_L = \left(\frac{\rho_0 - \rho_N}{\rho_L - \rho_0} \right) f_N \quad (4.3)$$

where f_N is the flow rate of N_2 gas into the test vessel, ρ_L is the concentration of O_2 in the laboratory, ρ_N is the O_2 impurity present in the N_2 bottle, and ρ_0 is the O_2 reading from the LabView interface.




Figure 4.13: Leak rate of the 47 chambers built at Tennessee with the maximum failure rate at 0.25 ml/hr shown (*Courtesy of Charles Hughes*).

A maximum leak rate of 0.25 mL/hour was placed as the rejection criteria for a given chamber. If a chamber had a leak rate above this the N_2 gas could get swapped out with a helium gas tank and we could use a helium sniffer to identify the areas on the IROC causing the leak and patch the area with epoxy. Figure 4.13 shows the leak rate for all IROCs produced at Tennessee, due to none of the chambers surpassing the leak threshold the helium sniffer was not used in the production of the IROCs.

Once an IROC chamber was fitted with the 4 GEM foils at Yale and sent to CERN a spark test was performed over the entire chamber. The test involved placing each chamber next to the LHC beam line, flushing with the nominal TPC gas, and placing the nominal voltage across each GEM and monitoring the spark rate once a beam was present in the LHC. Figure 4.14 shows myself installing the completed IROC chambers next to the LHC beam line in front of ALICE and the output from the spark monitor. The voltage across each chamber could be varied in real time in order to minimize sparking while maintaining

the IROC performance, thus reducing the rate of degradation on a per chamber basis once installed in the TPC.




Figure 4.14: The author testing spark testing chambers next to the LHC beam line (*left*) and real time output from the spark test during a live beam (*right*).

4.0.6 Outlook

As of November 2018, the 47 IROCs assembled at Tennessee have been received at CERN. None of the built chambers have surpassed the gas leak test performed at CERN and so far only 4 chambers exhibit sparking rates above the threshold. By May of 2019, the LHC beamline around ALICE will be decommissioned and the TPC will be moved to a surface level clean room for the installation of the chambers. The installation of the new FEE electronics and the ROC chambers should take place throughout the summer of 2019. Afterwards there will be a 10 month commissioning with the TPC, during which the performance of the upgraded TPC will be evaluated with cosmic rays. By the end of 2020, the TPC should be

back in the ALICE cavern and the beam line will have been re-installed. The Hi-Lumi run of the LHC is expected to start in March of 2021.

Chapter 5

Analysis Methods and Corrections

Beginning in March of 2012, the LHC began seven months of pp collisions at $\sqrt{s} = 8$ TeV.

During the 7 months of data taking the ALICE Experiment was recording data almost 60% of the time, Figure 6.11.




Figure 5.1: LHC state during the 8 TeV run.

This large amount of data taking produced almost 200 million events recorded from ALICE that satisfied the Min Bias trigger. The proton-proton Min Bias trigger is defined that at

least one hit was recorded in the SPD or V0. The 8 TeV data set also recorded almost high- p_T events using the EMCAL triggered data, which in the case of the Gamma trigger was a trigger patch threshold of 8 GeV.

This large amount of data offers a unique chance to explore QCD phenomena using jets with a large dataset. In order to measure the jet cross section the following formula is used,

$$\frac{d^2\sigma^{jet}}{d\eta dp_T} = \frac{1}{\epsilon_{reco} \epsilon_{trigger}(p_T)} \times C_{MC} \times \frac{1}{A(p_T)} \times \frac{1}{\mathcal{L}_{int}} \times \frac{dN_{jet}^2}{dp_T d\eta} \quad (5.1)$$

where,

- ϵ_{reco} is the efficiency for reconstructing the jet in the ALICE detector.
- $A_{trigger}$ is the acceptance for EMCAL triggered events and $\epsilon_{trigger}(p_T)$ is the EMCAL trigger efficiency. These factors correct for imperfections in the electronics of the EMCAL and the overall factors are equal to one in minimum bias events.
- C_{MC} is a correction factor due to detector effects and it allows for comparisons between the ALICE experiment to other experiments or theoretical calculations. Unfolding is used to determine this factor.
- \mathcal{L}_{int} is the integrated luminosity during the period when the data was recorded.
- $A(p_T)$ is the geometrical detector acceptance.
- $\frac{dN_{jet}^2}{dp_T d\eta}$ is the inclusive jet momentum spectra.

The following sec

5.1 Raw Jet Spectra

This thesis measured inclusive jet results for radii between 0.1 and 0.5. Furthermore, jet results for radii $R = 0.2$ to $R = 0.4$ will be presented in the body of this chapter while results from the other jet radii are still being investigated. Figures 5.2 5.3 5.4 show the raw (uncorrected) p_T spectra for inclusive jets from both MB and EMCAL triggered data. It is also evident from Figures that the EMCAL triggered data extends the p_T reach of the spectra.




Figure 5.2: Raw inclusive $R = 0.2$ jet spectra from the 8 TeV Min Bias and EMCAL triggered data

The next sections of this chapter will discuss the QA implemented to the data, trigger scaling, unfolding, and other corrections

5.2 8 TeV Data Quality

ALICE is a state-of-the-art experiment with excellent tracking and particle identification capabilities as discussed in Chapter 3. However, just like any real world experiment, it contains a number of inefficiencies and imperfections. This means that the data collected




Figure 5.3: Raw inclusive $R = 0.3$ jet spectra from the 8 TeV Min Bias and EMCAL triggered data

during the 8 TeV pp collision must be examined and any inaccuracies in the data must be removed before hard physics conclusions may be reached. Data may be compromised at both the event-level, the experiment erroneously recorded something as an event, or at the constituent-level, one of the subdetectors mismeasured a feature of a particle, and these outliers must be accounted for and removed

Event Selection

During the 8 TeV data collection period approximately 180 million minimum bias events were recorded, as summarized in table ?? . These events are separated into periods, which dictate the particular beam and detector configurations during the data taking. The 8 TeV data is broken into 7 periods with approximately 181 million minimum bias events recorded. This minimum bias sample corresponds to an integrated luminosity, \mathcal{L}_{int} , of 8.95 pb^{-1} during this time period[104].




Figure 5.4: Raw inclusive $R = 0.4$ jet spectra from the 8 TeV Min Bias and EMCAL triggered data

Approximately, 15% of the data sampled is unusable due to malfunctions in TPC chambers, EMCAL super modules, the electronics for the EMCAL or TPC,. LHC12a,b triggered data is not used in this analysis due to the trigger threshold being varied from the other periods.

Period	# of runs	# of Min Bias events
LHC12c	89	~24 M
LHC12d	140	~62 M
LHC12e	5	~2 M
LHC12f	56	~15 M
LHC12g	8	~0.4 M
LHC12h	159	~75 M
LHC12i	40	~3 M
Total	497	~181 M

Table 5.1: 2012 8 TeV data taking period.

Monte Carlo Anchored Data

The ALICE Collaboration produced two Monte Carlo data sets anchored to the detector performance during the 8 TeV run. LHC15l2a1 is a PYTHIA anchored data set that consisted of about 17 million Min Bias triggered events and LHC15l2a2 is a PHOjet anchored data set consisting of about 21 million Min Bias triggered Events. Neither of these data driven Monte Carlos modeled the EMCal triggers.

General Event Section Criteria

For an event to be selected into a physics analysis it must pass a number of quality control tests. For example, the LHC must have be in a state of stable beams, cosmic rays must be excluded by only accepting tracks that originate from a vertex inside the detector, and the relevant detectors for a given analysis must be functioning as intended. Event selection and QA is implemented via a centralized class, AliEventCuts, within the AliRoot framework. This class contains a number of corrections including:

- The event has a primary vertex reconstructed.
- The primary vertex occurs within a 10 cm window of the primary interaction point.
- The vertex resolution must be below 0.25 cm.
- The event passes basic pile-up checks based on the V0 and T0 signals.

A summary of the rejection reasons at the event level is summarized in Figure 5.5.

Figure 5.6 shows the reconstructed vertex for the accepted Min Bias events. We see that the vertex distribution peaks at the primary interaction point as expected. It should also




Figure 5.5: Min Bias event rejection summary.

be noted that a similar set of event QA was implemented to the EMCal triggered data (*not shown*) and that the results were consistent with the Min Bias data.

5.2.1 EMCal Cluster Selection

Corrections are performed on EMCal cells including; removing hot and dead towers (bad channels) based on the average occupancy and energy of the towers, calibrations to cell timing caused by the physical layout of the EMCal (such as differences in cabling length), and an energy calibration is implemented based on the π^0 mass. After these corrections are applied to the towers are grouped together into clusters using the v2 algorithm. The v2 algorithm has a minimum tower seed, $E_{seed} = 300$ MeV, after which all adjacent towers with a minimum energy, $E_{cell} \geq 100$ MeV, are iteratively added until a local minimum is reached. The cluster energy is the sum of the seed and grouped neighbor tower energies.

Figure 5.7 shows the bad channel map after removing bad towers, the ϕ distribution is segmented into 5 parts representing the five super modules of the EMCal. After the cells




Figure 5.6: Vertex displacement from primary interaction point for accepted Min Bias events.

are clustered together the clusters are corrected for exotics. This correction is performed by cutting all clusters with a $F_{cross} \geq 0.97$, where

$$F_{cross} = 1 - \frac{E_{cross}}{E_{cell}}, \quad (5.2)$$

E_{cross} is the sum of the four cells sharing a full edge with the leading cell. The main source of exotic clusters in the EMCAL is due to a hadron hitting the Avalanche Photodiode (APD) in a tower. This will concentrate the energy of the cluster into a single tower while the adjacent towers will contain only a small fraction of the cluster energy. These clusters are removed before jet finding occurs as they are not part of the jet energy.

The EMCAL is optimized to measure the energy of electrons and photons as they tend to fully shower inside the EMCAL structure. Hadrons are detected by the EMCAL but will only shower a fraction of their intrinsic energy. A hadronic correction is performed in order to account for this missing energy due to the partial hadron shower. Charged tracks from the outer layer of the TPC are propagated to the EMCAL and the centroids of the clusters




Figure 5.7: EMCal cell occupancy after bad channels removed.

and tracks are matched together. Figure 5.8 shows the distance between the centroid of a cluster in the EMCal to the nearest track propagated from the TPC. Hadrons are identified by requiring the matched distance to be, $\sqrt{\Delta\phi^2 + \Delta\eta^2} \leq 0.015$, which is within one EMCal tower distance.




Figure 5.8: Matched track-cluster distance.

Corrections for the double counting from hadrons is based on correcting the EMCAL cluster energy by a weight function,

$$E_{corr} = E_{clust} - f_{sub} \times \sum p, \quad (5.3)$$

where $\sum p$ is the magnitude of the 3-momentum of the hadron and $f_{sub} = 1$ is the nominal value for the weight. If $E_{corr} \leq 0$ the cluster is removed, this may be caused by cluster pile-up and only accounts for a small fraction of the clusters. In order for a cluster to be accepted $E_{corr} \geq 300$ MeV was required. The 300 MeV threshold is required because a minimum ionizing particle (MIP) will on average deposit 280 MeV in the EMCAL.

A final cut was performed on the cluster time.




Figure 5.9: EMCAL cluster time distribution before cuts.

Cutting on the cluster time is done in order to readout only the particles created from an event and to limit the contamination due to slower particles from previous events. The main source of the slow moving particles are neutrons and K_L^0 and this analysis limits cluster time to $t_{clus} \in [-50 \text{ ns}, 100 \text{ ns}]$.




Figure 5.10: Corrected EMCal cluster yield.

Figure fig:EMCalfinal shows the final cluster energy distribution with all the cuts and corrections previously discussed applied and makes up the set of clusters over which the jet finding was performed. The same cuts and QA was applied to the EMCAL triggered data.

5.2.2 TPC Track Selection

Tracks are reconstructed in the TPC using a Kalman filtering which helps alleviate any corrections needed due to multiple scatterings, dead sectors, etc. Jet finding was performed using ‘hybrid’ tracks. Hybrid tracks consist of two track sets the first being all the tracks with at least one hit in the SPD (Global) and the second set being all tracks that can be constrained to the primary vertex (Complimentary). For this analysis, the minimal $p_{T,track}$ is 150 MeV/c and the track must be constrained to: $-0.9 \leq \eta \leq 0.9$ and $0 \leq \phi \leq 2\pi$, shown in Figure 5.11. The spatial distributions of the hybrid tracks remain relatively flat as expected in the 8 TeV data set.




Figure 5.11: Hybrid Track η and ϕ yields.

The quality of the jet p_T resolution was maintained by only accepting jets into the jet finder with a resolution below 1%, Figure 5.12, and this p_T distribution may be seen in Figure 5.13.




Figure 5.12: Accepted hybrid track resolution.

These cuts followed a number of previous track cuts seen in a number of jet results published from ALICE[105].




Figure 5.13: Accepted track p_T yield.

5.3 Jet Selection

Accepted tracks and clusters are feed into the anti- kt jet reconstruction algorithm to reconstruct inclusive jets. A minimum threshold of 5 GeV was used to reconstruct a jet in this analysis. A high- p_T track threshold of 100 GeV is placed on the reconstructed jets. This is motivated by the degradation of the momentum resolution above that energy range. In addition a cut was applied that a jet must be composed of at least one constituent, Figure 5.14 and 5.15.




Figure 5.14: $R = 0.2$ leading track p_T per jet p_T .




Figure 5.15: $R = 0.2$ number of constituents in a jet per jet p_T .

5.3.1 $z_{leading}$ cut

The fragmentation function for the leading track in a jet,

$$z_{leading} = \frac{p_{leading,proj}}{p_{jet}}, \quad (5.4)$$

may be artificially high due to misidentifying secondary decay particles as primary vertex tracks and assigning them a much larger p_T . Additionally, fake clusters, such as exotics, may skew the jet p_T to apparently large values and thus make $z_{leading}$ infinitesimal.




Figure 5.16: $R = 0.2$ $z_{leading}$ from the Min Bias data sample.

Figure 5.16 shows the projection of the hadronic 3-momentum onto the jet axis. We observe an excess of jets, especially at low jet p_T , of z values close to 1 or zero. To help remove these biased jets a cut on $z_{leading} \geq 0.03$ and $z_{leading} \leq 0.97$ is imposed. It should also be noted that due to QCD hadronization $z_{leading} \sim 1$ corresponds to a singular particle jet which is highly rare. In between .03 and .97 we see the $z_{leading}$ is continuous and uniform as expected.

5.3.2 Jet Area Cut

A jet area of, A_{jet} , cut was imposed on accepted jets.

$$A_{jet} \geq 0.6\pi R_{jet}^2 \quad (5.5)$$

The area is estimated in FastJet using ‘ghost’ particles. As jet reconstruction is being performed these fake particles with infinitesimal p_T are placed randomly through the event. The number of ghost particles captured in a jet is proportional to the jet area, thus the precision of the jet area is sensitive to the reconstruction of soft particles. Figure ?? shows the rejection reason for a jet with the dominate reasoning due to the area cut and this distribution skewed towards low- pT jets.

5.3.3 NEF cut

The Neutral Energy Fraction (NEF) is the total jet energy carried by the neutral components of the jet, i.e. EMCal clusters. Figure 5.17 shows the NEF for $R = 0.2$ jets from the Min Bias sample.




Figure 5.17: $R = 0.2$ NEF per jet P_T .

We observe an excess in jets at low- p_T with NEF values around zero or one, similar to what was seen with the $z_{leading}$ distribution. The cause for these excesses were explored in this analysis but no hard source was identified and no cut to the NEF was used.

The jets that passed all of the selection criteria and cuts make up the signal jets that are reported in the cross section and spectra. These criteria were implemented for the $R = 0.3$ and $R = 0.4$ along with the EMCal triggered data.

5.4 EMCal Triggered Data

Figures 5.2 - 5.4 show the raw Min Bias and EMCal triggered data for the 8 TeV run and that the triggered data greatly extends the kinematic reach for the spectra. This thesis looked at the two primary Level-1 triggers configured for the EMCal, the jet trigger (EJE) and the gamma (EGA) trigger[106]. Although both of the Level-1 triggers were investigated in this analysis, only the EGA trigger was ultimately corrected and used for the final jet cross sections and spectra. The EJE trigger utilizes a trigger patch consisting of 32 x 32 EMCal towers sliding by 8 towers until a patch is reconstructed that meets the minimum predefined

trigger energy threshold. Once this threshold is surpassed the event is recorded and tagged as a kEMCEJE triggered event. A similar procedure is followed for the EGA trigger, but with a smaller patch region of 4×4 towers with a sliding window of 2 towers and its own predefined trigger threshold. The bump at low- p_T seen in figures 5.2 - 5.4 is the trigger turn on curve and the peak corresponds to the trigger threshold for the EGA trigger. Ideally, a jet in the EMCAL acceptance should fire the EJE trigger and the EGA should fire due to the presence of a photon or electron. Since, the triggers increase the yield of jets. In order to calculate the trigger bias two macros were used, AliEMCalTriggerMaker.cxx reconstructs the EMCAL and trigger patch geometries while AliEMCalSetupTask.cxx allows a user to set the trigger thresholds for the EkEMCEJE and kEMCEGA triggers. The trigger bias was calculated by reconstructing trigger patches from the Pythia Min Bias sample production created by ALICE and matching those reconstructed patches to jets within the EMCAL acceptance.




Figure 5.18: Distance to closest reconstructed EJE patch to $R = 0.2$ jet with the Min Bias Pythia Monte Carlo.

Figures 5.18 and 5.19 show the distance between a reconstructed jet and its closest reconstructed EMCAL trigger patch for $R = 0.2$ jets in the Pythia Monte Carlo. We can




Figure 5.19: Distance to closest reconstructed EGA patch to $R = 0.2$ jet with the Min Bias Pythia Monte Carlo.

see that with the EGA trigger that the peak of the distribution is within the jet radius of $R = 0.2$ while with the EJE trigger the distribution is more uniformly distributed. The EJE distribution can be accounted for the fact that the EJE trigger is large enough to have multiple jets matched to EJE trigger patch or vice-versa. For the trigger reconstruction I used the EGA trigger and performed a geometrical matching requirement to the spatial center of the trigger patch,

$$\sqrt{(\phi_{jet} - \phi_{EGA,patch})^2 + (\eta_{jet} - \eta_{EGA,patch})^2} \leq R_{jet} \quad (5.6)$$

Equation 5.6 ensures that any EGA trigger patch that fired due to the presence of a jet was confined within the radius parameter of the jet. Double counting may occur if multiple triggers are reconstructed within the same jet. In order to account for this the simulated Monte Carlo spectra is scaled by $N_{Triggers}$ matched to a jet.

5.5 Acceptance Correction

Jet spectra, cross sections, and ratios of cross sections are reported over the full azimuth angle and pseudorapidity acceptance. However, due to jets being constrained to the EMCAL, a geometric factor is used to correct for the limited acceptance of the detector. This thesis uses a maximum jet radius of 0.5 to help study the effects of wide angle radiation on jet fragmentation. Heavy-ion use smaller jet radii, typically of 0.2, to help negate the high multiplicity background. Due to these geometric corrections the centroid of a jet is constrained to,

$$|\eta_{jet}| \leq 0.7 - R, \quad 1.4 + R \leq \phi_{jet} \leq 3.14 - R. \quad (5.7)$$

$$A(p_T) = \frac{(1.4 - 2R) \times (1.745 - 2R)}{2\pi}. \quad (5.8)$$

For jets between $R = 0.1$ through $R = 0.5$ the following jet acceptance corrections are used.

Jet R	$A(p_T)$
0.1	0.296
0.2	0.214
0.3	0.146
0.4	0.091
0.5	0.048

Table 5.2: EMCAL jet acceptance for radii 0.1 - 0.5.

5.6 Unfolding

The reconstructed jet p_T has a number of detector effects ‘folded’ into the measurement.

These effects included such things as:

- Tracking inefficiencies from the TPC and ITS.
- Missing jet energy components from long-lived particles, such as the K_L^0 and neutron, that are cut by the EMCAL timing requirement.
- TPC track p_T and EMCAL cluster energy resolutions.
- Hadronic corrections to the EMCAL cluster spectrum.
- Material loss in the detectors.

Unfolding is the method by which these detector effects are removed from the raw inclusive jet spectra and a ‘true’ jet spectra may be obtained and compared with theoretical calculations or other experimental results. In order to unfold a jet spectra it is necessary to generate a response matrix that simulates the described effects above, after the response matrix is generated a number of statistical approaches including, Bayesian, Singular Value Decomposition (SVD), or Bin-by-Bin, may be applied to unfold the raw jet spectra. In order to generate the response matrix we embed a Pythia generated event into a GEANT3 simulation of the ALICE detector. Due to the fact that the performance and efficiency of the ALICE detector may change between the data taking periods each simulation is ‘anchored’ to a given LHC, these anchors contain all the hot and dead sectors for the subdetectors, along with their calibrated performance during that specified data taking. Two Monte Carlo

data sets were produced with the MB trigger for the full 8 TeV run, the first was a Pythia generator using the Monash-2013 tune and the second was a MB tune of the PHOjet Monte Carlo package. Both data sets were explored for this thesis and it was decided that the final corrected spectra would be obtained via unfolding with the Pythia MC data set. The magnitude of any one of the effects unfolding is supposed to account for is not expected to be very large, but combined may be significant, thus unfolding is an important step in this analysis.

5.6.1 Response Matrix

Given a truth-level particle jet p_T we wish to reconstruct that jet's p_T at the detector-level. The particle-level pythia jets are constructed from the primary particles generated via Pythia while excluding any daughter decay particles in order to avoid double counting. In addition the tracking efficiency in Pythia is known to deviate from nature. This is due to Pythia under predicting the production of strange quarks. Constructing the response matrix in this case is calculated on a jet-by-jet basis. The particle-level jet centroid $(\phi_{part}, \eta_{part})$ is matched to the detector-level jet via a constrain on the displaced distance between the two jet centroids in (ϕ, η) . This distance was constrained to: $\Delta R = \sqrt{(\phi_{part} - \phi_{det})^2 + (\eta_{part} - \eta_{det})^2} \leq 0.25$. Once a jet is matched at the detector level to a jet generated from the particle level the response matrix is incremented by jet p_T at both the detector and Monte Carlo levels. The response matrix is generated with a fine binning with a width of 1 GeV per bin.

Figure 5.20 shows the response matrices for the $R = 0.2$ (top left), $R = 0.3$ (top right), and $R = 0.4$ (bottom) jets generated with the prescribed manner. The response matrices




Figure 5.20: Response Matrices for $R = 0.2$, $R=0.3$, and $R = 0.4$ jets.

display a linear relationship below 50 GeV on both axis and above ~100 GeV the matrices are statistics starved. This is primarily due to the Monte Carlo Pythia and PhoJet data sets generated for the 8 TeV pp run did not model the high- p_T triggers associated with the EMCAL. The particle jet finders configured for the response matrices allowed for jet finding down to a 100 MeV jet candidate at the particle level with no constraints on the minimum particle momentum or energy for a constituent. The detector level jet finders were configured in the same manner as the jet finders configured for the raw jet spectra measurement.

5.6.2 Corrections to Particle Level

Unfolding was performed using the `RooUnfold`[107] software package. Corrections are applied using the bin-by-bin[108] algorithm.

$$C_{MC}(p_T^{low} : p_T^{high}) = \frac{\int_{p_T^{low}}^{p_T^{high}} dp_T \frac{dF_{meas}^{uncorr}}{dp_T} \times \frac{d^2 N_{MC}^{particle}/d\eta dp_T}{d^2 N_{MC}^{detector}/d\eta dp_T}}{\int_{p_T^{low}}^{p_T^{high}} dp_T \frac{dF_{meas}^{uncorr}}{dp_T}} \quad (5.9)$$

where $d^2 N_{MC}^{particle}/dp_T d\eta$ is the PYTHIA level inclusive jet spectra, $d^2 N_{MC}^{detector}/dp_T d\eta$ is the GEANT 3 level inclusive jet spectra, dF_{meas}^{uncorr}/dp_T is a weight function which minimizes the dependence on the two simulation spectra shapes, finally p_T^{low} and p_T^{high} are the lower and upper bin limits. Due to the limited statistics derived from the Monte Carlos available the unfolding procedure was stable only in unfolding the truth level jet spectra for the range: $p_{T,jet} \in [10 \text{ GeV}, 120 \text{ GeV}]$ for both the raw Min Bias and Emcal triggered data sets. Due to the final truth value for the jet spectra will be reported in this range.

5.6.3 Unfolded MB Spectra




Figure 5.21: Unfolded jet spectra with fine binning for $R = 0.3$

Figure 5.21 shows an example of the output from the bin-by-bin unfolding with the fine binning for $R = 0.3$ jets. It should be noted that at low- p_T it was observed that unfolding

increased the yield of the spectra while at high- $p_T \geq 40$ GeV the yield was decreased for all jet radii in this analysis. This is most likely due to the lack of statistics in the response matrix. Once the bin-by-bin unfolding has been performed for the fine binned spectra the output along with the bin-by-bin correction factors are rebinned using a variable binning between 10 GeV and 120 GeV.




Figure 5.22: Unfolded Min Bias $R = 0.2$ jet spectra with corrections factors using a variable binning.

5.6.4 Unfolded EMCAL Triggered Spectra

The unfolding procedure is repeated again for the EMCAL triggered jet spectra. The response matrix from the Min Bias sample is used for the bin-by-bin unfolding and performed using a fine binning. The detector level and particle level jets are configured in the same manner as above and the output from the unfolded triggered spectra are reported after rebinning to a variable size over the same kinematic range as the Min Bias sample.




Figure 5.23: Unfolded Min Bias $R = 0.3$ jet spectra with corrections factors using a variable binning.

Due to the limitations on the response matrix the bin-by-bin unfolding of the EMCal triggered data was only stable up to 120 GeV. It should be noted that the hump in the EMCal jet spectra is due to the firing threshold of the trigger. The unfolded Emcal jet spectra was used to estimate the ratio the the jet yields between the Min Bias and triggered data samples, from this point the trigger scaling was calculated. Due to the lack of a trigger modeled with the 8 TeV Monte Carlo productions the inability to extend the kinematic range of the jet spectras beyond 120 GeV presents a missed opportunity in terms of the recorded data from the 8 TeV runs. In order to address this issue a new Monte Carlo production will need to be requested from the ALICE collaboration.




Figure 5.24: Unfolded Min Bias $R = 0.4$ jet spectra with corrections factors using a variable binning.

5.6.5 Jet Reconstruction and Matching Efficiency

In order to quantify the inefficiencies due to unfolding along with inefficiencies in the ALICE experiment in reconstructing jets we quantify the jet reconstruction efficiency, $\epsilon_{reco}(p_{T,jet})$, and the jet matching efficiency, $\epsilon_{match}(p_{T,jet})$.

$$\epsilon_{reco}(p_{T,jet}) = \frac{N_{reco}(p_{T,jet})}{N_{Truth}(p_{T,jet})} \quad (5.10)$$

$$\epsilon_{match}(p_{T,jet}) = \frac{N_{match}(p_{T,jet})}{N_{Truth}(p_{T,jet})} \quad (5.11)$$

where $N_{reco}(p_{T,jet})$ is the reconstructed jet yield at the detector level per p_T bin, $N_{match}(p_{T,jet})$ is the reconstructed jet at the detector level that was matched to a particle level jet per p_T bin, and $N_{Truth}(p_{T,jet})$ is the truth-level jet yield from the Pythia embedded event per p_T bin. These quantities




Figure 5.25: Unfolded EMCAL triggered $R = 0.2$ jet spectra with corrections factors using a variable binning.




Figure 5.26: Unfolded EMCAL triggered $R = 0.3$ jet spectra with corrections factors using a variable binning.




Figure 5.27: Unfolded EMCAL triggered $R = 0.4$ jet spectra with corrections factors using a variable binning.

Chapter 6

Conclusion and Outlook

The final portion of this chapter will pull all of the results and corrections together and present the final cross sections and the comparison to different theoretical models.

6.1 Systematic Uncertainties

Systematic uncertainties arise due to our limited knowledge of the precise operating conditions and performance of the experiment and also due to any bias in our understanding of how to fundamental model the interactions. They systematics may therefore be broken into two components: uncertainties to the jet energy scale (JES) which shifts the momentum spectra along the x-axis and uncertainties in the jet yield which shift the spectra along the y-axis. The systematical and statistical uncertainties presented in this analysis will be presented as errors to the yield of the spectra. Due to the fact that the p_T distribution follows a power law function, $dN/dp_T \sim p_T^{-5}$ uncertainties in the JES are converted to yield uncertainties by dividing each one by 5. Due to the low statistics at the highest p_T bins in this

analysis, uncertainties in this regime may have large statistical fluctuations. Small systematic variations for the input of the jet spectra will have a dramatic effect over sparsely filled bins versus bins with a low granularity. As such it may be necessary to extrapolate the systematic from a low p_T bin to those at the highest p_T range. The systematic were performed on both the MB and EMCAL triggered data samples but no large variation was observed between the two, thus only the uncertainties from the MB sample are shown and are extrapolated to the triggered data.

6.1.1 Systematic Uncertainty to Jet Energy Scale

The following sections present and discuss the uncertainties cause by shifts to the JES.

Tracking Efficiency Sensativity

Only a fraction of charged tracks generated by the hard scattering of two protons will be detected in the TPC due to finite track efficiency. Uncertainties in the efficiency of the TPC were studied and found to account for a 5% discrepancy[109]. Reproducing this efficiency was performed by randomly throwing out 5% of the tracks from each event from the 8 TeV data samples and remeasuring the jet spectra. All of the inputs for jet finding were maintained.

Figure 6.1 shows the systematical uncertainties for $R = 0.2$ (top left), $R = 0.3$ (top right), and $R = 0.4$ (bottom) jets. A 10% systematic was applied to $R = 0.2$ and $R = 0.3$ jets while a 15% systematic uncertainty was given to $R = 0.4$ jets for this analysis.




Figure 6.1: Systematic due to TPC tracking efficiency; $R = 0.2$ (top left), $R = 0.3$ (top right), $R = 0.4$ (bottom).




Figure 6.2: Systematic due to hadronic correction efficiency; $R = 0.2$ (top left), $R = 0.3$ (top right), $R = 0.4$ (bottom).




Figure 6.3: Systematic due to EMCAL clusterization algorithm; $R = 0.2$ (*top left*), $R = 0.3$ (*top right*), $R = 0.4$ (*bottom*).

Hadronic Correction Systematic

Sensitivity to EMCAL Clusterization Algorithm

As previously stated, the clusterizer used in this thesis was the v2 algorithm. This algorithm was used in both the detector-level Monte Carlo and data analysis. In order to test the sensitivity the JES has to the clusterization algorithm a different algorithm was chosen and a new spectra was generated. The v1 algorithm was choosen and is similar to the v2 algorithm with the exception that the total size of the cluster is forced to be smaller then nine towers. Similar to the other systematic presented we see a large anti-correlated bin-to-bin variations at high- p_T due to sparsely field binning.

6.1.2 Systematic Uncertainty to Jet Yield

The following sections discuss the systematic uncertainties affecting the jet yield.

Track p_T resolution

The momentum resolution of TPC is estimated using the covariance matrix, Figure 6.4, generated from a Kalman filtering[110] pad signal on the TPC read-out region. To estimate the systematic due to the p_T resolution tracks are smeared




Figure 6.4: Inclusive track resolution, Min Bias 8 TeV.

Most tracks have below 1% momentum resolution and the tracks with $\sigma(p_T)/p_T \geq 1\%$ tend to be tracks below 200 MeV tracks. To measure the p_T resolution I smeared each track from the Min Bias data by a Gaussian function which would alter the track p_T by up to 1% of its original value and remeasure the spectra.

Cluster Energy resolution

Min Bias Luminosity and Uncertainty

The luminosity of a hadronic collider, \mathcal{L} , is given by the expression




Figure 6.5: P_T resolution; $R = 0.2$ (top left), $R = 0.3$ (top right), $R = 0.4$ (bottom).




Figure 6.6: energy resolution; $R = 0.2$ (top left), $R = 0.3$ (top right), $R = 0.4$ (bottom).

$$\mathcal{L} = \frac{R}{\sigma} \quad (6.1)$$

where R is the interaction rate and σ is the visible cross section. Due to the fact that we only measure events within a 10 cm window within the primary vertex region we must scale the total luminosity to that which is delivered within the primary vertex region of the ALICE experiment. This scale factor is determined by dividing the total number of MB events to those accepted within the 10 cm window. $N_{MB}^{tot}/N_{MB}^{10cmvertex} = 1.024$ from the acceptance criteria held in this analysis. The luminosity along with its uncertainty were determined during a special Van der Meer scan run in April of 2012[104]. The total systematic uncertainty for the minimum bias (MB) trigger were obtained by measuring the visible cross section using the T0 and V0 detectors. The MB trigger was defined as V0AND which required a hit in both the V0A and V0C. The cross section was reported as being a combined average for MB with the V0AND as,

$$\sigma_{V0} = (55.8 \pm 1.2)mb \quad (6.2)$$

with a combined systematic uncertainty of 2.19% on the visible cross section and 2.60% on the luminosity.

Total Uncertainty

A summary of the total systematic errors used in the final analysis.

Systematic Errors			
Systematic	R = 0.2 Jets	R = 0.3 Jets	R = 0.4 Jets
Clusterization (low- p_T)	1.0%	1.0%	3.0%
(high- p_T)	5.0%	10.0%	10.0%
Hadronic (all bins)	5.0%	4.0%	5.0%
Track Eff (low- p_T)	20.0%	15.0%	15.0%
(high- p_T)	25.0%	20.0%	25.0%
Unfolding (all bins)	6.0%	6.0%	6.0%
p_T Resolution	2.0%	1.0%	4.0%
E Resolution	2.0%	1.0%	5.0%
Luminosity (all bins)	2.2%	2.2 %	2.2%
Total Sys (low- p_T)	8.9%	6.6%	10.9%
(high- p_T)	10.3%	9.1 %	14.5%

The systematics from the yield and JES are added in quadrature together and this is

combined in quadrature with the statistical errors.

6.2 8 TeV Inclusive Jet Results from CMS and ATLAS

CMS[31] and ATLAS[32] both reported the double differential cross section for inclusive jets at 8 TeV.




Figure 6.7: 8 TeV CMS inclusive jet cross sections with radii of $R = 0.7$ and binned by jet rapidity compared to NLO calculations with non-perturbative corrections[31].

6.3 Inclusive Jet Spectra and Cross Section Ratios at 2.76 TeV

Inclusive jet spectra and cross section ratios were measured in the ALICE experiment using a 2011 pp 2.76 TeV data sample[111]. Jets were reconstructed using TPC tracks and EMCAL clusters with the FastJet Anti- K_T algorithm. Tracks with a minimum $p_T \geq 150$ MeV and constrained to within 10 cm of the primary vertex were accepted into the jet finder. EMCAL clusters were




Figure 6.8: $R = 0.4$ inclusive jet cross section at 8 TeV from ATLAS in binned by jet rapidity compared to NLO QCD predictions[32].




Figure 6.9: The 8 TeV ATLAS jet cross sections rescaled to better show comparisons with NLO and non-perturbative calculations at low p_T [32].




Figure 6.10: Inclusive differential cross section from the 2.76 TeV proton proton run with ALICE




Figure 6.11: LHC state during the 8 TeV run.




Figure 6.12: Minimum Bias Event Rejection




Figure 6.13: Emcal Triggered Event Rejection

Bibliography

- [1] C. Patrignani et al. Review of Particle Physics. *Chin. Phys.*, C40(10):100001, 2016. [xiii, 2](#)
- [2] Vardan Khachatryan et al. Measurement of the inclusive 3-jet production differential cross section in proton–proton collisions at 7 TeV and determination of the strong coupling constant in the TeV range. *Eur. Phys. J.*, C75(5):186, 2015. [xiii, 6](#)
- [3] Jets at cms and the determination of their energy scale, July 2012. [xiii, 6](#)
- [4] Matt Dobbs and Jorgen Beck Hansen. The HepMC C++ Monte Carlo event record for High Energy Physics. *Comput. Phys. Commun.*, 134:41–46, 2001. [xiii, 8](#)
- [5] J. Feltesse. Introduction to Parton Distribution Functions. *Scholarpedia*, 5(11):10160, 2010. revision #186761. [xiii, 11](#)
- [6] Martin Erdmann. Investigation of quark - anti-quark interaction properties using leading particle measurements in e+ e- annihilation. *Phys. Lett.*, B510:29–35, 2001. [xiii, 12](#)
- [7] B. R. Webber. Hadronization. In *Proceedings: Summer School on Hadronic Aspects of Collider Physics, Zuoz, Switzerland, Aug 23-31, 1994*, pages 49–77, 1994. [xiii, 12, 13](#)
- [8] Bo Andersson, Sandipan Mohanty, and Fredrik Soderberg. Recent developments in the Lund model. In *36th Annual Winter School on Nuclear and Particle Physics (PINP 2002) and 8th St. Petersburg School on Theoretical Physics St. Petersburg, Russia, February 25-March 3, 2002*, 2002. [xiii, 14](#)

- [9] Jan Rak and Michael J. Tannenbaum. *QCD in hard scattering*, pages 214–243. Cambridge Monographs on Particle Physics, Nuclear Physics and Cosmology. Cambridge University Press, 2013. [xiii, 15](#)
- [10] Gerald C. Blazey et al. Run II jet physics. In *QCD and weak boson physics in Run II. Proceedings, Batavia, USA, March 4-6, June 3-4, November 4-6, 1999*, pages 47–77, 2000. [xiv, 16](#)
- [11] Ryan Atkin. Review of jet reconstruction algorithms. *J. Phys. Conf. Ser.*, 645(1):012008, 2015. [xiv, 16, 20](#)
- [12] Bedangadas Mohanty. Exploring the QCD phase diagram through high energy nuclear collisions: An overview. *PoS*, CPOD2013:001, 2013. [xiv, 26, 28](#)
- [13] J. D. Bjorken. Highly relativistic nucleus-nucleus collisions: The central rapidity region. *Phys. Rev. D*, 27:140–151, Jan 1983. [xiv, 27](#)
- [14] Morad Aaboud et al. Measurement of the nuclear modification factor for inclusive jets in Pb+Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV with the ATLAS detector. *Phys. Lett.*, B790:108–128, 2019. [xiv, 29](#)
- [15] Esma Mobs. The CERN accelerator complex. Complex des accélérateurs du CERN. Jul 2016. General Photo. [xiv, 32](#)
- [16] W. M. Alberico, A. Beraudo, A. De Pace, A. Molinari, M. Monteno, M. Nardi, and F. Prino. Heavy-flavour spectra in high energy nucleus-nucleus collisions. *Eur. Phys. J.*, C71:1666, 2011. [xiv, 35](#)

- [17] Jaroslav Adam et al. Centrality dependence of the charged-particle multiplicity density at midrapidity in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV. *Phys. Rev. Lett.*, 116(22):222302, 2016. [xiv](#), [37](#)
- [18] J. Alme et al. The ALICE TPC, a large 3-dimensional tracking device with fast readout for ultra-high multiplicity events. *Nuclear Instruments and Methods in Physics Research A*, 622:316–367, October 2010. [xiv](#), [38](#)
- [19] Ralf Diener. Gas amplification with micro pattern gas detectors. [xiv](#), [39](#)
- [20] Betty Bezverkhny Abelev et al. Performance of the ALICE Experiment at the CERN LHC. *Int. J. Mod. Phys.*, A29:1430044, 2014. [xiv](#), [41](#)
- [21] G Bourdaud. Gamma-jet physics with the electro-magnetic calorimeter in the alice experiment at lhc. *Journal of Physics: Conference Series*, 110(3):032006, 2008. [xiv](#), [42](#)
- [22] U. Abeysekara et al. ALICE EMCal Physics Performance Report. 2010. [xiv](#), [43](#)
- [23] Cristiane Jahnke. J/ψ production as a function of event multiplicity in pp collisions at $\sqrt{s} = 13$ TeV using EMCal-triggered events with ALICE at the LHC. In *14th International Workshop on Hadron Physics (Hadron Physics 2018) Florianopolis, Santa Catarina, Brazil, March 18-23, 2018*, 2018. [xv](#), [45](#)
- [24] B Abelev et al. Upgrade of the ALICE Experiment: Letter of Intent. Technical Report CERN-LHCC-2012-012. LHCC-I-022. ALICE-UG-002, CERN, Geneva, Aug 2012. [xv](#), [47](#), [48](#), [49](#)

- [25] Erik Brücken and Timo Hildén. GEM Foil Quality Assurance For The ALICE TPC Upgrade. *EPJ Web Conf.*, 174:03004, 2018. [xv](#), [49](#), [57](#)
- [26] Purba Bhattacharya, Bedangadas Mohanty, Supratik Mukhopadhyay, Nayana Majumdar, and Hugo Natal da Luz. 3D simulation of electron and ion transmission of GEM-based detectors. *Nucl. Instrum. Meth.*, A870:64–72, 2017. [xv](#), [50](#)
- [27] Dick Majka. The alice tpc upgrade project. Quark Matter 2017, 2017. [xv](#), [xvi](#), [52](#), [56](#)
- [28] Upgrade of the ALICE Time Projection Chamber. Technical Report CERN-LHCC-2013-020. ALICE-TDR-016, Oct 2013. [xv](#), [49](#), [52](#)
- [29] Addendum to the Technical Design Report for the Upgrade of the ALICE Time Projection Chamber. Technical Report CERN-LHCC-2015-002. ALICE-TDR-016-ADD-1, Feb 2015. [xv](#), [xvi](#), [49](#), [54](#), [55](#)
- [30] Jens Erik Brücken and Timo Eero Hildén. The GEM QA Protocol of the ALICE TPC Upgrade Project. In *5th International Conference on Micro Pattern Gas Detectors (MPGD2017) Temple University, Philadelphia, USA, May 22-26, 2017*, 2018. [xvi](#), [57](#), [58](#)
- [31] CMS Collaboration. Measurement of the double-differential inclusive jet cross section at $\sqrt{s} = 8$ TeV with the CMS detector. 2013. [xviii](#), [98](#), [99](#)
- [32] Morad Aaboud et al. Measurement of the inclusive jet cross-sections in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *JHEP*, 09:020, 2017. [xviii](#), [98](#), [100](#)

- [33] Paul Langacker. Introduction to the Standard Model and Electroweak Physics. In *Proceedings of Theoretical Advanced Study Institute in Elementary Particle Physics on The dawn of the LHC era (TASI 2008): Boulder, USA, June 2-27, 2008*, pages 3–48, 2010. [1](#)
- [34] Tatsumi Aoyama, M. Hayakawa, Toichiro Kinoshita, and Makiko Nio. Tenth-Order Electron Anomalous Magnetic Moment — Contribution of Diagrams without Closed Lepton Loops. *Phys. Rev.*, D91(3):033006, 2015. [Erratum: *Phys. Rev.*D96,no.1,019901(2017)]. [1](#)
- [35] Michael Riordan. The discovery of quarks. *Science*, 256(5061):1287–1293, 1992. [4](#)
- [36] Frank Wilczek. Asymptotic freedom: From paradox to paradigm. *Proc. Nat. Acad. Sci.*, 102:8403–8413, 2005. [Rev. Mod. Phys.77,857(2005)]. [5](#)
- [37] J. D. Bjorken. Asymptotic sum rules at infinite momentum. *Phys. Rev.*, 179:1547–1553, Mar 1969. [7](#)
- [38] J. D. Bjorken. Can We Measure Parton Parton Cross-Sections? *Phys. Rev.*, D8:4098, 1973. [7](#)
- [39] Wolfgang Kurt Hermann Panofsky. Low q electrodynamics, elastic and inelastic electron (and muon) scattering. 1968. [10](#)
- [40] K. Kovarik, T. Jezo, A. Kusina, F. I. Olness, I. Schienbein, T. Stavreva, and J. Y. Yu. CTEQ nuclear parton distribution functions. *PoS*, DIS2013:274, 2013. [10](#)

- [41] Richard D. Ball, Valerio Bertone, Stefano Carrazza, Christopher S. Deans, Luigi Del Debbio, Stefano Forte, Alberto Guffanti, Nathan P. Hartland, José I. Latorre, Juan Rojo, and Maria Ubiali. Parton distributions for the LHC Run II. *JHEP*, 04(arXiv:1410.8849. EDINBURGH 2014-15. IFUM-1034-FT. CERN-PH-TH-2013-253. OUTP-14-11P. CAVENDISH-HEP-14-11):040. 138 p, Oct 2014. Comments: 138 pages, 64 figures. Several typos in text and references corrected, one reference added. [10](#)
- [42] Harald Fritzsch. QCD: 20 years later. In *QCD 20 Years Later: Proceedings, Workshop, Aachen, Germany, June 9-13, 1992*, pages 827–852, 1992. [11](#), [12](#)
- [43] John C. Collins, Davison E. Soper, and George F. Sterman. Factorization of Hard Processes in QCD. *Adv. Ser. Direct. High Energy Phys.*, 5:1–91, 1989. [11](#)
- [44] Nora Brambilla, Xavier Garcia i Tormo, Joan Soto, and Antonio Vairo. The Logarithmic contribution to the QCD static energy at $N^{**}4$ LO. *Phys. Lett.*, B647:185–193, 2007. [12](#)
- [45] B. R. Webber. A QCD Model for Jet Fragmentation Including Soft Gluon Interference. *Nucl. Phys.*, B238:492–528, 1984. [14](#)
- [46] L. L. Jenkovszky, V. K. Magas, J. T. Londergan, and A. P. Szczepaniak. Explicit Model Realizing Parton-Hadron Duality. *Int. J. Mod. Phys.*, A27:1250157, 2012. [14](#)
- [47] J E Huth, N Wainer, K Meier, N J Hadley, F Aversa, Mario Greco, P Chiappetta, J P Guillet, S Ellis, Zoltán Kunszt, and Davison Eugene Soper. Toward a standardization of jet definitions. (FERMILAB-CONF-90-249-E):7 p, Dec 1990. [16](#)

- [48] Matteo Cacciari, Gavin P. Salam, and Gregory Soyez. FastJet User Manual. *Eur. Phys. J.*, C72:1896, 2012. [21](#)
- [49] Megan Connors, Christine Nattrass, Rosi Reed, and Sevil Salur. Jet measurements in heavy ion physics. *Rev. Mod. Phys.*, 90:025005, 2018. [22](#)
- [50] R Brun, F Bruyant, M Maire, A C McPherson, and P Zanarini. *GEANT 3: user's guide Geant 3.10, Geant 3.11; rev. version*. CERN, Geneva, 1987. [22](#)
- [51] Torbjorn Sjostrand, Stephen Mrenna, and Peter Z. Skands. A Brief Introduction to PYTHIA 8.1. *Comput. Phys. Commun.*, 178:852–867, 2008. [23](#)
- [52] Peter Zeiler Skands. Tuning Monte Carlo Generators: The Perugia Tunes. *Phys. Rev.*, D82:074018, 2010. [23](#)
- [53] A. Capella, U. Sukhatme, C.-I. Tan, and J. Tran Thanh Van. Dual parton model. *Physics Reports*, 236(4):225 – 329, 1994. [23](#)
- [54] Cheuk-Yin Wong. *Introduction to high-energy heavy-ion collisions*. World Scientific, Singapore, 1994. Erratum. [23](#)
- [55] Fritz W. Bopp, R. Engel, and J. Ranft. Rapidity gaps and the PHOJET Monte Carlo. In *High energy physics. Proceedings, LAFEX International School, Session C, Workshop on Diffractive Physics, LISHEP'98, Rio de Janeiro, Brazil, February 16-20, 1998*, pages 729–741, 1998. [23](#)
- [56] M. Bahr et al. Herwig++ Physics and Manual. *Eur. Phys. J.*, C58:639–707, 2008. [24](#)

- [57] T. Csorgo. New form of matter at CERN SPS: Quark matter but not quark gluon plasma. *Nucl. Phys. Proc. Suppl.*, 92:62–74, 2001. [,62(2000)]. [25](#)
- [58] I. Arsene et al. Quark gluon plasma and color glass condensate at RHIC? The Perspective from the BRAHMS experiment. *Nucl. Phys.*, A757:1–27, 2005. [25](#)
- [59] K. Adcox et al. Formation of dense partonic matter in relativistic nucleus–nucleus collisions at rhic: Experimental evaluation by the {PHENIX} collaboration. *Nuclear Physics A*, 757(1–2):184 – 283, 2005. First Three Years of Operation of {RHIC}. [25](#)
- [60] B.B. Back et al. The {PHOBOS} perspective on discoveries at {RHIC}. *Nuclear Physics A*, 757(1–2):28 – 101, 2005. First Three Years of Operation of {RHIC}. [25](#)
- [61] J. Adams et al. Experimental and theoretical challenges in the search for the quark–gluon plasma: The {STAR} collaboration’s critical assessment of the evidence from {RHIC} collisions. *Nuclear Physics A*, 757(1–2):102 – 183, 2005. First Three Years of Operation of {RHIC}. [25](#)
- [62] Barbara V. Jacak and Berndt Müller. The exploration of hot nuclear matter. *Science*, 337(6092):310–314, 2012. [26](#)
- [63] Shanshan Cao, Tan Luo, Guang-You Qin, and Xin-Nian Wang. Heavy and light flavor jet quenching at rhic and lhc energies. *Physics Letters B*, 777:255 – 259, 2018. [28](#)
- [64] Michael L. Miller, Klaus Reygers, Stephen J. Sanders, and Peter Steinberg. Glauber modeling in high energy nuclear collisions. *Ann. Rev. Nucl. Part. Sci.*, 57:205–243, 2007. [28](#)

- [65] Klochkov and I Selyuzhenkov and. Centrality determination in heavy-ion collisions with the CBM experiment. *Journal of Physics: Conference Series*, 798:012059, jan 2017. [29](#)
- [66] James L. Nagle and William A. Zajc. Small System Collectivity in Relativistic Hadronic and Nuclear Collisions. *Ann. Rev. Nucl. Part. Sci.*, 68:211–235, 2018. [30](#)
- [67] Wenbin Zhao, You Zhou, Hao jie Xu, Weitian Deng, and Huichao Song. Hydrodynamic collectivity in proton–proton collisions at 13 tev. *Physics Letters B*, 780:495 – 500, 2018. [30](#)
- [68] STEPHEN MYERS. The large hadron collider 2008–2013. *International Journal of Modern Physics A*, 28(25):1330035, 2013. [31](#)
- [69] Thomas Taylor and Daniel Treille. The Large Electron Positron Collider (LEP): Probing the Standard Model. *Adv. Ser. Direct. High Energy Phys.*, 27:217–261, 2017. [31](#)
- [70] G. Aad et al. The ATLAS Experiment at the CERN Large Hadron Collider. *JINST*, 3:S08003, 2008. [31](#)
- [71] S. Chatrchyan et al. The CMS Experiment at the CERN LHC. *JINST*, 3:S08004, 2008. [31](#)
- [72] A. Augusto Alves, Jr. et al. The LHCb Detector at the LHC. *JINST*, 3:S08005, 2008. [31](#)

[73] K. Aamodt et al. The ALICE experiment at the CERN LHC. *JINST*, 3:S08002, 2008.

31

[74] Serguei Chatrchyan et al. Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC. *Phys. Lett.*, B716:30–61, 2012. 31

[75] Georges Aad et al. Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC. *Phys. Lett.*, B716:1–29, 2012. 31

[76] C. Fabjan and J. Schukraft. The Story of ALICE: Building the dedicated heavy ion detector at LHC. In '*The Large Hadron Collider: A marvel technology*', EPFL-Press Lausanne, Switzerland, 2009 (Editor: L. Evans), chapter 5.4, 2011. 34

[77] Vladimir V. Gligorov, Simon Knapen, Benjamin Nachman, Michele Papucci, and Dean J. Robinson. Leveraging the ALICE/L3 cavern for long-lived exotics. 2018.

36

[78] M. Bondila et al. ALICE T0 detector. *IEEE Trans. Nucl. Sci.*, 52:1705–1711, 2005.

36

[79] E. Abbas et al. Performance of the ALICE VZERO system. *JINST*, 8:P10016, 2013.

36

[80] Rick Field. Min-Bias and the Underlying Event at the LHC. *Acta Phys. Polon.*, B42:2631–2656, 2011. 36

[81] Constantin Loizides. Glauber modeling of high-energy nuclear collisions at the subnucleon level. *Phys. Rev.*, C94(2):024914, 2016. 37

- [82] S. Beolè. The alice inner tracking system: Performance with proton and lead beams. *Physics Procedia*, 37:1062 – 1069, 2012. Proceedings of the 2nd International Conference on Technology and Instrumentation in Particle Physics (TIPP 2011). [37](#)
- [83] Jaroslav Adam et al. Particle identification in ALICE: a Bayesian approach. *Eur. Phys. J. Plus*, 131(5):168, 2016. [37](#)
- [84] Christian Lippmann. Performance of the alice time projection chamber. *Physics Procedia*, 37:434 – 441, 2012. Proceedings of the 2nd International Conference on Technology and Instrumentation in Particle Physics (TIPP 2011). [40](#)
- [85] A Fantoni and the ALICE collaboration. The alice electromagnetic calorimeter: Emcal. *Journal of Physics: Conference Series*, 293(1):012043, 2011. [41](#)
- [86] Olivier Bourrion, R Guernane, B Boyer, JL Bouly, and G Marcotte. Level-1 jet trigger hardware for the alice electromagnetic calorimeter at lhc. *Journal of Instrumentation*, 5(12):C12048, 2010. [44](#)
- [87] O Bourrion, N Arbor, G Conesa-Balbastre, C Furget, R Guernane, and G Marcotte. The alice emcal l1 trigger first year of operation experience. *Journal of Instrumentation*, 8(01):C01013, 2013. [44](#)
- [88] Jiri Kral, Terry Awes, Hans Muller, Jan Rak, and Joachim Schambach. {L0} trigger for the {EMCal} detector of the {ALICE} experiment. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 693:261 – 267, 2012. [44](#)

- [89] G Apollinari, I Béjar Alonso, O Brüning, M Lamont, and L Rossi. High-Luminosity Large Hadron Collider (HL-LHC) : Preliminary Design Report. 2015. [46](#)
- [90] Levente Molnar and the ALICE Collaboration. Upgrade and physics perspective of alice at the lhc. *Journal of Physics: Conference Series*, 589(1):012014, 2015. [47](#)
- [91] T L Karavicheva and ALICE Collaboration. The fast interaction trigger detector for the alice upgrade. *Journal of Physics: Conference Series*, 798(1):012186, 2017. [47](#)
- [92] B Abelev and all. Technical Design Report for the Upgrade of the ALICE Inner Tracking System. Technical Report CERN-LHCC-2013-024. ALICE-TDR-017, Nov 2013. [47](#)
- [93] Technical Design Report for the Muon Forward Tracker. Technical Report CERN-LHCC-2015-001. ALICE-TDR-018, Jan 2015. [47](#)
- [94] P Buncic, M Krzewicki, and P Vande Vyvre. Technical Design Report for the Upgrade of the Online-Offline Computing System. Technical Report CERN-LHCC-2015-006. ALICE-TDR-019, Apr 2015. [48](#)
- [95] Radoslaw Ryblewski and Michael Strickland. Dilepton production from the quark-gluon plasma using leading-order (3+1)D anisotropic hydrodynamics. *Acta Phys. Polon. Supp.*, 8(2):445, 2015. [48](#)
- [96] Mauricio Martinez and Michael Strickland. Measuring QGP thermalization time with dileptons. *Phys. Rev. Lett.*, 100:102301, 2008. [48](#)

- [97] F. Sauli. GEM: A new concept for electron amplification in gas detectors. *Nucl. Instrum. Meth.*, A386:531–534, 1997. [49](#)
- [98] Maxim Titov. Perspectives of Micro-Pattern Gaseous Detector Technologies for Future Physics Projects. In *Proceedings, CMS Workshop: Perspectives on Physics and on CMS at Very High Luminosity, HL-LHC: Alushta, Crimea, Ukraine, May 28–31, 2012*, pages 241–258, 2013. [49](#)
- [99] M Tang, A Q Liu, A Agarwal, and M H Habib. A single-mask substrate transfer technique for the fabrication of high-aspect-ratio micromachined structures. *Journal of Micromechanics and Microengineering*, 17(8):1575, 2007. [51](#)
- [100] W. You, Y. Zhou, J. Liu, M. Shao, C. Li, H. Chen, Z. Tang, Y. Sun, Y. Zhang, Y. Lv, G. Song, X. Wang, D. Hu, and D. Hong. A new technique for assembling large-size gem detectors. *Journal of Instrumentation*, 12(06):C06036, 2017. [51](#)
- [101] Fabio Sauli. The gas electron multiplier (gem): Operating principles and applications. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 805:2 – 24, 2016. Special Issue in memory of Glenn F. Knoll. [51](#)
- [102] Marco Santimaria and Davide Pinci. Study of the GEM detector for the LHCb experiment upgrade, Nov 2013. Presented 19 Dec 2013. [51](#)
- [103] S. Agostinelli et al. GEANT4: A Simulation toolkit. *Nucl. Instrum. Meth.*, A506:250–303, 2003. [51](#)
- [104] ALICE luminosity determination for pp collisions at $\sqrt{s} = 8$ TeV. Mar 2017. [66](#), [97](#)

- [105] Shreyasi Acharya et al. Charged jet cross section and fragmentation in proton-proton collisions at $\sqrt{s} = 7$ TeV. *Phys. Rev.*, D99(1):012016, 2019. [74](#)
- [106] O. Bourrion, R. Guernane, B. Boyer, J. L. Bouly, and G. Marcotte. Level-1 jet trigger hardware for the ALICE electromagnetic calorimeter at LHC. *JINST*, 5:C12048, 2010. [78](#)
- [107] Tim Adye. Unfolding algorithms and tests using RooUnfold. In *Proceedings, PHYSTAT 2011 Workshop on Statistical Issues Related to Discovery Claims in Search Experiments and Unfolding, CERN, Geneva, Switzerland 17-20 January 2011*, pages 313–318, Geneva, 2011. CERN, CERN. [84](#)
- [108] G. Cowan. A survey of unfolding methods for particle physics. *Conf. Proc.*, C0203181:248–257, 2002. [,248(2002)]. [84](#)
- [109] Betty Bezverkhny Abelev et al. Energy Dependence of the Transverse Momentum Distributions of Charged Particles in pp Collisions Measured by ALICE. *Eur. Phys. J.*, C73(12):2662, 2013. [92](#)
- [110] R. Fruhwirth. Application of Kalman filtering to track and vertex fitting. *Nucl. Instrum. Meth.*, A262:444–450, 1987. [95](#)
- [111] Rongrong Ma. Measurements of the inclusive jet cross section and jet fragmentation in pp collisions with the alice experiment at the lhc. *Nuclear Physics A*, 910-911:319 – 322, 2013. Hard Probes 2012. [99](#)

Appendices

Vita

Andrew John Castro was born in San Jose, CA. He grew up in the San Francisco Bay Area and Mexico City. In 2012 he was awarded Bachelors of Science from the California Polytechnic University in Pomona, CA. During his undergraduate education Andrew worked with Dr. Alex Small on simulating tumor growths and Dr. Antonio Auerilia on high energy physics. In 2011, Andrew spent 3 months at the Euorpean Organization for Nuclear Research (CERN) working with Dr. Tancredi Carli of the ATLAS experiment on parton distribution functions and jet physics. After his undergraduate studies Andrew entered the University of Tennessee pursuing a Ph.D in Physics. Andrew defended his thesis in 2019. During his time at Tennessee Andrew worked on developing trigger code for the ALICE EMCal, heavy flavor jets, light-nuclei production, hadronization, and Monte Carlo simulations. He also contributed to the upgrade of the ALICE Time Projection Chamber to a continuous readout mode using micro pattern gaseous detectors. Andrew currently works for the Space Radiation Analysis Group with NASA in Houston, Texas. His work is focused on the impact that heavy-ion radiation has to biological material and solutions to shield future manned missions in space. He lives with his fiance and three cats.