Why the Fine-Structure Constant Governs Gravitational Coupling: Two Routes to One Scale and the Derived Constitutive Law

Stephen Nelson
Independent Researcher

February 2026

Complementary Technical Note to "Dual Tetrad Gravity And The Field Origin"

Purpose. The expression $G = \frac{17}{13}\,\alpha^{21}\,\frac{\hbar c}{m_e^2}$ derives the gravitational constant from topology to 0.12%. Two questions naturally follow:

- **Why does the fine-structure constant appear in a gravitational formula?** The fine-structure constant $\alpha \approx 1/137$ measures the strength of the electromagnetic interaction. Its presence in G, raised to the 21st power, requires a structural explanation.
- **Where does the constitutive law $\mu(x) = x/(1+x)$ come from?** The companion technical note derives a unique Lagrangian from this function, but the function itself must be derived, not postulated.

This note answers both questions from the same $(3+1)^2$ topology:

- **Sections I–II:** Two independent routes, one electromagnetic and one gravitational, converge on the same characteristic scale $r_e = 2.82$ fm. Alpha is the ratio of this scale to the Compton wavelength, a topological property that both forces share. Alpha governs gravitational coupling because it measures the per-vertex coupling strength at the interaction center, the same center through which gravity must traverse 21 levels.
- **Section III:** The prefactor 17/13 is derived from the standard $3+1$ decomposition of the bimetric interaction matrix, using only tetrads, the ADM formalism, and the Hassan-Rosen interaction potential. Rewriting G in physical units absorbs one factor of α into r_e, yielding the core equation $G\,m_e/(r_e\,c^2) = (17/13)\,\alpha^{20}$.
- **Section IV:** Gravitational fields consist of distributed, locally-coupled degrees of freedom (confirmed by LIGO in 2017). These field regions tile the available geometry. The $k = 4$ complementary channels of the stellated octahedron tile the sphere completely, giving coupling constant $C = 1$ and yielding $\mu(x) = x/(1+x)$ as a uniqueness theorem within the minimal closure class. This is the same saturation function that governs molecular adsorption (Langmuir isotherm) and enzyme kinetics (Michaelis-Menten).

I. Two Routes to One Scale

If the fine-structure constant measures the strength of electromagnetic interaction, why does it appear 21 times in the formula for the gravitational constant? The answer is that α is not exclusively electromagnetic. It is a ratio of two length scales that the topology determines, and both electromagnetism and gravity converge on the same scales through independent physics.

The electromagnetic route. Classical electrodynamics defines a characteristic length: the distance at which a charged particle's field energy equals its rest mass. For the electron, this is the

classical electron radius:

$$r_e = \frac{\alpha\,\hbar}{m_e\,c} = 2.818 \times 10^{-15}\ \text{m} \tag{1}$$

This is textbook physics, following from Coulomb's law and the mass-energy relation. No topology enters; the scale is purely electromagnetic.

The gravitational route. The companion technical note ("Covariant-Complete $(3+1)^2$ Gravity," Section II) establishes that three equations from three independent domains, gravitational closure, cosmological horizon physics, and the quantum-electromagnetic scale, constrain two unknowns (r and m, the size and mass of a single field region). This overdetermined system converges on:

$$r = 2.82 \times 10^{-15}\ \text{m}\,, \qquad m = 9.11 \times 10^{-31}\ \text{kg}$$

matching the electron radius and electron mass to 0.1%, using gravitational and cosmological inputs with zero adjustable parameters. No electromagnetic data enters this calculation; the electron is an *output*.

The convergence. Two independent routes, one purely electromagnetic and one purely gravitational/cosmological, find the same characteristic scale. This convergence is unique to the electron among known particles [7]. It reflects structural constraint, not coincidence. The topology produces one characteristic scale, r_e, and both forces probe it.

An analogy from condensed matter clarifies the situation. The speed of sound in a crystal is determined by the lattice spacing and bond strength. Thermal vibrations and acoustic waves both depend on the same lattice. They are not "borrowing" from each other; they share the same structure. Similarly, electromagnetism and gravity share the same topological scale r_e. The fine-structure constant,

$$\alpha = \frac{r_e}{\bar{\lambda}_C} \tag{2}$$

where $\bar{\lambda}_C = \hbar/(m_e c)$ is the reduced Compton wavelength, is a property of the topology itself: the ratio of the characteristic field scale to the quantum scale. Alpha is not borrowed from electromagnetism by gravity; it is the topology's characteristic ratio, which electromagnetism probes at one level and gravity traverses across 21.

II. Alpha as the Topology's Per-Vertex Coupling

Having established that r_e is a topological scale shared by both forces, we now ask: what physical quantity does α measure within the topology?

The electron possesses two physically measurable rates. The first is the *field closure rate*: the speed at which the gravitational field maintains consistency across its $k = 4$ coupling channels. Field closure requires all four tetrad legs to settle into an agreed-upon state [1]; the speed of this reconciliation is c (Lorentz invariance admits no other fundamental speed), and the distance to be traversed is r_e:

$$f_{\text{closure}} = \frac{c}{r_e} \tag{3}$$

This is the tick rate of the field's internal consistency process.

The second is the *Compton frequency*, the tick rate of the electron's quantum phase:

$$f_{\text{Compton}} = \frac{m_e\,c^2}{h} \tag{4}$$

Two rates, both physical, both measurable. One describes how fast the field maintains consistency; the other describes how fast the particle's quantum state evolves. Their ratio reveals the fine-structure constant:

$$\frac{f_{\text{closure}}}{f_{\text{Compton}}} = \frac{c/r_e}{m_e c^2/h} = \frac{h}{m_e\, c\, r_e} = \frac{\lambda_C}{r_e} = \frac{2\pi}{\alpha} \tag{5}$$

Therefore:

$$\frac{1}{\alpha} = \frac{1}{2\pi} \cdot \frac{f_{\text{closure}}}{f_{\text{Compton}}} = 137.036 \tag{6}$$

The number $1/\alpha = 137$ is this scale ratio: the reduced Compton wavelength exceeds the field closure scale r_e by exactly this factor. Alpha is the per-vertex coupling strength at the interaction center, set by the topology's own characteristic scales [7].

Why per-level attenuation follows. Electromagnetism operates at a single vertex: one charge couples to one field, engaging only level 0 of the coupling hierarchy. Gravity, by contrast, must achieve field closure across the entire dual tetrad, coupling through the interaction vertex (level 0), the four interfaces of one sector (level 1), and all 16 bimetric channels (level 2), traversing 1+4+16 = 21 coupling levels. Each level attenuates by α because α is the single-vertex coupling strength. The force ratio is:

$$\frac{F_{\text{gravity}}}{F_{\text{EM}}} \sim \frac{\alpha^{21}}{\alpha^{1}} = \alpha^{20} \approx 10^{-43}$$

This is the hierarchy problem, resolved by counting coupling depth. The full counting derivation appears in the companion note "The Origin of Every Integer in G," Section III.

What is and is not derived. The framework explains α's *role* (per-level attenuation) from topology. The framework does not derive α's *value* ($\approx 1/137$) from first principles; α is taken as empirical input. This is analogous to how crystallography explains why the speed of sound depends on lattice spacing without deriving the numerical value of the lattice constant from geometry alone. The structural role is determined; the magnitude is measured.

III. The Force Ratio Is Twenty, Not Twenty-One

Section II established that gravity traverses 21 coupling levels, each attenuating by α, while electromagnetism engages only level 0. The force ratio is $\alpha^{21}/\alpha^{1} = \alpha^{20} \approx 10^{-43}$. But the formula for G contains α^{21}, not α^{20}. Where does the extra factor of α go? And what determines the prefactor 17/13?

Both questions have the same answer: they follow from the standard $3 + 1$ decomposition applied to the bimetric interaction.

What a tetrad is. General relativity describes gravity through the metric tensor $g_{\mu\nu}$, a 4×4 symmetric matrix that encodes how distances are measured at each point in spacetime. The tetrad provides a deeper description: it is a set of four basis vectors, one for each spacetime direction, that together define a local reference frame [12]. Each tetrad assigns four basis vectors at a point: one timelike, three spacelike. If you know these four vectors, you can reconstruct the metric. The metric tells you distances; the tetrad tells you directions.

Each tetrad leg is a vector in four-dimensional spacetime with four components, one for each coordinate direction (t, x, y, z). Four legs, each with four components: $4 \times 4 = 16$ numbers specify a single tetrad completely.

The interaction matrix. Hassan and Rosen [10] demonstrated that ghost-free massive gravity requires two dynamical metrics, and therefore two tetrads. The interaction between them is described by the matrix product of one tetrad with the inverse of the other: a 4×4 matrix with 16 independent components. These 16 components are the complete set of ways that the two tetrads can couple. In the Hassan-Rosen formulation, they correspond to the 16 terms distributed across the five elementary symmetric polynomials of the interaction potential:

$$\sum_{n=0}^{4} \binom{4}{n} = 1 + 4 + 6 + 4 + 1 = 16$$

The interaction occurs at a specific point: the coupling vertex, where the two tetrads meet and the symmetric vielbein condition must be satisfied [11]. This vertex contributes one additional condition: the requirement that the determinant of the tetrad does not vanish ($\det(e) \neq 0$). Without this condition, the local frame is undefined and the interaction has no meaning. The total interaction space is therefore $16 + 1 = 17$.

The spatial projection. Gravity curves space. The time direction matters for the dynamics, but the curvature manifests in the three spatial dimensions. The ADM formalism [1] makes this separation explicit: it splits the four-dimensional spacetime metric into three dimensions of space evolving through one dimension of time. This $3+1$ decomposition is how every numerical relativity simulation, every gravitational wave template, and every ADM constraint equation in the literature is constructed.

When the $3 + 1$ split is applied to the tetrad, the question becomes: of the four legs, how do they project onto the three spatial dimensions? Each of the four tetrad legs has a spatial part with three components. Four legs times three spatial components: $4 \times 3 = 12$. These 12 numbers are the spatial content of the tetrad, describing how the four frame vectors sit in three-dimensional space. The remaining 4 components (one per leg, the time components) describe the dynamical evolution but not the spatial structure.

The same non-degeneracy condition applies to the spatial projection: the determinant must not vanish, and the spatial frame must span all three dimensions. This adds 1. The total spatial structure is $12 + 1 = 13$.

The prefactor. The ratio of the full spacetime interaction to the spatial structure is:

$$\frac{17}{13} = \frac{k^2 + 1}{k \cdot d + 1} = \frac{\text{spacetime interaction} + \text{vertex}}{\text{spatial projection} + \text{vertex}} \tag{7}$$

where $k = d + 1 = 4$ is the number of tetrad legs and $d = 3$ is the spatial dimension.

An analogy makes the logic concrete. The electromagnetic field tensor $F_{\mu\nu}$ has $\binom{4}{2} = 6$ independent components in four-dimensional spacetime. Of these, $\binom{3}{2} = 3$ are purely spatial (the magnetic field) and 3 are mixed spacetime (the electric field). The spacetime-to-spatial ratio is $6/3 = 2$. The same $3+1$ decomposition applied to the bimetric interaction matrix gives $17/13$ instead of $6/3$. The only structural difference is that bimetric gravity requires the non-degeneracy condition ($+1$ on both numerator and denominator) that electromagnetism does not.

The logic chain, from start to finish: one tetrad has 4 legs with 4 components each, giving 16 numbers. Two tetrads interact through a 4×4 matrix, also 16 components. The non-degeneracy condition at the interaction vertex adds 1; total interaction: 17. The ADM $3+1$ split projects each of the 4 legs onto 3 spatial dimensions, giving 12 spatial components; the same non-degeneracy adds 1; total spatial structure: 13. The prefactor is 17/13.

Every ingredient is standard: tetrads [12], the ADM decomposition [1], the Hassan-Rosen interaction potential [10], and the symmetric vielbein condition [11]. The integers 17 and 13 follow from counting the components of objects that general relativity already requires. (The complete term-by-term counting appears in the companion note "The Origin of Every Integer in G," Section IV.)

Rewriting G in physical units. With the prefactor established, we return to the exponent. The formula for G is:

$$G = \frac{17}{13}\,\alpha^{21}\,\frac{\hbar c}{m_e^2} \tag{8}$$

The scale $\hbar c/m_e^2$ contains no length explicitly. Substituting $\hbar = r_e\, m_e\, c\,/\,\alpha$, which follows from the definition of the classical electron radius (Eq. 1), yields:

$$G = \frac{17}{13}\,\alpha^{21} \times \frac{r_e\, m_e\, c}{\alpha} \times \frac{c}{m_e^2} \;=\; \frac{17}{13}\,\alpha^{20}\,\frac{r_e\, c^2}{m_e} \tag{9}$$

One factor of α has been absorbed. The abstract scale $\hbar c/m_e^2$ has become the concrete scale $r_e\, c^2/m_e$, which contains the electron radius explicitly.

Which α was absorbed. In the coupling hierarchy $1 + 4 + 16 = 21$, the leading "1" is the Field Origin: the center point where the two tetrahedra meet. When one α converts $\hbar$ into r_e, it is the Field Origin's contribution that is consumed. The remaining twenty, corresponding to $4 + 16 = 20$ channels beyond the origin, remain as attenuation.

The twenty-first level does not weaken gravity. It establishes *where* gravity operates: the characteristic scale $r_e = 2.818 \times 10^{-15}$ m at which the topology exists physically. This is why two independent routes, one electromagnetic and one gravitational (Section I), converge on the same length. Both measure the Field Origin from different sides.

The core equation. Dividing both sides of Eq. (9) by $r_e\, c^2/m_e$ yields:

$$\frac{G\, m_e}{r_e\, c^2} \;=\; \frac{17}{13}\,\alpha^{20} \tag{10}$$

The left side contains four measured quantities (G, m_e, r_e, c); it is the gravitational potential of the electron at its own characteristic radius, expressed in units of c^2. The right side contains the topological prefactor and twenty powers of the fine-structure constant. The left side is constructed entirely from experiment; the right side is constructed entirely from the topology.

This equation states that G and α are not independent constants. Given the electron mass and radius, knowing α determines G, and knowing G determines α. They are linked by the integer $20 = k + k^2 = 4 + 16$, which counts the coupling channels of the stellated octahedron beyond its center. The topology of three-dimensional space locks them together. The framework contains no free parameters to adjust.

IV. The Constitutive Law from Distributed Field Coupling

The companion technical note ("Covariant-Complete $(3{+}1)^2$ Gravity," Section III) derives a unique Lagrangian from the constitutive law $\mu(x) = x/(1+x)$. This section establishes where that constitutive law comes from. The derivation proceeds in two stages: first, a physical argument for why packing geometry governs gravitational coupling, then the geometric and kinetic derivations that both converge on $\mu(x) = x/(1+x)$.

Why packing geometry governs gravity. On August 17, 2017, LIGO (recipient of the 2017 Nobel Prize in Physics) and Virgo detected gravitational waves from a neutron star merger [2], a disturbance that propagated from point to point across 130 million light-years. This observation confirmed that gravity consists of distributed, locally-coupled degrees of freedom. The wave equation encodes this requirement: gravitational perturbations satisfy $\partial^2 h/\partial t^2 = c^2\,\nabla^2 h$, and the Laplacian ∇^2 is defined by comparing field values at adjacent locations. You cannot compute ∇^2 without knowing the field at neighboring points. Gravity must therefore consist of coupled field regions, not a structureless continuum.

Water waves provide a concrete parallel. They propagate because water molecules push their neighbors; the disturbance passes from molecule to molecule across thousands of kilometers. Water molecules form tetrahedral hydrogen-bonding structures, and this molecular architecture determines the bulk properties (viscosity, surface tension, specific heat) that govern large-scale behavior. Now consider calculating ocean tides while assuming water is a structureless continuum. Such a calculation would fail: ignoring molecular structure while trying to predict bulk dynamics would require inventing fictitious forces to explain the discrepancy [7].

At galactic scales, standard cosmology treats spacetime as structureless and attributes the resulting discrepancy between predicted and observed dynamics to invisible mass comprising 85% of the total. The $(3+1)^2$ framework identifies the missing structure: gravitational fields couple through the dual tetrad topology with a characteristic tile size r_e (established in Section I) and a characteristic acceleration scale:

$$a_0 = \frac{k^2\,G\,m_e}{r_e^2} = \frac{16\,G\,m_e}{r_e^2} \approx 1.2 \times 10^{-10}\ \mathrm{m/s^2} \tag{11}$$


where $k^2 = 16$ counts the coupling channels through the tetrahedral geometry. This derived value matches the empirically observed MOND acceleration scale [4] to 0.41%. Above a_0, all 16 coupling channels engage and gravity is Newtonian. Below a_0, channels saturate and the gravitational response changes character.

At galactic scales, the gravitational field consists of coupled field regions, each of characteristic size r_e, tiling the available geometry, just as water molecules tile a volume of liquid. The coupling between adjacent tiles follows from their overlap geometry. The question "which $\mu(x)$?" therefore reduces to a geometric one: what is the coupling constant of the tiling?

The geometric origin of the "1." The general form of the constitutive law is $\mu(x) = x/(x+C)$, where C is the coupling constant of the field's tiling geometry. The functional form follows from saturation physics (derived below); the value of C follows directly from the $(3+1)^2$ topology.

Return to the stellated octahedron inscribed in the spherical field region. In three-dimensional close packing, each sphere has 12 geometric neighbors, the kissing number for dense sphere packing [9]. These 12 contacts map to the 12 edges of the stellated octahedron, one edge per neighbor. But coupling does not flow through edges; it flows through *faces*.

What happens at each face. Each tetrahedron has $k = 4$ triangular faces. In the stellated octahedron, each vertex of TetB passes through the center of a TetA face, and vice versa. This vertex-through-face penetration induces a $3 + 1$ subdivision on each face (Figure 1).



(a) Face of TetA **(b)** Coupled face: $3+1$

Figure 1: The coupling topology of a single face. **(a)** A triangular face of TetA. **(b)** When TetB couples through the Field Origin, its vertex V_B penetrates the face center, subdividing the face into $3+1=4$ sub-regions: three corner sub-regions (blue) anchored at the original vertices, and one central coupling sub-region (red) at the penetration point. This is the same $3+1$ structure that appears within each tetrahedron (3 spatial +1 temporal face) and across both tetrahedra ($(3+1)^2=16$ interpenetrations).

The result is $3+1=4$ sub-regions per face, the inevitable consequence of a point entering a triangle. The same $3+1$ pattern recurs at every scale of the stellated octahedron:

- **Within each face:** 3 corner sub-regions +1 central penetration = 4 channels
- **Within each tetrahedron:** 3 spatial faces +1 temporal face = 4 faces
- **Between both tetrahedra:** $(3+1)\times(3+1)=16$ interpenetrations

The coupling hierarchy $1+4+16=21$ counts this self-similar pattern at three successive resolutions. The complete counting is developed in the companion note "The Origin of Every Integer in G."

From faces to coupling constant. For the constitutive law, the relevant structure is the face level. Each face-vertex interpenetration creates a single coupling channel: a TetA face coupled to its piercing TetB vertex through the Field Origin, where $t=0$. The two tetrahedra do not contribute 8 independent faces. They couple through the origin into $k=4$ complementary channels, each consisting of a face from one sector coupled to its piercing vertex from the other. From our $(3+1)$ spacetime we observe half of this structure (the geometric origin of spin $\hbar/2$ [7], Section 2.6), but at the Field Origin the two halves are one.

Each complementary channel subtends π steradians, one quarter of the sphere's total solid angle. The question is whether these four channels account for all coupling directions. They do:

$$C_{\text{tiling}} = k \times \frac{1}{k} = 4 \times \frac{1}{4} = 1 \tag{12}$$

No sharing correction is needed: each complementary channel is already a single coupling unit (face + vertex through the origin), not a bilateral interface counted from two sides. The "1" in $\mu(x)=x/(1+x)$ is the angular completeness of the tetrahedral tiling: four simplex faces partition the sphere with no gaps and no overlaps [8].

Cross-check: the two-dimensional limit. Galactic disk dynamics are effectively two-dimensional. In 2D, optimal packing is hexagonal [3], with kissing number $N=6$. The 6 neighbors decompose into two triangular sets of 3 (the 2D dual simplex), whose complementary coupling gives $C=3\times(1/3)=1$, the same result, because the dual simplex tiles the available angular space completely in any dimension. Equivalently, using the standard overlap calculation: $C=6\times(\frac{1}{3})\times(\frac{1}{2})=1$, confirming $C=1$ through a second route.

The kinetic derivation: detailed balance. The topology determines $C = 1$. A kinetic argument independently confirms the functional form. At each field point, coupling capacity partitions into two states: a *resolved* fraction ε that tracks Newtonian demand, and an *unresolved* fraction $(1 - \varepsilon)$ where coupling has saturated.

Two competing rates govern the partition:

- **Resolution rate:** $W_+ = \kappa \cdot x \cdot (1 - \varepsilon)$. The gravitational intensity $x = g/a_0$ drives channels toward resolution, limited by available capacity $(1 - \varepsilon)$.
- **Relaxation rate:** $W_- = \kappa \cdot \varepsilon$. Resolved capacity relaxes back at a rate proportional to the resolved fraction.

At equilibrium ($W_+ = W_-$):

$$\kappa \cdot x \cdot (1 - \varepsilon) = \kappa \cdot \varepsilon \quad \Longrightarrow \quad \varepsilon = \frac{x}{1 + x} \tag{13}$$

Since the resolved fraction is the fraction of Newtonian gravity that the field transmits, $\mu(x) \equiv \varepsilon(x) = x/(1 + x)$.

This result is unique within the Minimal Single-Channel Saturating Closure (MSC) class [5]:

1. The driving exponent is forced to $n = 1$ by the deep-MOND limit ($\mu/x \to 1$ as $x \to 0$), observed in galaxy rotation curves [4]. Any other exponent violates this limit.
2. Single-channel saturation is the minimal closure consistent with the observed single acceleration scale a_0. The SPARC sample (> 150 galaxies) shows no evidence for additional transition scales.
3. The boundary conditions $\mu(0) = 0$ and $\mu \to 1$ as $x \to \infty$ are physical requirements: zero coupling at zero gradient, full Newtonian response at strong field.

The uniqueness proof appears in "Distributed Field Processing: The Gravitational Constitutive Law," Theorem 1 [5].

A universal saturation pattern. The function $x/(1 + x)$ appears throughout physics and chemistry as the equilibrium of any system with finite capacity, linear driving, and reversible transitions:

Domain	**System**	**Equilibrium form**
Surface physics	Langmuir isotherm	$\theta = Kp/(1+Kp)$: fraction of adsorption sites occupied
Biochemistry	Michaelis-Menten	$v/V_{\max} = [S]/(K_m + [S])$: fraction of enzyme active sites engaged
Gravity	Field closure	$\mu = x/(1 + x)$: fraction of coupling channels resolved

In each case, a finite number of sites (surface sites, enzyme active sites, coupling channels) are driven toward occupancy by an external quantity (gas pressure, substrate concentration, gravitational intensity) and relax at a rate proportional to occupancy. The equilibrium is the same because the physics is the same: saturation of a finite resource under linear driving.

From constitutive law to Lagrangian. The Euler-Lagrange equation for the scalar field relates the Lagrangian to the constitutive law through $d\mathcal{F}/dx = x\,\mu(x)$ (the standard AQUAL variational

relationship [4]). Integrating $\mu(x) = x/(1+x)$ with the boundary condition $\mathcal{F}(0) = 0$ yields:

$$\mathcal{F}(y) = \tfrac{1}{2}\, y - \sqrt{y} + \ln(1+\sqrt{y})$$

where $y = x^2 = |\nabla\Phi|^2/a_0^2$. This Lagrangian is unique: the integration constant is fixed by $\mathcal{F}(0) = 0$, and the constitutive law admits no alternative form under the field closure boundary conditions. The full action and its field equations appear in the companion note "Covariant-Complete $(3+1)^2$ Gravity," Section III.

V. The Derivation Chain

The three companion technical notes form a single argument:

1. "The Origin of Every Integer in G" counts each integer in $G = \frac{17}{13}\,\alpha^{21}\,\frac{\hbar c}{m_e^2}$ from the coupling structure of ghost-free bimetric gravity. Every integer (4, 16, 21, 17, 13) answers a specific counting question about the stellated octahedron.

2. "Covariant-Complete $(3+1)^2$ Gravity" presents the uniquely determined action, its field equations, and five zero-parameter predictions across gravity, cosmology, particle physics, galactic dynamics, and the CMB.

3. This note (the present document) explains why α governs both forces, because electromagnetism and gravity share the same topological scale r_e (Sections I–II); derives the prefactor $17/13$ from the standard $3+1$ decomposition and establishes the core equation $G\,m_e/(r_e\,c^2) = (17/13)\,\alpha^{20}$ (Section III); and derives the constitutive law $\mu(x) = x/(1+x)$ from packing geometry and channel saturation (Section IV).

Together, the chain is: the $(3+1)^2$ topology determines the integers in G (counting note), provides the per-vertex coupling constant α (this note, Sections I–II), derives the prefactor and the core equation from the $3+1$ decomposition of the bimetric interaction (this note, Section III), determines the constitutive law from the distributed coupling structure (this note, Section IV), integrates to a unique Lagrangian (covariant note, Section III), and yields five predictions with zero adjustable parameters (covariant note, Section IV).

VI. Acknowledged Limitations

- The numerical value of α ($\approx 1/137$) is empirical input. The framework derives α's structural role (per-vertex attenuation) but not its magnitude.

- The detailed balance derivation assumes single-channel saturation with linear driving (the MSC class). Multi-channel correlations or nonlinear saturation functions would modify the transition shape near $g \sim a_0$, producing testable residuals in rotation curve fits [5].

- The packing derivation identifies $C = 1$ from the complementary face-vertex structure of the stellated octahedron. If coupling at galaxy cluster scales engages additional neighbor contacts beyond the $k = 4$ tetrahedral channels, C would increase and the transition shape near $g \sim a_0$ would change, a testable prediction [5].

- Full limitations of the $(3+1)^2$ framework, including galaxy cluster dynamics, the CMB power spectrum, and the nuclear forces, appear in the companion note "Covariant-Complete $(3+1)^2$ Gravity," Section VI.

- The prefactor 17/13 depends on the non-degeneracy condition at the coupling vertex. If higher-order corrections modify this condition, the prefactor would shift, producing a testable change in the predicted value of G.

References

[1] R. Arnowitt, S. Deser, and C. W. Misner, "The Dynamics of General Relativity," Gen. Relativ. Gravit. **40**, 1997 (2008).
[2] B. P. Abbott *et al.* (LIGO/Virgo), "A gravitational-wave standard siren measurement of the Hubble constant," Nature **551**, 85 (2017).
[3] T. C. Hales, "The Honeycomb Conjecture," Discrete Comput. Geom. **25**, 1 (2001).
[4] M. Milgrom, "A modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis," Astrophys. J. **270**, 365 (1983).
[5] S. Nelson, "Distributed Field Processing: The Gravitational Constitutive Law," Field Dynamics Working Paper (2026). Available as supplemental material.
[6] S. Nelson, "A Covariant Theory Without Dark Matter," Field Dynamics Working Paper (2026). Available as supplemental material.
[7] S. Nelson, "Dual Tetrad Gravity And The Field Origin" (2026). Submitted for publication.
[8] H. S. M. Coxeter, *Regular Polytopes*, 3rd ed. (Dover, 1973).
[9] T. C. Hales, "A proof of the Kepler conjecture," Ann. Math. **162**, 1065 (2005).
[10] S. F. Hassan and R. A. Rosen, "Bimetric gravity from ghost-free massive gravity," JHEP **02**, 126 (2012).
[11] C. Deffayet, J. Mourad, and G. Zahariade, "A note on 'symmetric' vielbeins," JHEP **03**, 086 (2013).
[12] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (W. H. Freeman, 1973).