Dual Tetrad Topology and the Field Origin: From Nuclear Decay to Galactic Dynamics

Bidirectional Validation of a Zero-Parameter Gravitational Field Theory

Stephen Nelson[1,*]

[1] *Independent Researcher*

ORCID: 0009-0008-6190-3269

(Dated: February 2026)

Abstract

Gravity occupies a unique position among the fundamental forces: it is dynamical spacetime itself, requiring self-consistency at every point. General relativity encodes this through field closure, the requirement that the metric remain non-degenerate ($\det(g) \neq 0$) everywhere. In tetrad formulation, closure forces four basis vectors at each point to span the tangent space completely, defining a tetrahedron at each field node. Ghost-free bimetric gravity (Hassan-Rosen 2012) couples two such tetrad fields through shared boundary faces. Field coupling closure in (3+1) dimensions determines a unique minimal topology: the stellated octahedron, two tetrahedra with opposite orientations coupled at a shared origin.

Deriving a covariant scalar-tensor action from that coupling hierarchy yields quantitative predictions. The topological constant $k = 4$ (from simplex packing in (3+1) dimensions) predicts the neutron lifetime beam-bottle discrepancy (1.084% vs observed 1.081%, agreement to 0.3%) and Milky Way rotation velocity (229 km/s, exact match). Bidirectional validation across 36 orders of magnitude, from nuclear physics (10^{-15} m) to galactic dynamics (10^{21} m) to cosmological structure (10^{26} m), with zero adjustable parameters.

When dual tetrahedral field regions couple, the interaction occurs at three structural levels: field existence (1), propagation through vierbein faces ($k = 4$), and Hassan-Rosen bimetric channels ($k^2 = 16$). The coupling polynomial $f(k) = 1 + k + k^2 = 21$ is not phenomenological but the architecture that field coupling closure requires. Structural identification of the 16 bimetric interaction channels with the coupled face channels of the stellated octahedron yields a zero-parameter covariant completion, distinct from Hassan-Rosen bimetric gravity which contains five free functions. The resulting scalar-tensor action passes 14 empirical tests spanning GW speed (exact to 10^{-15}), Mercury precession, lunar laser ranging, wide binary plateaus, rotation curves, and cosmological expansion, with zero fitted parameters compared to ΛCDM's 15-20.

Read for coupling depth, the hierarchy yields $G = (17/13)\alpha^{21}\hbar c/m_e^2$ (0.12% tree-level, 0.001% one-loop). Read for interaction breadth, it yields $1/\alpha = 137.036$ (0.00021% agreement). The pattern across independent tests demonstrates that the missing 85% attributed to dark matter reflects missing structure in how we model field dynamics, not missing particles. Fundamental constants are topological consequences of how gravitational fields couple in (3+1) dimensions.

PACS numbers: 04.60.-m, 04.50.Kd, 04.20.Cv, 06.20.Jr, 03.65.Ta

I. INTRODUCTION

The fundamental constants of physics have been measured with extraordinary precision. Newton's gravitational constant G has been the subject of increasingly refined experiments since Cavendish's 1798 torsion balance measurement, yet for over three centuries no theory has calculated its value from first principles. The Standard Model of particle physics requires over two dozen such parameters [**?**], each determined by experiment but not derived from any deeper principle. That these constants take the values they do is what makes atoms stable, chemistry possible, and structure viable at every scale. Explaining why they take precisely these values has remained one of the deepest open questions in physics.

The stakes extend beyond unexplained constants. Galactic rotation curves, gravitational lensing, and the cosmic microwave background all indicate that approximately 85% of the universe's gravitating mass has never been directly detected [**?**]. Four decades of dedicated searches have produced no confirmed dark matter particle. The assumption underlying standard cosmology is that spacetime is a structureless continuum whose gravitational response is fully characterized by a single constant, G. But if the gravitational field possesses internal coupling structure that modifies its response at different scales, the question changes: is the missing 85% missing mass, or is it missing structure?

In other branches of physics, the relationship between internal topology and observable properties is well established. The molecular lattice of a ruby crystal determines the discrete energy levels available to electrons passing through it; the result is stimulated emission, the physical basis of the laser. Semiconductor band gaps, superconducting transition temperatures, and phonon spectra all follow from the coupling architecture of their respective lattices. In each case, the internal topology of the medium determines the measurable constants that describe it. Just as the angles of a crystal are determined by its lattice, the constants of nature may be determined by the topology of the gravitational field. They would not be free parameters that could have been different; they would be structural consequences of living in three spatial dimensions.

Gravity, too, possesses an internal coupling structure. The tetrad formalism [**?** **?**] defines the "square root of the metric," $g_{\mu\nu} = \eta_{ab}\, e^a_\mu\, e^b_\nu$, decomposing each metric into a local frame

* Contact author: stephen@fielddynamics.org

whose spatial triad defines a tetrahedron [**?**], the minimal solid in three dimensions. Ghost-free bimetric gravity [**?** **?**] operates with two such tetrad fields, $g = E^T \eta\, E$ and $f = L^T \eta\, L$, coupled at a single interaction vertex through a potential with five free coefficients β_0 through β_4. Ghost-freedom requires opposite spatial orientations for the two tetrads [**?**]; general covariance requires them to share compatible (3+1) decompositions [**?**]. Together, these constraints define a dual tetrahedron field topology whose field coupling hierarchy has not been systematically read out. This paper reads that hierarchy and shows what it produces.

Gravity occupies a unique position among the fundamental forces: it is not a field on a background spacetime but the dynamical spacetime itself, and it must therefore be self-consistent at every point with no external structure to enforce coherence. General relativity encodes this through the non-degeneracy of the metric ($\det(g) \neq 0$), which requires the field to close: to define a complete coupling structure at every point, with no gaps and no missing directions. When field configurations couple, they must settle into a compatible state with consistent geometry across all coupling channels. In this paper, we refer to this process as *field closure* (FC_Σ): the continuous enforcement of self-consistency that the gravitational field must satisfy. Closure forces the field to develop internal architecture, creating the coupling chambers through which neighboring field regions interact. Each spacetime point carries a local frame of four basis vectors that couple to neighboring frames through shared boundary faces. The pattern of these couplings constitutes the field's topology. If this topology is constrained tightly enough by self-consistency, the fundamental constants may not be free parameters at all but consequences of how the gravitational field couples to itself.

Atomic structure emerges from electromagnetic field coupling: the interaction between charged particles produces persistent configurations with definite internal topology, and that topology determines measurable properties from spectral lines to chemical bonds. Gravitational field coupling produces structure by the same principle. When field regions interact through the (3+1) tetrad frames that general relativity already requires, the coupling creates a self-consistent field architecture whose topology is not a free choice but a consequence of the field's own consistency requirements.

The physical picture begins with field packing. Gravitational field regions, being distributed and locally coupled, pack in three dimensions. The simplex theorem establishes that $k = 4$ faces are both necessary and sufficient to enclose a volume in three dimensions, and these

four faces define a tetrahedral field chamber, the minimal solid enclosing the coupling region. A single tetrahedral field chamber, however, is not self-consistent. Four independent results from the literature, spanning five decades and over 2,300 combined citations, all demonstrate this: spinor theory demands the SL(2,$\mathbb{C}$) double cover [**?** **?**]; Ashtekar's chiral decomposition requires both self-dual and anti-self-dual connections [**?**]; ghost-free bimetric gravity requires two dynamical metrics [**?**]; and stability analysis confirms that a single orientation produces a scalar ghost [**?**]. Self-consistency requires a second tetrahedron with opposite orientation, coupled to the first at a shared center: the Field Origin. The resulting structure, two interlocking tetrahedra sharing a common center, is the stellated octahedron [**?**].

The field coupling structure of this stellated octahedron is a polynomial in $k = (3{+}1) = 4$:

$$f(k) = 1 + k + k^2 \tag{1}$$

whose three terms correspond to structures already recognized in the literature: the Field Origin (1) [**?**], the vierbein face ($k = 4$) [**?**], and the full bimetric field coupling ($k^2 = 16$) [**?**]. The stellated octahedron has three structural regions (TetA, the Field Origin, TetB), and evaluating $f(k)$ in each yields three integers: $f(4) = 21$ (the full coupling depth), $f(4)|_{s=0} = 17$ (the coupling capacity at the Field Origin), and $f(4)|_{s=-1} = 13$ (the spatial structure of the inverted sector). These three integers, together with the quantum scale $\hbar c/m_e^2$, determine G:

$$G = \frac{17}{13} \times \alpha^{21} \times \frac{\hbar c}{m_e^2} \tag{2}$$

Topology: $G_{\text{topo}} = 6.666 \times 10^{-11}\ \text{m}^3\,\text{kg}^{-1}\,\text{s}^{-2}$

Measured: $G_{\text{CODATA}} = 6.674 \times 10^{-11}\ \text{m}^3\,\text{kg}^{-1}\,\text{s}^{-2}$ [**?**]

Agreement: 0.12% at tree level; 0.001% at one loop. Not fitted.

The polynomial is not fitted to G. It is the field coupling structure that the stellated octahedron's topology requires. The entire framework rests on a single discrete fact: space has three dimensions ($d = 3$). From this one input, every other number follows through geometry, linear algebra, and dimensional analysis. Zero free parameters are adjusted. The test is simple: start from $d = 3$, follow the chain, and if any single prediction disagrees with measurement, the framework fails without adjustment. There are no dials to turn.

If this field topology derives G, does it also explain other fundamental relationships? The answer, developed in Sections **??** through **??**, is that the same dual tetrahedron field topology produces quantitative results across classical, relativistic, and quantum physics. The inverse square law follows from field spreading in three dimensions. Mass-energy equivalence follows from the bimetric field structure and its two closure speeds. The force hierarchy follows from the depth of field coupling within the 21-level structure. Electron spin $\hbar/2$ follows from the face ratio of the stellated octahedron. Dark matter observations are accounted for by the constitutive law that field closure imposes. These are not separate explanations for separate phenomena. They are aspects of one field topology, the topology that yields G.

A. Scope and Limits

This paper presents what the dual tetrahedron topology explains about fundamental physics. It does not claim to explain everything. The fine-structure constant α is derived in Section **??** from the Hamiltonian interaction states of the bimetric topology, yielding $1/\alpha = 137.036$ to 0.00021%. The electron mass m_e is derived in Section **??** from the exactly determined closure system. No particle-physics input is required; the Hubble constant H enters as a cosmological observable and is itself predicted by the topology (Section **??**). The cosmological constant, the strong force, and the weak force are not directly addressed. The goal is clarity about what the topology does explain, not a claim to have solved all of physics.

B. Overview

Section **??** establishes the theoretical foundations: LIGO confirms gravitational propagation, the wave equation requires neighbors, and field closure is what GR already enforces through non-degeneracy. Section **??** reviews the dual tetrahedron topological structure from which the derivations follow. Section **??** develops the coupling polynomial, the heart of the paper. Section **??** derives the fundamental equations: G from tetrad coupling, the inverse square law, mass-energy equivalence, the force hierarchy, and quantum consistency. Section **??** derives the fine-structure constant. Section **??** addresses dark matter observations at galactic scales and predicts the Hubble constant. Section **??** presents the covariant completion.

Section **??** derives the electron mass. Section **??** presents the precision context, distinctive predictions, and the complete dependency chain.

II. WHY GRAVITY HAS TOPOLOGY

This section establishes that topology is not an addition to general relativity but what GR already requires. The gravitational field propagates (LIGO confirmed this in 2015). Propagation requires neighbors (the wave equation demands it). And neighbors that must be self-consistent at every point, with no external structure to enforce coherence, must achieve field closure (GR has enforced this for a century through non-degeneracy). Once there is structure, counting begins. Numbers emerge from counting, not from fitting. Everything that follows in this paper is an exercise in counting the right geometric object.

A. The Gravitational Field Propagates

General Relativity describes gravity through the metric tensor $g_{\mu\nu}$, a dynamical field satisfying Einstein's equations. This is standard physics, confirmed by a century of observations. GR works beautifully. But when we ask how the gravitational field couples to itself, how one region of spacetime influences its neighbors, the answer lies in a reformulation that many physicists learn but few fully explore: tetrad gravity.

Tetrad gravity, also called vierbein or frame-field gravity, is textbook material [**?** **?**]. It describes the same physics as metric GR but makes the field's internal structure explicit. The tetrad is defined as the "square root of the metric": the decomposition $g_{\mu\nu} = \eta_{ab}\, e^a_\mu\, e^b_\nu$ expresses the metric as the inner product of two copies of the tetrad field. In the (3+1) decomposition, the three spacelike legs of the tetrad (the spatial triad) define a tetrahedron [**?**], the minimal solid in three dimensions. This tetrahedral structure at each point couples to neighboring tetrahedra through shared faces. The mathematics has been in graduate textbooks for decades, yet the physical implications of this coupling topology remain underappreciated.

Why does gravity, uniquely among the fundamental interactions, require us to ask about its internal structure? Because gravity is not a field that lives *on* spacetime. Gravity *is*

spacetime. The electromagnetic field inherits its topology from the background metric: it knows what "neighbors" means because spacetime tells it. But the gravitational field is the metric itself. It cannot borrow structure from a background, because it is the background. If the field must be self-consistent and no external structure exists to enforce that consistency, the field must provide its own closure mechanism. This is the physical content of background independence, one of GR's deepest features.

LIGO's first detection of gravitational waves in September 2015, which earned the 2017 Nobel Prize in Physics, confirmed that gravity propagates as a wave through locally-coupled structure. The subsequent detection on August 17, 2017, when LIGO and Virgo observed gravitational waves from a neutron star merger arriving 1.7 seconds before the gamma-ray burst [**?**], confirmed this structure with multi-messenger precision. The gravitational field has spatially distributed, locally-coupled degrees of freedom: the disturbance propagated from point to point, each region of the field influencing its neighbors across 130 million light-years.

The ocean makes this concrete. Water waves propagate because water molecules push their neighbors; the disturbance passes from molecule to molecule across thousands of kilometers. Water molecules must form closed electron shells, and this closure creates tetrahedral hydrogen-bonding structures, the same four-fold topology that appears in tetrad gravity. In both systems, three independent spatial directions from a common origin define a tetrahedron: field closure in three dimensions requires no fewer than four faces.

The wave equation encodes this requirement mathematically. Gravitational perturbations satisfy:

$$\frac{\partial^2 h}{\partial t^2} = c^2 \nabla^2 h \tag{3}$$

The Laplacian ∇^2 appears on the right-hand side. By definition, this operator requires evaluating the field at neighboring points:

$$\nabla^2 \Phi = \lim_{\epsilon \to 0} \frac{\Phi(x+\epsilon) + \Phi(x-\epsilon) - 2\Phi(x)}{\epsilon^2} \tag{4}$$

The logic is inescapable: gravitational waves propagate (LIGO observed this). Propagation requires ∇^2 (the wave equation (**??**) demands it). ∇^2 requires neighbors (by definition). Therefore, gravity must consist of distributed, locally-coupled degrees of freedom. Figure **??**

illustrates this schematically: each field region overlaps with its immediate neighbors, coupling through shared field volume, and the pattern of these couplings constitutes the field's topology.




FIG. 1. Two-dimensional cross-section illustrating the coupling topology of gravitational field regions. Each circle represents a field region; where adjacent regions overlap (red shading), the fields couple through shared volume. One interior region (red) and its $k = 4$ nearest neighbors (orange) are highlighted, with the four coupling zones visible as lens-shaped intersections. In three spatial dimensions, the simplex theorem (Section **??**) requires $k = d + 1 = 4$ coupling faces per region; these four coupled faces enclose a tetrahedral volume, the field chamber.

The question is not *whether* gravity has topology; LIGO and tetrad theory both confirm it does. The question is: *what is* that topology?

B. Field Closure: The Requirement GR Already Enforces

If gravity consists of distributed, locally-coupled degrees of freedom, how do these degrees of freedom achieve consistency?

Consider what "locally-coupled" means. At each point, the field must define a local reference frame: which direction is timelike, which directions are spacelike. Neighboring points must agree on how their frames relate. If point A says "my timelike direction points this way" and neighboring point B says "that same direction is spacelike," physics would be undefined at the boundary.

For the field to be well-defined everywhere, the local frames must form a complete, non-

singular structure. At every point, all four directions (one time, three space) must be specified. No direction can be missing; no direction can be collapsed. This requirement, that the frame structure be complete everywhere, is what we call **field closure**.

This is not a new requirement. General Relativity has enforced it for a century, using different language. In metric GR, the condition is called non-degeneracy: $\det(g_{\mu\nu}) \neq 0$. Without this condition, the inverse metric $g^{\mu\nu}$ is undefined, indices cannot be raised, and the field equations cannot be written. Field closure and non-degeneracy are two names for the same physics.

The tetrad formalism reveals what this condition means geometrically [**?** **?**]. The equivalence principle requires physics to be locally Minkowski at every point:

$$g_{\mu\nu} = \eta_{ab}\, e^a_\mu\, e^b_\nu \tag{5}$$

where $\eta_{ab} = \text{diag}(-1, +1, +1, +1)$ is the Minkowski metric and e^a_μ are four orthonormal basis vectors defining the local frame. Taking determinants:

$$\det(g) = -[\det(e)]^2 \tag{6}$$

Therefore $\det(g) \neq 0$ if and only if $\det(e) \neq 0$: the four tetrad legs must span the tangent space completely. **GR's non-degeneracy condition and field closure are mathematically equivalent.**

In simplicial quantum gravity, Barbieri [**?**] showed that the analogous closure constraint (four face normals summing to zero) uniquely defines a tetrahedron at each node of the gravitational field. The continuous condition $\det(e) \neq 0$ and the discrete condition $\sum \vec{n}_i = 0$ are two expressions of the same geometric requirement: four independent directions must span a complete structure.

Concept	GR Language	Tetrad Language
Field defined everywhere	Non-degenerate metric	Field closure
Mathematical condition	$\det(g) \neq 0$	$\det(e) \neq 0$
Structure	3+1 decomposition	4 tetrad legs
Constraints	4 ADM equations	4 projections onto legs

TABLE I. GR non-degeneracy and field closure are two names for the same requirement. Practitioners of General Relativity have enforced field closure for a century.

Field closure is the spark plug of the framework: a binary condition. Either the field encloses volume ($\det(e) \neq 0$) or spacetime has collapsed ($\det(e) = 0$). This is the $k^0 = 1$ term in the coupling polynomial (Section **??**). It is always on, independent of direction, orientation, or dynamics. It is the precondition for everything else.

Field closure converts a qualitative demand ("spacetime must not degenerate") into a quantitative constraint: how many faces must a local field region have to guarantee enclosure in d dimensions? The answer is $d + 1$, which leads directly to k.

C. Why $k = 4$: The Uniqueness of Four

Before examining the mathematics, consider what $k = 4$ means physically. The parameter k counts how many neighbors each field region must touch to achieve field closure. In three-dimensional space, what is the minimum number of neighbors required to enclose a region?

Nature answers this question everywhere we look. Stack oranges at a grocery store: in close-packing each orange touches 12 neighbors, but the gaps between groups of three touching spheres form tetrahedral pockets, the simplest three-dimensional cavities. Examine water molecules: each forms hydrogen bonds with up to four neighbors, creating the tetrahedral network that gives water its remarkable properties. Study crystal structures: atoms pack in arrangements where tetrahedral coordination dominates. In 1611, Kepler conjectured that spheres in three dimensions pack most efficiently in arrangements with tetrahedral local structure; Thomas Hales proved this rigorously in 1998 [**?**]. Four-fold coordination is not a

choice. It is what three-dimensional space requires for efficient packing.

The simplex theorem [**?**] formalizes this observation: the minimum convex polytope enclosing volume in d dimensions has $d+1$ faces. For $d=3$ spatial dimensions, this gives 4 faces: a tetrahedron. The tetrahedron is the simplest possible three-dimensional container. Just as a triangle is the minimum polygon in 2D (you cannot enclose area with fewer than 3 sides), a tetrahedron is the minimum polyhedron in 3D (you cannot enclose volume with fewer than 4 faces). The tetrahedron is to 3D gravity what the triangle is to 2D tiling: you do not choose it; three dimensions force it.

The value $k=4$ is squeezed from both sides by independent constraints:

From below ($k \geq 4$): Fewer than 4 constraints cannot close a 3D evolution problem. The simplex theorem establishes this geometrically: minimal enclosure in d dimensions requires $d+1$ boundary conditions.

From above ($k \leq 4$): The symmetric metric tensor in four dimensions has 10 independent components. The k constraint equations and k gauge freedoms together remove $2k$ components, leaving $10-2k$ propagating degrees of freedom. LIGO measured exactly two graviton polarizations [**?**]. For $k=4$, this gives $10-8=2$, matching observation. With $k=5$ or higher, $10-2k<2$: fewer than 2 propagating modes, violating the observation that gravity propagates.

Combining: $k \geq 4$ (field closure) and $k \leq 4$ (graviton polarizations). The only solution is $k=4$.

D. Why $d=3$: The Tiling Condition

The preceding subsection established $k=d+1$. This subsection establishes why $d=3$, removing the last degree of freedom.

The dual tetrahedron structure requires two simplices to tile a region of space. The face count of two simplices in d dimensions is $2(d+1)$, which grows linearly. The vertex count of the dual lattice is 2^d, which grows exponentially. For the dual structure to close consistently, these two counts must match:

$$2(d+1) = 2^d \tag{7}$$

Evaluating for small d: $d = 1$ gives $4 \neq 2$; $d = 2$ gives $6 \neq 4$; $d = 3$ gives $8 = 8$; $d = 4$ gives $10 \neq 16$. The linear and exponential functions cross at exactly one integer: $d = 3$. This is arithmetic, not fitting.

Two additional constraints independently select $d = 3$:

Orbital stability. Bertrand's theorem [**?**] establishes that closed orbits exist only for $1/r^2$ and r^2 force laws. In d spatial dimensions, the gravitational force falls as $1/r^{d-1}$. Closed orbits require $d - 1 = 2$, hence $d = 3$. In $d \geq 4$ dimensions, orbits are unstable: small perturbations send planets spiraling inward or outward.

Atomic stability. In d spatial dimensions, the hydrogen atom has no bound states for $d \geq 4$: the centrifugal barrier cannot prevent collapse of the electron into the nucleus. Stable matter requires $d \leq 3$.

Three different domains of physics, topological tiling, orbital mechanics, and quantum stability, intersect at one value. The dimension d is not a parameter; it is an intersection point. Parameters can be adjusted. The intersection of three constraints cannot.

E. The Characteristic Scale

The tetrad formalism specifies topology ($k = 4$ faces) but not scale. At what physical radius does this internal structure exist?

Two independent routes converge on the same characteristic length:

Route 1: Electromagnetic Stability. At some critical radius, the energy stored in the electron's electric field equals its rest mass energy $m_e c^2$. The radius where these two energies balance is the classical electron radius [**?**]:

$$r_e = \frac{e^2}{4\pi\varepsilon_0 m_e c^2} = 2.818 \times 10^{-15} \text{ m} \tag{8}$$

This calculation uses only electromagnetism (e, ε_0) and special relativity (c). No gravity. No cosmology.

Route 2: Gravitational Topology. When gravity couples through the dual tetrahedron structure, all $k^2 = 16$ coupling channels engage (Section **??**). The characteristic radius

emerges when the tetrad's gravitational acceleration equals the cosmic threshold a_0:

$$r_e = k\sqrt{\frac{Gm_e}{a_0}} = 2.82 \times 10^{-15} \text{ m} \tag{9}$$

This calculation uses gravity (G), cosmology (a_0), and topology ($k = 4$). No electromagnetism.

Route	**Constants used**	**Result**	**Domain**
EM stability	e, ε_0, m_e, c	$r_e = 2.818$ fm	Electromagnetism
Cosmic topology	G, H, m_e, k, c	$r_e = 2.82$ fm	Gravity + Cosmology

TABLE II. Two independent routes converge on the same scale to 0.21%.

Match: **0.21%**. Two independent domains of physics, using different constants and different reasoning, converge on the same scale. If the same topology governs both gravity and electromagnetism, the characteristic scale must be shared. The fact that it is confirms the hypothesis quantitatively.

Once the scale is identified, the fine-structure constant becomes geometric:

$$\alpha = \frac{r_e}{\bar{\lambda}_C} \qquad \text{where } \bar{\lambda}_C = \frac{\hbar}{m_e c} \tag{10}$$

Alpha is the ratio of the field scale (where the topology lives) to the quantum scale (where coherence sets the wavelength). It is the lattice spacing divided by the quantum coherence length. Alpha is not a running coupling that happens to be near 1/137; it is a topological ratio.

Summary. The theoretical foundations are established in the literature: the gravitational field propagates (LIGO), propagation requires field topology (wave equation), field topology requires field closure (non-degeneracy), field closure requires $k = 4$ faces (simplex theorem), the spatial dimension is $d = 3$ (tiling condition, orbital stability, atomic stability), and the scale is the electron radius (convergence). What remains is to show the structure that these constraints produce, and what it yields.

III. THE DUAL TETRAHEDRON

Section **??** established that field closure requires $k = 4$ faces and $d = 3$ spatial dimensions. A single tetrahedron satisfies the face count, but not the consistency requirements. This section shows that four independent arguments require a second tetrahedron with opposite orientation, introduces the Field Origin where they couple, and derives the face structure and geometric elements that determine the coupling hierarchy.

A. Four Roads to Two Tetrahedra

Section **??** established that field closure requires $k = 4$ faces and $d = 3$ spatial dimensions. A single tetrahedron satisfies the face count. But is a single tetrahedron sufficient for a consistent gravitational field? Four independent arguments from the literature, spanning five decades and over 2,300 combined citations, all establish that it is not:

1. **Stability** (Alexander, Marcianò, Smolin 2014 [**?**]): A single tetrahedral field chamber has a preferred direction, an apex pointing somewhere. This asymmetry produces a *ghost*: a field mode with negative kinetic energy that destabilizes the vacuum. Two opposite field chambers cancel the asymmetry, and the ghost disappears.

2. **Spinor structure** (Weinberg 1995 [**?**]; Geroch 1968 [**?**]): Fermions require the SL(2,$\mathbb{C}$) double cover of the Lorentz group. Rotating a fermion 360 degrees does not return it to its original state; a full 720-degree rotation is needed. This is bimetric structure: one 360-degree rotation brings TetA back into alignment but leaves TetB out of phase; the second rotation realigns both. Each tetrahedron realizes one sheet of the double cover.

3. **Chirality** (Ashtekar 1986 [**?**]): Gravity decomposes into self-dual and anti-self-dual components, $A^+ \oplus A^-$, like left-handed and right-handed gloves that are mirror images but cannot be superimposed. Full gravitational field closure requires both halves. Each half corresponds to one tetrahedron.

4. **Bimetric gravity** (Hassan, Rosen 2012 [**?**]; Hinterbichler, Rosen 2012 [**?**]): Ghost-free bimetric gravity requires two dynamical metrics, each decomposed into its own

tetrad field: $g_{\mu\nu} = \eta_{ab}\, E^a_\mu\, E^b_\nu$ and $f_{\mu\nu} = \eta_{ab}\, L^a_\mu\, L^b_\nu$. The spatial triad of each tetrad defines a tetrahedron [**?**]; two tetrads define two tetrahedra. The two fields couple through an interaction potential $V(\mathcal{S}) = \sum_{n=0}^{4} \beta_n\, e_n(\mathcal{S})$, where $\mathcal{S} = \sqrt{g^{-1} f}$ is a 4×4 matrix and β_0 through β_4 are free coefficients [**?**]. Hassan and Kocic [**?**] proved that the two metrics must share compatible (3+1) decompositions. The theory is ghost-free at the full nonlinear level, with seven propagating degrees of freedom.

When four independent roads converge on the same destination, the destination is not hypothesized. These are textbook results from different subfields: stability analysis, quantum field theory, canonical gravity, and massive gravity. All four independently require the same field topology: two tetrahedra with opposite orientations, coupled at a shared center, the Field Origin.

Each argument constrains the structure in a specific way. Stability constrains *orientation*: the tetrahedra must point in opposite directions. Spinor structure constrains *topology*: the two tetrahedra must be genuinely distinct, realizing the two sheets of the SL(2,$\mathbb{C}$) double cover. Chirality constrains *completeness*: each tetrahedron carries one half of gravity. Bimetric structure constrains *interaction*: the two tetrad fields must couple at a single Field Origin [**?**], with opposite spatial orientations, because same-orientation coupling produces the Boulware-Deser ghost [**?**].

Together, these constraints uniquely determine the arrangement: two tetrahedra, opposite orientations, interpenetrating so that each vertex of one lies at the center of a face of the other. This is the stellated octahedron [**?**], the unique compound of two tetrahedra satisfying all four requirements. The resulting figure has:

- 8 vertices (4 from each tetrahedron)
- 12 edges (6 from each tetrahedron)
- 8 triangular faces (4 from each tetrahedron)
- A central octahedral region where the two tetrahedra overlap
- Inversion symmetry: reflecting any point through the center ($\mathbf{r} \to -\mathbf{r}$) maps the structure onto itself, exchanging TetA and TetB

These 28 geometric elements $(8 + 12 + 8)$ constitute the complete topological scaffold. This count will prove significant when the coupling hierarchy is read for its total breadth rather than its depth (Section **??**).

Other arrangements, tetrahedra side by side, apex to apex, or partially overlapping, either break the central symmetry required by bimetric gravity or fail to provide the complete spinor doubling. Only the stellated octahedron has both the dual orientation needed for stability and the shared coupling point needed for bimetric interaction. The structure is not chosen; it is required.




FIG. 2. The stellated octahedron (Dual Tetrahedron): two tetrahedra with opposite orientations sharing a common center. TetA (thick lines) points upward; TetB (thinner lines) points downward. The structure has 28 geometric elements: 8 vertices, 12 edges, and 8 faces. At the geometric center lies the Field Origin, where the two spacetime sectors couple. This is the dual tetrad topology from which the fundamental equations derive.

B. The Field Origin

Bimetric gravity requires that two dynamical metrics interact. The Hassan-Rosen formulation [**?**] and the vielbein extension by Deffayet, Mourad, and Zahariade [**?**] establish that the two spin-2 fields must couple at a shared spacetime point, what the literature calls an *interaction vertex*. Without such a vertex, the two metrics would be causally disconnected:

two separate gravitational fields with no physical relationship. We call this point the **Field Origin**.

The stellated octahedron provides exactly this Field Origin. At the geometric center, equidistant from all eight vertices, the two tetrahedra share a single point. This is where the two spacetime sectors meet, where information from one tetrahedron can couple to the other.

Think of the Field Origin as the neck of an hourglass. Sand flows from one chamber through the neck into the other; nothing can pass between the two halves except through that single constriction. The Field Origin is the topological bottleneck through which all coupling between TetA and TetB must pass.

Why does time vanish at this point? Consider the light-travel time from the coupling center to any point at distance r:

$$\tau_{\text{prop}} = \frac{r}{c} \tag{11}$$

At the center itself, $r = 0$, so $\tau_{\text{prop}} = 0/c = 0$. The propagation time vanishes. But this equation says more than "time is zero here." Read in the other direction, $\tau_{\text{prop}} = r/c$ means time *increases* with distance from the origin. Time radiates outward. The Field Origin is not merely a point where time happens to vanish; it is the source from which all temporal separation emerges. The analogy is the eye of a hurricane: perfectly still, yet generating the entire storm.

This property is what makes the Field Origin physically meaningful. The two tetrahedra represent two separate spacetime sectors with opposite orientations. For ghost-free coupling, they must meet at a point where their interaction can be instantaneous, unmediated by propagation delay. The Field Origin, where $\tau_{\text{prop}} = 0$, satisfies this requirement. But why is instantaneous coupling physically necessary?

Consider a field object: a macroscopic object such as an asteroid composed of $\sim 10^{33}$ coupled fields. This is not 10^{33} separate entities; it is one unified field configuration: iron atoms with their binding fields, crystalline lattice with its modes, all coupled into a single gravitational entity. We refer to such unified configurations as *field objects*. Like a musical chord, where the relationships between notes define the chord (not the notes individually), a field object is defined by the correlations between its components. You cannot play half a chord; the chord exists only when all notes and their relationships are present. Similarly, FC_Σ cannot

couple to part of a field object. It must read the entire unified configuration in a Full State Snapshot.

Here is the constraint: FC_Σ propagates at speed c between spatially separated field regions, confirmed by gravitational wave observations to fifteen decimal places [**?**]. If FC_Σ within the field object configuration were also limited by this propagation speed, the field could not achieve Full State Snapshot. Information would require time $\tau = r/c$ to traverse the object's spatial extent r, preventing the atomic read that FC_Σ requires. The unified configuration could not settle simultaneously; by the time one part was read, other parts would have evolved. FC_Σ is atomic: partial reads produce inconsistent state.

Why must this read be atomic? Three requirements from different areas of physics converge on the same answer. Textbook general relativity requires $\det(g) \neq 0$ everywhere [**?** **?**]; this non-degeneracy condition ensures the Laplacian operator, which appears throughout gravitational physics and is defined by neighbor comparison, can function. Distributed field values must reach consistency simultaneously. Hassan and Kocic [**?**] proved that bimetric gravity requires compatible (3+1) decompositions with ADM constraints satisfied simultaneously across all degrees of freedom: two coupled fields settle together, not sequentially. Observation confirms what theory requires: gravitational fields exhibit no wakes, drag, or hysteresis as objects move through them. The coupling is lossless and history-independent. Partial settlement would leave observable traces; we see none. These three requirements (GR's constraint structure, bimetric well-posedness, and observational fact) establish that the field configuration must settle simultaneously into a globally consistent state. This is what we call a Full State Snapshot of the field object configuration. The Field Origin, which is the interaction vertex of the Hassan-Rosen bimetric Hamiltonian, provides the mechanism that satisfies these requirements. Where both tetrad fields couple at $r = 0$ through the nonlinear interaction, propagation time vanishes ($\tau_{\mathrm{prop}} = 0$) and both field configurations can settle simultaneously, exactly the Full State Snapshot that field closure demands.

The Field Origin resolves this through topology, not propagation. At $r = 0$, where $\tau_{\mathrm{prop}} = 0$, field closure can complete without waiting for signals to propagate across space. In the Hassan-Rosen bimetric formalism, this is the interaction vertex where both metric fields couple through the interaction potential—both fields are evaluated at the same spacetime point, so no propagation is needed. Fields still propagate at speed c, but the Field Ori-

gin provides a nonlinear coupling boundary where propagation time vanishes, enabling the atomic read that field closure requires.

Why everything spins

One of the deepest questions in physics: why does everything rotate? Stars spin. Galaxies spin. Electrons spin. Angular momentum pervades the universe at every scale.

Einstein-Cartan theory, developed in 1922, provides a hint [**?**]. Cartan showed that spacetime geometry and intrinsic spin are intimately connected: the spin of matter couples directly to spacetime torsion. Where there is torsion, there is spin. Where there is spin, there is torsion.

The dual tetrahedral field structure makes this connection concrete. When two tetrahedra with opposite orientations couple at a shared point, they create a continuous rotational field topology. Consider the analogy of water flowing in a river whirlpool (Figure **??**, left panel). At the surface, water spirals inward. As it approaches the center, it accelerates, passes through the central axis, and disperses below. The vortex core is not where motion stops; it is where rotation vanishes but through-flow continues.




FIG. 3. The Field Origin as a vortex core. **Left:** A river whirlpool viewed from the side. Water spirals inward at the surface, passes through the vortex core, and disperses below. **Right:** The dual tetrahedral field structure. TetA (purple, top) with apex-up orientation, the Field Origin (red, center) at the coupling point where $\tau_{\text{prop}} = 0$, and TetB (blue, bottom) with apex-down orientation. Angular momentum does not stop at the Field Origin but flows through it.

The dual tetrahedral field structure shares this vortex topology. Angular momentum propagates through both spacetime sectors, passing through the Field Origin that connects them. At the Field Origin itself, the rotational component vanishes (because $\tau_{\text{prop}} = 0$ there), yet angular momentum continues through. This is the topological origin of spin: angular momentum flowing through the neck of the hourglass.

How much angular momentum passes through this vortex eye? The answer depends on the internal face structure of the stellated octahedron. Spin lives at the Field Origin, and its value is determined by the partition of faces between the two tetrahedral sectors. Angular momentum divides 3:2:3 across TetA (3/8), the Field Origin (2/8), and TetB (3/8). The fraction flowing through the Field Origin is $(2/8) \times 2\hbar = \hbar/2$. Spin is not a property added to matter from outside. It is the angular momentum flowing through the topological connection between the two tetrahedra. Section **??** develops this in full.

C. The Face Structure and Spin

The 3+1 split of spacetime is often treated as a coordinate choice, a matter of convenience for the ADM formalism. In the stellated octahedron, it is a counting fact about the solid. The faces tell us what spacetime is made of.

Each tetrahedron has 4 triangular faces. The stellated octahedron, composed of two interlocking tetrahedra, therefore has 8 faces in total. These 8 faces divide into two types based on their orientation relative to the Field Origin:

- **TIME faces** (2 total): In each tetrahedron, the face opposite the apex points inward, toward the geometric center where the Field Origin lies. The base of TetA (apex-up) faces downward toward the center, and the base of TetB (apex-down) faces upward toward the center. These two faces point toward the coupling point where $\tau_{\text{prop}} = 0$. They are called TIME faces because they connect to the temporal origin.

- **SPATIAL faces** (6 total): The remaining three faces of each tetrahedron share the apex vertex and point outward, away from the Field Origin into the spatial dimensions. Three faces from TetA radiate upward and outward; three faces from TetB radiate downward and outward.

Why exactly 2 faces point inward is a geometric fact, not a parameter. A tetrahedron has one apex and one base. The base, the single face opposite the apex, points toward the center of the stellated octahedron. The other three faces share the apex and point away. With two tetrahedra, two bases point inward and six apex-sharing faces point outward. The ratio 6:2 (spatial:temporal) is the geometric origin of spacetime's (3+1) signature. It is a counting fact about the solid, not a postulate about spacetime.

Electron spin from the same topology

The face structure determines not only the signature of spacetime but also the spin of the fundamental fermion. Spin is computed from the topology, not added as an external quantum number.

Route 1: The face ratio. The stellated octahedron has 8 faces: 2 TIME faces pointing inward toward the Field Origin, 6 SPATIAL faces pointing outward. Each tetrahedron carries quantized angular momentum $\hbar$ (one closed circuit through 4 faces); the stellated octahedron carries total angular momentum $2\hbar$. Only the 2 TIME faces point toward the vortex eye; the 6 SPATIAL faces carry orbital angular momentum that does not pass through the Field Origin. The observable intrinsic spin is:

$$S = \frac{N_{\text{TIME}}}{N_{\text{total}}} \times L_{\text{total}} = \frac{2}{8} \times 2\hbar = \frac{\hbar}{2} \tag{12}$$

This is the vortex aperture argument: the Field Origin admits angular momentum only through the 2 faces that point toward it.

Route 2: The structural state differences. The structural states (Section **??**) reveal a second, independent path. In TetA ($s = +1$), the full hierarchy has 21 field coupling levels. At the Field Origin ($s = 0$), 17 remain. The $21 - 17 = 4$ that vanish are the face channels that require spatial extent. Similarly, $17 - 13 = 4$ channels belong to TetB's sector. The total face contribution across both sectors is $4 + 4 = 8 = 2k$. An observer in our (3+1) spacetime has access to only one sector's face channels: 4 of the 8. With quantized angular momentum $\hbar$ per closed circuit through k faces:

$$S = \frac{k}{2k} \times \hbar = \frac{4}{8} \times \hbar = \frac{\hbar}{2} \tag{13}$$

Two independent readings of the same field topology, one topological (counting faces) and one algebraic (counting coupling channels from the polynomial), both yield $\hbar/2$. The intermediate numbers differ (2/8 versus 4/8, $2\hbar$ versus $\hbar$), but both arrive at $\hbar/2$. Their agreement from independent starting points is a self-consistency check on the topology. The 1/2 follows from the bimetric structure: an observer occupies one of two coupled spacetime sectors.

Calibration note. Both routes assume that the total angular momentum per closed circuit is $\hbar$, the experimentally established quantum of angular momentum. The topology determines the fraction (1/2); the magnitude $\hbar$ is an empirical input, not derived here.

D. The 28 Geometric Elements

Count what you can build from the stellated octahedron. The answer is 28, and this number enters the most precisely measured constant in physics.

The total count of geometric elements is:

$$V + E + F = 8 + 12 + 8 = 28 \tag{14}$$

This can be verified by building a physical model. Eight vertices, twelve edges, eight triangular faces. The count is geometry, not conjecture.

What makes 28 significant is not that it appears once but that it appears three times, from three independent countings:

Counting 1: Geometric elements. As above, $V + E + F = 8 + 12 + 8 = 28$. This counts every distinct structural feature of the stellated octahedron.

Counting 2: Face-channels. Each of the 8 faces of the stellated octahedron has $k - 1 = 3$ independent coupling channels (a triangular face has 3 vertices, and excluding the Field Origin leaves $k - 1$). But the 2 TIME faces are shared between sectors and contribute a single coupling each. An effective count gives 7 independent face types times 4 channels per face type: $7 \times 4 = 28$. More precisely, $k(k + d) = 4 \times 7 = 28$, where $k + d = 7$ is the number of effective face modes.

Counting 3: Bimetric degrees of freedom. Ghost-free bimetric gravity has 7 propagating degrees of freedom (2 from a massless graviton, 5 from a massive graviton). Each

propagating mode phase-locks across 4 face channels: $7 \times 4 = 28$.

Three different enumerations, geometric, face-based, and dynamical, all yield 28. This triple convergence is an algebraic identity: $V+E+F = k(k+d)$ for the compound of two simplices in d dimensions. It is not a coincidence to be explained away; it is the statement that the geometric scaffold, the face coupling structure, and the dynamical degrees of freedom all describe the same object counted from different perspectives.

This number enters the fine-structure constant. The fractional part of $1/\alpha$ is:

$$\frac{1}{\alpha} = 137 + \frac{1}{28} = 137.0357\ldots \tag{15}$$

The most precisely measured constant in physics carries, in its decimal tail, the element count of the stellated octahedron. Section **??** derives this in full. When the fraction is derived from geometry and matches measurement to 0.00021%, the geometry is telling the truth.

IV. THE COUPLING POLYNOMIAL

Section **??** established the dual tetrahedron: two interlocking tetrahedra sharing a Field Origin. This section develops the coupling polynomial $f(k) = 1 + k + k^2$, the mathematical heart of the paper. The polynomial counts the depth of field coupling within the stellated octahedron: how many levels the gravitational field must traverse to achieve complete coupling between the two sectors. Every derivation in subsequent sections flows from this polynomial and the three integers it produces.

A. Why the Polynomial Has Three Terms

Field coupling between the two tetrahedra does not happen all at once. It has levels, each contributing to the total interaction strength. Consider what must connect for gravity to operate. The answer is a causal sequence: you cannot propagate what does not exist, and you cannot interact without first reaching. The polynomial's three terms are not a mathematical convenience; they are the three stages of this causal chain.

1. **EXIST: The Field Origin ($k^0 = 1$).** Field coupling begins at the origin, where the two tetrahedra meet. This is the foundation: without the Field Origin, there is no coupling at all. This term is binary: either the field encloses volume ($\det(e) \neq 0$) or spacetime has collapsed ($\det(e) = 0$). It is always on, independent of direction, orientation, or dynamics.


2. **PROPAGATE: One coupled face ($k^1 = 4$).** From the Field Origin, the field propagates through one face of the tetrahedron, which subdivides into $3 + 1 = 4$ subregions (three spatial, one coupling) as shown in Figure **??**. This is how the field extends through one sector of the dual tetrahedral field chamber.

3. **INTERACT: The full bimetric Field Coupling ($k^2 = 16$).** Both tetrahedra interact fully. Each has (3+1) dimensions, so the complete field coupling space is (3+1) $\times$ (3+1) = 16-dimensional. This is the 4×4 frame transition matrix $\mathcal{S}$ of Hassan-Rosen bimetric gravity [**?**].

The causal ordering is forced: EXIST before PROPAGATE before INTERACT. You cannot propagate what does not exist. You cannot interact without first reaching. This is not physics; it is logic. The causal sequence *is* the polynomial $f(k) = 1 + k + k^2$.

The total field coupling hierarchy is:

$$f(k) = 1 + k + k^2 = 1 + (3{+}1) + (3{+}1)^2 = 1 + 4 + 16 = 21 \tag{16}$$



(a) Face of TetA **(b)** Coupled face: $3 + 1$

FIG. 4. The Field Coupling topology of a single face. **(a)** A triangular face of TetA. **(b)** When TetB couples through the Field Origin, its vertex V_B penetrates the face center, subdividing the face into $3 + 1 = 4$ sub-regions: three corner sub-regions (blue) anchored at the original vertices, and one central coupling sub-region (red) at the penetration point. The $3 + 1$ structure recurs at every scale of the hierarchy.

Why there is no fourth term. A k^3 term would represent a 3-dimensional coupling cell on the boundary of the tetrahedron. But coupling occurs on the *surface* of the simplex, which is 2-dimensional (triangular faces). There is no 3-cell on the boundary of a tetrahedron. Numerically, adding $k^3 = 64$ would give an exponent of 85, and α^{85} is effectively zero: the coupling would contribute nothing observable. The polynomial has exactly three terms because the simplex surface has exactly three types of cells: 0-cells (vertices), 1-cells (edges), and 2-cells (faces).

The gravitational field traverses all 21 coupling levels. Each level attenuates by the fine-structure constant α (Section **??** derives α as the topology's own scale ratio), giving the coupling depth α^{21} in the expression for G. The number 21 is derived from the field topology, not imported from elsewhere. Having established what each term in the polynomial represents, Section **??** now demonstrates why this specific form is the unique solution to three independent physical constraints.

B. Why This Form Is Unique

Section **??** explained what each term in the polynomial represents: existence (1), propagation (k), and bimetric interaction (k^2). The natural question follows: why must the polynomial take precisely this form? Could it be $1 + k$ alone, or contain additional terms such as k^3

or k^4? This subsection demonstrates that $f(k) = 1 + k + k^2$ is the unique solution to three independent physical constraints established in Sections **??** and **??**. We present two complementary derivations: one proves no other polynomial satisfies the constraints, the other derives this polynomial directly from component counting.

Constraint-based uniqueness

Three established constraints from observational physics and geometric necessity uniquely determine the polynomial form.

Constraint 1: LIGO graviton polarizations. The LIGO/Virgo collaboration measured exactly two tensor polarizations in gravitational waves from binary neutron star merger GW170817 [**?**]. The metric tensor $g_{\mu\nu}$ is a symmetric 4×4 matrix with 10 independent components. Physical degrees of freedom equal total components minus constraints minus gauge freedoms. Observing 2 polarizations gives $10 - 2k = 2$, which uniquely determines $k = 4$.

Constraint 2: Simplex theorem. In d dimensions, exactly $d + 1$ points are required to enclose a bounded region [**?**]. This is geometric necessity, not convention. Fewer points cannot form a closed boundary; additional points are redundant. For 3-dimensional space, $k = d + 1 = 3 + 1 = 4$. The tetrahedron (4 vertices, 4 faces) is the minimal 3D solid. Any coupling structure in 3D space must have $k = 4$ faces.

Constraint 3: Ghost-free bimetric coupling. Hassan-Rosen bimetric gravity eliminates the Boulware-Deser ghost by introducing a second metric tensor [**?**]. The symmetric vielbein condition $g^{\mu\nu} e^A_\mu e^B_\nu = g^{\mu\nu} e^B_\mu e^A_\nu$ requires the two tetrads to couple through a common point where their interaction satisfies ghost-free constraints. Geometric analysis demonstrates this coupling point must lie at the center of a dual tetrahedral structure where both tetrads meet. The two tetrads have opposite orientations (apexes at antipodal points), which is the geometric condition for ghost elimination.

These three constraints force the polynomial structure. The "1" term represents the Field Origin where both tetrads couple at $r = 0$ (satisfying $\det(e) \neq 0$ for field existence). The "k" term represents propagation through $k = 4$ vierbein faces within one tetrad (required by the simplex theorem for 3D space). The "k^2" term represents bimetric interaction: k faces

in TetA coupling to k faces in TetB gives $k \times k = 16$ total channels (required by ghost-free bimetric theory). No other polynomial can satisfy all three constraints simultaneously. A polynomial lacking the "1" term violates the existence condition. A polynomial lacking the "k" term cannot describe propagation through spatial channels (contradicting Constraint 2). A polynomial lacking the "k^2" term cannot capture bimetric coupling (contradicting Constraint 3). Additional terms such as k^3 or k^4 would contradict Constraint 1, which uniquely determines $k = 4$, and Constraint 2, which establishes 4 as the minimum for 3D enclosure.

The form $f(k) = 1 + k + k^2$ is therefore the unique solution.

Component-counting derivation

The polynomial can be derived independently by counting coupling channels in the bimetric ADM Hamiltonian.

The vierbein e^a_μ connects coordinate basis dx^μ to orthonormal tetrad basis θ^a. It carries two indices: frame index $a \in \{0, 1, 2, 3\}$ (4 values: 1 timelike plus 3 spacelike directions) and coordinate index $\mu \in \{0, 1, 2, 3\}$ (4 spacetime coordinates). Total tetrad components: $k \times k = 4 \times 4 = 16 = k^2$. The metric tensor is reconstructed via the tetrad as $g_{\mu\nu} = \eta_{ab}\, e^a_\mu\, e^b_\nu$, where $\eta_{ab} = \mathrm{diag}(1, -1, -1, -1)$ is the Minkowski metric. The summation over frame indices a and b activates all $k^2 = 16$ coupling channels.

The Hassan-Rosen bimetric interaction Hamiltonian takes the form [**?**]:

$$H_{\mathrm{int}} = m^2 \sqrt{g} \sum_{n=0}^{4} \beta_n\, e_n(\mathcal{S}) \tag{17}$$

where $\mathcal{S}^a_b = (e^{-1} f)^a_b$ is the frame transition matrix linking the two tetrads, and $e_n(\mathcal{S})$ are elementary symmetric polynomials in the eigenvalues of $\mathcal{S}$. Each e_n counts n-fold index contractions:

$$e_0(\mathcal{S}) = 1 \quad \text{(existence condition: } \det(e) \neq 0) \tag{18}$$

$$e_1(\mathcal{S}) = \mathrm{Tr}(\mathcal{S}) = \sum_a \mathcal{S}^a_a \quad \text{(single-index sum: } k \text{ channels)} \tag{19}$$

$$e_2(\mathcal{S}) \propto \sum_{a,b} \mathcal{S}^a_a\, \mathcal{S}^b_b \quad \text{(pairwise coupling: contains } k^2 \text{ terms)} \tag{20}$$

The polynomial $f(k) = 1 + k + k^2$ directly mirrors this structure: field existence (1), single-index propagation (k), and double-index bimetric coupling (k^2). This is not pattern matching; it is component counting.

If the dual tetrad topology correctly identifies the gravitational field structure, the coupling ratio $a_0/(Gm_e/r_e^2)$ should equal $k^2 = 16$, where $a_0 = 1.22 \times 10^{-10}$ m/s^2 is the MOND acceleration scale, $G = 6.674 \times 10^{-11}$ m^3/(kg·s^2) is Newton's constant, $m_e = 9.109 \times 10^{-31}$ kg is the electron mass, and $r_e = 2.818 \times 10^{-15}$ m is the classical electron radius. Direct calculation gives:

$$\frac{a_0}{Gm_e/r_e^2} = \frac{1.22 \times 10^{-10}}{7.66 \times 10^{-12}} = 15.93 \tag{21}$$

The predicted value is $k^2 = 16$. The measured value is 15.93. Agreement: 0.41%. This empirical verification confirms that counting tetrad components yields the correct polynomial form.

Synthesis

Two independent derivation paths converge on $f(k) = 1 + k + k^2$. The first demonstrates uniqueness through constraint satisfaction: three established physical requirements (LIGO measurement determining $k = 4$, simplex theorem requiring $k = 4$ for 3D enclosure, ghost-free bimetric coupling requiring two tetrads) uniquely determine the three-term polynomial structure. No other polynomial satisfies all three constraints simultaneously. The second demonstrates construction through component counting: the tetrad is a $4 \times 4 = 16$-component object, and the Hassan-Rosen Hamiltonian naturally contains terms e_0, e_1, and e_2, corresponding to existence (1), single-index propagation (k), and double-index coupling (k^2). Direct counting yields the polynomial form, and empirical verification of $k^2 = 16$ to 0.41% confirms the structure.

The polynomial $f(k) = 1 + k + k^2$ is therefore both necessary (no other form works) and sufficient (this form emerges from the topology). It is derived from the fundamental structure of bimetric gravity operating in 3-dimensional space with constraints set by observation, not identified by pattern matching.

The geometric series identity

The polynomial $f(k) = 1 + k + k^2$ has a revealing algebraic identity:

$$f(k) = \frac{k^3 - 1}{k - 1} = \frac{k^d - 1}{k - 1} = \frac{64 - 1}{3} = 21 \tag{22}$$

This is the standard geometric series with ratio k and $d = 3$ terms. The identity connects the polynomial directly to the dimension of space: $d = 3$ terms in the series because $d = 3$ spatial dimensions. The series sums $(k^d - 1)/(k - 1)$, which for a simplex in d dimensions counts the total number of coupling levels on the boundary. The connection between the polynomial and the dimension is not imposed; it is the definition of a geometric series on a simplex.

Clarification: the "1" versus the "+1"

The hierarchy contains a "1" and the spacetime structure contains a "+1." These are related but distinct.

Symbol	**What it represents**	**Where it applies**
The "+1" in (3+1)	Time as a dimension	Throughout spacetime
The "1" in the hierarchy	Time's origin (EXIST)	At $r = 0$, where $t = r/c = 0$

TABLE III. The distinction between time as dimension (the "+1" in 3+1) and the temporal origin (the "1" in $f(k)$).

Self-Similarity: The Requirement of Scale-Independent Closure

The polynomial structure $f(k) = 1 + k + k^2$ is not merely a property of a single tile: it is a requirement imposed by field closure under finite propagation speed.

The Physical Constraint. Field closure, expressed mathematically as $\det(e) \neq 0$, must hold at every scale. Gravitational waves propagate at speed c [**?**], a consequence of the hyperbolic structure of Einstein's equations in tetrad formulation [**?**]. Consider a field

region of characteristic size R. Closure at scale R must complete within the causal time $\tau = R/c$.

The Self-Similar Structure. At the fundamental tile scale r_e, the coupled face architecture $1 + k + k^2$ achieves closure. For closure to hold at scale $R > r_e$, the same architecture must operate at scale R, with each level recapitulating the full hierarchy. The alternative (a hierarchy that does not recurse) would produce a scale above which the field cannot close within the causal constraint set by c, causing $\det(e)$ to fail.

Therefore, the polynomial $f(k) = 1 + k + k^2$ serves as the recursion relation of a necessarily self-similar closure structure. Each coupled face at level n contains the full $(1 + k + k^2)$ structure at the next level down. This self-similarity is visible in the stellated octahedron's geometry: each of the 8 faces subdivides into 1 coupling vertex + 4 sub-regions + 16 cross-modes, reproducing the 21-level hierarchy at smaller scale.

Physical Consequence. The weight k^n at level n counts the number of closure channels active at that hierarchical depth. The self-similarity requirement guarantees that field closure, which we observe to hold throughout the universe, remains valid from the Planck scale to cosmological scales.

C. Why Each Coefficient Is One

The polynomial $f(k) = 1 + k + k^2$ has unit coefficients: each power of k appears exactly once. This is not a choice; it is forced by the structure.

Consider what each coefficient counts:

- $a = 1$ (coefficient of k^0): Existence is binary. The field either encloses volume ($\det(e) \neq 0$) or it does not ($\det(e) = 0$). There is one existence condition, not two. One Field Origin, not two.
- $b = 1$ (coefficient of k^1): Each face couples once. The connection between a field region and its neighbor is a single shared boundary. One connection per neighboring cell.
- $c = 1$ (coefficient of k^2): Each entry of the frame transition matrix counts once. The matrix $\mathcal{S}^a_b$ is a map from one frame to another, not a weighted sum. One matrix entry per pair of frame directions.

Each coefficient counts "how many of this type," and the answer is always one. Unit coefficients are the statement that the geometry is a simplex and the coupling is democratic: no face is privileged, no coupling mode is weighted.

The geometric series identity (Eq. **??**) provides the mathematical confirmation. The standard geometric series $\sum_{n=0}^{d-1} k^n = (k^d - 1)/(k - 1)$ has all coefficients equal to 1 by definition. The coupling polynomial is this geometric series with $d = 3$. If any coefficient departed from unity, the polynomial would no longer equal $(k^d - 1)/(k - 1)$, and the connection to simplex geometry would be broken.

This can be tested. Suppose $f(k) = 1 + 2k + k^2 = 25$. Then $f(k) \neq (k^3 - 1)/(k - 1)$, the polynomial no longer represents the dimensional sum on a simplex boundary, the geometric series identity fails, and the three structural states no longer satisfy the arithmetic mean relation (Section **??**). Departing from unity destroys the structural identity. The coefficients are 1 because the geometry is a simplex and the coupling is democratic. There is no freedom to adjust them.

D. The Three Structural States

The 21 coupling levels of Section **??** assume all channels are active. In the actual stellated octahedron, however, the coupling hierarchy is recursive: structure couples through origin to structure. Different structural regions see different channels because the Field Origin is the nonlinear coupling boundary between structures, not a structural region itself.

Of the polynomial's three terms, each maps onto a level of the recursive structure. The Field Origin (1) is the nonlinear coupling boundary where $r = 0$, present in every field configuration. The bimetric interaction ($k^2 = 16$) is a relationship between two structures at that boundary: if TetA couples to TetB, TetB couples to TetA, requiring no preferred direction. The middle term, $k = 4$, is different in kind: it represents field propagation through the four channels of one coupled face, the linear transport of the field through one sector of the stellated octahedron. Field propagation requires spatial extent ($r > 0$ and direction) and therefore exists only within a structural region, never at the coupling boundary.

The stellated octahedron has three distinct structural regions, and the state coefficient s


determines whether face channels contribute:

- In **TetA** $(s = +1)$, the field propagates outward through the four face channels with full coupling. The middle term contributes $+k$.
- At the **Field Origin** $(s = 0)$, the two tetrahedra couple at $r = 0$. This is the nonlinear coupling boundary, not a structural region: no spatial extent exists for face channels to operate. The middle term contributes 0.
- In **TetB** $(s = -1)$, the field propagates through the opposite sector's coupled face. The reversed orientation inverts the face contribution. The middle term contributes $-k$.

This structural distinction has a physical consequence. As established in Section **??**, the Field Origin enables coherent field configurations by providing a coupling point where $\tau_{\text{prop}} = 0$. Meanwhile, FC_Σ propagates causally at speed c through the $k = 4$ face channels. The three structural states map onto this physics: TetA and TetB represent propagation through spatial extent (causal, requiring $r > 0$), while the Field Origin represents the coupling boundary where coherence is established (structural, at $r = 0$). One is linear transport through channels; the other is nonlinear coupling at a point. The state coefficient s does not merely count active channels. It distinguishes whether the field operates in the propagation regime ($s = \pm 1$, where face channels carry information causally) or the coupling regime ($s = 0$, where the boundary enforces coherence). Both regimes are necessary: propagation governs how fields move through spacetime; coupling governs how fields settle into coherent configurations.

Evaluating $f(k) = 1 + s \cdot k + k^2$ in each structural state gives the three integers in G:

Structural state	**Polynomial**	**Value**	**Role in G**
TetA ($s = +1$, apex-up)	$1 + (3{+}1) + (3{+}1)^2$	$= 21$	exponent
Field Origin ($s = 0$)	$1 + \quad 0 \quad + (3{+}1)^2$	$= 17$	numerator
TetB ($s = -1$, apex-down)	$1 - (3{+}1) + (3{+}1)^2$	$= 13$	denominator



(a) EXIST: $k^0 = 1$ **(b) PROPAGATE:** $k^1 = 4$ **(c) INTERACT:** $k^2 = 16$

FIG. 5. The three topological components of $f(k) = 1 + s \cdot k + k^2$: **(a)** the Field Origin (1), **(b)** one coupled face ($k = 4$), and **(c)** the full stellated octahedron ($k^2 = 16$). The structural state s determines which components contribute.

All three integers in G emerge from one polynomial evaluated in the three structural regions of the stellated octahedron.

Why only the middle term changes

The structural argument is direct. Of the three terms, $k^0 = 1$ (a point) has no spatial extent to reverse. $k^2 = 16$ (a relationship) is symmetric: if TetA couples to TetB, then TetB couples to TetA regardless of orientation. But $k^1 = 4$ (propagation) has direction and spatial extent, and reversing the field orientation reverses that direction. Parity confirms this: under $k \to -k$, even powers (k^0, k^2) are unchanged, while the single odd power (k^1) changes sign.

The ghost-free condition fixes two opposite orientations (Section **??**), plus one boundary where they couple ($r = 0$), giving exactly three structural states. No fourth configuration exists. One polynomial, three regions, three integers.

The arithmetic mean lock

The three structural states are rigidly constrained by an algebraic identity:

$$\frac{f(+k) + f(-k)}{2} = \frac{21 + 13}{2} = 17 = f(0) \tag{23}$$

The Field Origin value is exactly the arithmetic mean of the TetA and TetB values. This holds for any polynomial of the form $1 + sk + k^2$; it is an algebraic identity, not a numerical coincidence. The three states are equally spaced: $21 - 17 = 17 - 13 = k = 4$. They are as rigidly constrained as three equally spaced marks on a ruler. There is no room to adjust one without breaking all three.

The prefactor: coupling capacity over spatial structure

The three structural states determine where each integer appears in G. The exponent 21 counts the full coupling depth from TetA, where all channels are active. The prefactor is a ratio comparing two structural states:

The numerator, 17, is the polynomial evaluated at the Field Origin ($s = 0$): $17 = k^2 + 1 = (3{+}1)^2 + 1$. At the coupling boundary, face channels are absent, leaving the 16 bimetric Field Coupling channels plus 1 for the non-degenerate coupling vertex itself. The numerator counts the coupling *capacity*.

The denominator, 13, is the polynomial evaluated in TetB ($s = -1$): $13 = kd + 1 = 3(3{+}1) + 1$. Each of the $k = 4$ face channels projects onto $d = 3$ spatial dimensions, giving $kd = 12$ spatial components plus 1 for non-degeneracy. The denominator counts the spatial *structure*.

$$\frac{17}{13} = \frac{k^2 + 1}{kd + 1} = \frac{(3{+}1)^2 + 1}{3(3{+}1) + 1} = \frac{\text{coupling capacity}}{\text{spatial structure}} \tag{24}$$

Neither is adjustable; both follow from $k = (3{+}1)$ and $d = 3$.

E. The Structural Identification

One polynomial, three structural states, the gravitational constant, electron spin, and a unique spatial dimension. Before deriving the equations, one question remains: the polynomial's bimetric interaction term $k^2 = 16$ is the same number as the 16 components of the frame transition matrix $\mathcal{S}$ in Hassan-Rosen gravity. Are these the same 16, or merely the same count?

The coupled face matrix

Each tetrahedron has 4 faces classified by orientation relative to the Field Origin: 3 spatial faces (pointing outward) and 1 time face (pointing inward). At the Field Origin, the coupled face matrix C_{ab} measures the coupling amplitude between face a of TetA and face b of TetB:

- Row index a: TetA's a-th face ($a = 0$ time, $a = 1, 2, 3$ spatial)
- Column index b: TetB's b-th face ($b = 0$ time, $b = 1, 2, 3$ spatial)
- Total: $4 \times 4 = 16$ coupling channels

The Hassan-Rosen frame transition matrix is $\mathcal{S}^a_{\ b} = (E^{-1})^a_{\ \mu}\, L^\mu_{\ b}$, where E^a_μ and L^a_μ are the vierbein fields of the two metrics. Its 16 components measure how each leg of one tetrad projects onto each leg of the other through the coupling point.

Proposition. *Under the conditions that (a) both tetrads share a common coupling point (the Field Origin of the Hassan-Rosen Hamiltonian), (b) the symmetric vielbein condition holds at that point (required for existence of a real square root [**?**]), and (c) compatible* (3+1) *decompositions exist (required for a well-posed Hamiltonian [**?**]), the coupled face matrix C_{ab} of the stellated octahedron is identically the Hassan-Rosen frame transition matrix $\mathcal{S}^a_{\ b}$.*

Proof by five identification conditions.

(i) Index identification. Barbieri [**?**] proved that at each 4-valent node of the gravitational field, the closure constraint $\sum_i \vec{N}_i = 0$ on four face normals uniquely defines a tetrahedron from the spatial triad. Each vierbein leg a therefore labels a face direction: $a = 0$ (timelike) corresponds to the time face, and $a = 1, 2, 3$ (the spatial triad) correspond to the three spatial faces. The Lorentz index on $\mathcal{S}$ and the face index on C label the same structural object. This correspondence is forced by the closure constraint, not chosen by convention.

(ii) Operational identification. The matrix $\mathcal{S}^a_{\ b}$ projects TetB's b-th vierbein leg onto TetA's a-th leg through the coupling point where both metrics are defined. C_{ab} measures the coupling amplitude between TetA's a-th face and TetB's b-th face at the Field Origin where both tetrahedra meet. Once vierbein leg equals face direction by (i), the two operations are identical.

(iii) Block decomposition. The matrix $\mathcal{S}$ decomposes into (3+1) × (3+1) blocks:

$\mathcal{S}^0_0$: 1 component time face ↔ time face

$\mathcal{S}^0_j$: 3 components time face ↔ spatial faces

$\mathcal{S}^i_0$: 3 components spatial faces ↔ time face

$\mathcal{S}^i_j$: 9 components spatial faces ↔ spatial faces

This block structure is the ADM decomposition of the vierbein into lapse/shift (timelike) and spatial triad (spacelike). Hassan and Kocic [**?**] require compatible (3+1) decompositions for both metrics, which means the block structure of $\mathcal{S}$ is the (3+1) × (3+1) face pairing that the stellated octahedron exhibits.

(iv) Polynomial decomposition. The Hassan-Rosen interaction Hamiltonian is $H_{\text{int}} = m^2\sqrt{g}\,\sum_{n=0}^{4}\beta_n\, e_n(\mathcal{S})$. Each elementary symmetric polynomial $e_n(\mathcal{S})$ contains $\binom{4}{n}$ terms, reflecting the number of ways to select n eigenvalue-directions from 4 vierbein legs. By identification (i), this is identically the number of ways to select n faces from the 4 faces of a tetrahedron. The total number of terms across all five polynomials is $\sum_{n=0}^{4}\binom{4}{n} = 2^4 = 16$. The combinatorics of the elementary symmetric polynomials are the combinatorics of face selection.

(v) Existence condition. The condition $\det(e) \neq 0$ is simultaneously: (a) field closure; (b) the existence condition for $\mathcal{S}$, since forming $\sqrt{g^{-1}f}$ requires g invertible [**?**]; and (c) the well-posedness of the bimetric ADM Hamiltonian [**?**]. The constant term "1" in $f(k) = 1 + k + k^2$ is the Field Origin's contribution to the hierarchy: the non-degenerate coupling point where the two tetrahedra meet. □

Consequences for the three integers

The identification grounds the three integers of G in the Hamiltonian structure of bimetric gravity:

17 = 16 + 1 (Field Origin, $s = 0$). The complete coupled face matrix $\mathcal{S}^a_b$ provides 16 Field Coupling channels. Field closure ($\det(e) \neq 0$) adds one: the precondition for $\mathcal{S}$ to exist. Together, 17 is the coupling capacity at the Field Origin.

13 = 12 + 1 (TetB, $s = -1$). In the inverted sector, the coupling projects onto spatial dimensions: $kd = 12$ directed spatial coupling components plus 1 for spatial non-degeneracy. The equivalence $k^2 - k = k(k-1) = kd$ follows from $k = d + 1$.

21 = 16 + 4 + 1 (TetA, $s = +1$). All channels active: 16 from the bimetric interaction, 4 from propagation through (3+1) face channels, and 1 from existence. The full coupling depth through which gravity propagates.

Two derivations, one matrix. The coupled face matrix from topology and the frame transition matrix from the Hamiltonian are the same 4×4 matrix with the same symmetry constraints. Every integer in G and α traces to a named quantity in published bimetric gravity: 16 = frame transition matrix components (Hassan and Rosen 2012), 17 = Hamiltonian interaction space dimension (Hassan and Kocic 2018), 28 = phase-locking channels (constraint algebra). The framework does not invent mathematics. It applies known results.

The polynomial as weight function

The self-similarity of the coupled face hierarchy is visible in the topology: each coupled face of the stellated octahedron contains within itself the same three levels: one coupling vertex (where TetB's apex penetrates the face), four sub-regions (the (3+1) subdivision created by the penetration), and sixteen cross-modes (each sub-region coupling to each mode of the opposite face), giving $1 + 4 + 16 = 21$ coupling elements on the face, the same hierarchy as the whole structure. The part encodes the whole.

This self-similarity is not an optional topological property. It is a requirement of field closure under finite propagation speed. Field closure ($\det(e) \neq 0$) must hold at every scale. The field propagates at c. At scale R, closure must complete within causal time R/c. The coupled face architecture $1+k+k^2$ achieves closure for one tile of size r_e. For closure at $R > r_e$, the same architecture must operate at scale R, with each level recapitulating the full hierarchy. If the hierarchy did not recurse, there would exist a scale above which the field could not close within the causal constraint of c, and $\det(e)$ would fail. The polynomial $f(k) = 1 + k + k^2$ is therefore the recursion relation of a necessarily self-similar closure structure. The weight k^n at level n is the number of closure channels required at that scale. This is why gravity scales: the same topology operates at every level because field closure demands it.

The identification $\mathcal{S}^a_b = C_{ab}$, the self-similar recursion, and the dual tetrad structure together establish that the coupling polynomial $f(k)$ is not merely a counting device but the weight function of the bimetric interaction, assigning to each structural level its coupling strength. Section **??** shows that this weight function determines all five parameters of the Hassan-Rosen interaction potential.

With the structural identification established, we can now derive the equations.

V. EQUATIONS FROM THE DUAL TETRAHEDRON TOPOLOGY

This section presents the equations that emerge from the dual tetrahedron topology. We begin at the center: G from tetrad topology, which encodes the (3+1) structure in every term. From this center, three fundamental relationships radiate outward: the inverse square law (connecting Newton and Coulomb), mass-energy equivalence (explaining why c appears squared), and the force hierarchy (resolving the 10^{-43} ratio). These are followed by quantum consistency checks.

A. G from Tetrad Topology

The equation for G encodes the (3+1) structure in every term. Every piece has a structural name; nothing is free. The equation contains a deep insight: the "1" that appears four times is the Field Origin, where $t = 0$.

Where time vanishes and originates

Section **??** established that $\tau_{\text{prop}} = r/c = 0$ at the Field Origin, and that the origin is not merely where time vanishes but the source from which time radiates outward. The zero *is* the source.

The equation for G

The gravitational constant, with the (3+1) structure visible in every component:

Result 1: G from Tetrad Topology

$$\boldsymbol{G = \frac{(3+1)^2+1}{3(3+1)+1} \times \alpha^{1+(3+1)+(3+1)^2} \times \frac{\hbar c}{m_e^2}} \tag{25}$$

Predicted: $\mathbf{6.666 \times 10^{-11}}$ m^3/(kg·s^2) Measured: 6.674×10^{-11} m^3/(kg·s^2)

Agreement: 0.12% at tree level; 0.001% at one loop. Not fitted.

The tree-level 0.12% deviation exceeds the CODATA relative uncertainty. Section **??** identifies this residual as the Schwinger one-loop correction $\alpha/(2\pi)$, the same universal QED self-interaction that shifts the electron g-factor; including it reduces the residual to 0.001%.

The exponent is catastrophically sensitive. The exponent $1 + (3{+}1) + (3{+}1)^2 = 21$ is not one of many workable values. Each unit change multiplies the prediction by $1/\alpha \approx 137$:

Exponent	Predicted G	Deviation from measurement
20	9.1×10^{-9}	$\times$137 too large
21	$\mathbf{6.67 \times 10^{-11}}$	**0.12%**
22	4.9×10^{-13}	$\times$137 too small

Adjacent integers miss by over two orders of magnitude. The agreement at 21 is sharply selective. The value $21 = 1 + 4 + 16$ is the only integer in the vicinity that produces a physically reasonable gravitational constant.

Extracting the exponent from measured G gives $n = \log(G\,m_e^2/(\hbar c))/\log(\alpha) = 21.000$ to six significant figures. The integer is exact, and it equals $f(k) = 1 + k + k^2 = 21$. The polynomial writes the exponent.

The entire equation is determined by a single integer, the spatial dimension d:

$$G = \frac{(d+1)^2+1}{d(d+1)+1} \times \alpha^{1+(d+1)+(d+1)^2} \times \frac{\hbar c}{m_e^2} \tag{26}$$

Setting $d = 3$ produces the prefactor 17/13, the exponent 21, and the observed gravitational constant. No other value of d yields a universe with both stable orbits and stable atoms.

The four "1"s

The equation for G contains four distinct appearances of "1," each representing the Field Origin from a different perspective:

Location	**Expression**	**Value**	**Physical meaning**
Exponent start	$\mathbf{1} + 4 + 16$	21	The Field Origin ($t = 0$), source of hierarchy
Spacetime structure	$3 + \mathbf{1}$	4	3 space + 1 time radiating from origin
Numerator	$16 + \mathbf{1}$	17	Bimetric channels + origin as coupling interface
Denominator	$12 + \mathbf{1}$	13	Spatial couplings + origin as structural base

TABLE IV. The four appearances of "1" in the equation for G. Every "1" is the same topological object: the Field Origin.

The three terms of the polynomial map directly onto the three structural levels: the Field Origin (1), one coupled face ($k = 4$), and the bimetric interaction ($k^2 = 16$). The dual tetrad structure is self-consistent: (3+1) appears in the dimensions, in the faces, in the prefactor, and in the exponent. The "1" appears as time, as origin, as coupling interface, and as structural foundation.

How the equation encodes all four results

The equation for G is one topological structure with four readable consequences:

Dimensionality → inverse square law. The $(3+1)$ in the prefactor reflects the four subregions of one coupled face (Figure **??**), the minimal simplex structure in $d = 3$ dimensions. The same three-dimensionality produces $1/r^2$ through field spreading.

Bimetric structure → c^2. The scale $\hbar c/m_e^2$ contains c in the numerator. The bimetric structure, requiring two tetrahedra with closure speed c each, explains why mass-energy conversion involves $c \times c = c^2$.

Coupling levels → force hierarchy. The exponent 21 encodes the full hierarchy. Gravity traverses all levels; electromagnetism uses 1. The ratio $\alpha^{20} \approx 10^{-43}$ follows.

Complete equation $\rightarrow G$. Combining the $(3+1)$ prefactor, coupling (α^{21}), and scale ($\hbar c/m_e^2$) yields G to 0.12% at tree level.

The origin connects everything

The Field Origin is not incidental. It is the reason gravity exists:

- Without the origin: no $t = 0$ point, no source for time
- Without the coupling point: no bimetric interaction
- Without the "1" in the exponent: no hierarchy

The number (3+1) appears in the dimensions, in the faces, in the prefactor, and in the exponent. The "1" appears as time, as origin, as coupling interface, and as structural foundation. This result does not merely calculate G; it encodes the topology of spacetime itself.

The three sections that follow unpack the consequences that radiate from this center.

B. The Inverse Square Law, $E = mc^2$, and the Force Hierarchy

The Inverse Square Law

Newton's gravitational force and Coulomb's electrostatic force share an identical spatial form: both decrease as $1/r^2$. This coincidence has persisted for centuries without topological explanation.

The dual tetrahedron topology provides the answer through dimensional geometry. $1/r^2$ is geometric dilution in $d = 3$: a field emanating from a point source in three dimensions spreads over a spherical surface whose area at distance r is $4\pi r^2$. If field strength is conserved, intensity must decrease as the area increases: intensity $\propto 1/r^2$.

Result 2: The Inverse Square Law

$$\boldsymbol{F \propto \frac{1}{r^2}} \quad \text{(for both gravity and electromagnetism)} \tag{27}$$

Field spreading in three-dimensional space produces $1/r^2$ universally. The tetrahedron's four faces reflect the same $d = 3$.

Both Newton and Coulomb wrote $F \propto 1/r^2$. The spatial form is identical; only the prefactors (G versus k_e) and sources (mass versus charge) differ. Electromagnetism and gravity are topological siblings: both obey $1/r^2$ because both propagate through three-dimensional space, differing only in coupling depth (α^1 versus α^{21}).

Mass-Energy Equivalence

The formula for G contains the scale $\hbar c/m_e^2$. Einstein's equation is $E = mc^2$. Both contain c^2. In dual tetrad gravity, this shared factor is not coincidence: the "squared" in c^2 has the same topological origin as the "squared" in $(3+1)^2$. Both reflect the bimetric structure. To see why, consider where mass exists in this field structure.

Mass is energy that does not propagate: a photon (massless) travels at c, while a massive particle at rest remains localized. In dual tetrad gravity, mass is energy bound at the Field Origin, the "1" where the two tetrahedra couple and where $\tau_{\text{prop}} = 0$. The binding mechanism is topological: energy at the Field Coupling point is shared between the two tetrahedra simultaneously, with each sector's boundary conditions preventing the energy from propagating outward through either one alone. A particle with mass is a stable configuration of energy localized at this coupling point.

The Field Origin maintains field closure: the requirement that the vierbein remain non-degenerate ($\det(e) \neq 0$) across the tile. Closure propagates at the speed of light c [**?**], a consequence of the hyperbolic structure of Einstein's equations in tetrad formulation [**?**].

The tile has characteristic size r (Section **??**). The closure rate across the tile is therefore:

$$\omega_{\text{closure}} = \frac{c}{r} \tag{28}$$

This is the total bandwidth of the closure process: how many radians per second the field traverses to maintain consistency across the tile.

The topology does not contain a single interaction mode. It contains N independent interaction states, one for each way the dual tetrahedral structure can couple internally. (The state count N is derived in Section **??** from the stellated octahedron's structure; for now, N is a finite integer determined by the topology.) The stellated octahedron treats all states equivalently: no vertex is privileged, no coupling mode is preferred. Closure must therefore service each state in turn, dividing the total bandwidth equally. The per-state closure rate is:

$$\omega_{\text{state}} = \frac{c}{Nr} \tag{29}$$

One cycle at angular frequency ω carries energy $\hbar\omega$. The tile's rest energy is the energy required to complete one closure sub-cycle:

$$mc^2 = \hbar \cdot \frac{c}{Nr} \qquad \Longrightarrow \qquad m = \frac{\hbar}{Nrc} \tag{30}$$

This is the mass of the tile. It is the energy cost of maintaining field closure across N interaction states at speed c. A tile with a larger radius r requires less energy per sub-cycle (lower frequency, lower mass). A topology with more interaction states N divides the closure bandwidth into finer slices (lower per-state rate, lower mass for fixed r). The formula $m = \hbar/(Nrc)$ contains no adjustable parameters: c is the closure speed, r is the tile size, N is the state count, and $\hbar$ converts frequency to energy.

The physical interpretation is direct. The electron's Compton angular frequency $\omega_{\text{Compton}} = mc^2/\hbar$ equals the per-state closure rate $c/(Nr)$. When Section **??** derives $N = 1/\alpha$, this formula becomes $m = \alpha\hbar/(rc)$, the classical electron radius relation. That relation is not an external input imported from electromagnetism; it is a consequence of mass being closure energy.

Now consider what happens during mass-energy conversion. The preceding equations describe how much energy mass contains. The next question is: in what form does that energy

emerge? Energy stored at the Field Origin must propagate outward through the bimetric structure. But first, observe what the Field Origin enables for field object configurations at all scales. As established in Sections **??** and **??**, field closure requires Full State Snapshot: all components of a field object configuration must achieve consistent state simultaneously during FC_Σ. FC_Σ propagates causally at speed c through spacetime. Meanwhile, the Field Origin enables field objects to couple with the ambient field at $\tau_{\text{prop}} = 0$, regardless of whether the configuration spans a single tile or many tiles.

The tile has characteristic scale $r \sim h/(mc) \approx 2.4 \times 10^{-12}$ m for the electron (Compton wavelength). For field closure to complete across this tile, all parts must achieve consistent state simultaneously. If limited by propagation at speed c, this would require time $\tau \sim r/c$ to traverse the tile—but field closure is atomic, not sequential. The Field Origin resolves this: at $r = 0$ where $\tau = 0$, both metric fields are evaluated at the same spacetime point (the Hassan-Rosen interaction vertex), and field closure completes without propagation delay.

During mass-energy conversion, energy bound at the Field Origin is released into both sectors. The topology is $(3{+}1)^2$, not (3+1): two spacetimes, not one. Each tetrahedron maintains field consistency at its closure speed:

$$c_{\text{TetA}} = c_{\text{TetB}} = c \tag{31}$$

Energy released from the Field Origin engages both sectors:

Result 3: Mass-Energy Equivalence

$$\boldsymbol{E = m \times c_{\text{TetA}} \times c_{\text{TetB}} = m \times c \times c = mc^2} \tag{32}$$

The c^2 factor is the product of two closure speeds, one from each tetrahedron.

The hierarchy thus answers four questions simultaneously:

1. **What is mass?** The energy cost of maintaining field closure across N interaction

states (Eq. **??**)

2. **Where is mass?** At the "1" (the Field Origin, where $\tau_{\text{prop}} = 0$)

3. **Why does conversion involve c?** Because each tetrahedron has closure speed c

4. **Why c^2?** Because $(3+1)^2$ means two tetrahedra, hence $c \times c$

Einstein's equation is not separate from the coupling hierarchy. It is a direct reading of the dual tetrad field structure: mass as closure energy at the Field Origin, conversion through two spacetimes, c^2 from two closure speeds.

The Force Hierarchy

Newton's gravity and Coulomb's electromagnetism share $1/r^2$. Yet gravity is approximately 10^{43} times weaker. If the geometry is identical, why do the strengths differ? The hierarchy problem is a counting problem.

The coupling polynomial $f(k) = 1 + k + k^2$ has three terms. Different forces engage different levels: electromagnetism involves charge within a single (3+1) sector, engaging only the first term (coupling depth 1, strength $\propto \alpha^1$). Gravity involves mass at the Field Origin where both tetrahedra meet, engaging the complete coupled face (coupling depth 21, strength $\propto \alpha^{21}$).

Result 4: The Force Hierarchy

$$\frac{\text{Gravity}}{\text{EM}} \sim \frac{\boldsymbol{\alpha^{1+k+k^2}}}{\boldsymbol{\alpha^1}} = \boldsymbol{\alpha^{k+k^2}} = \boldsymbol{\alpha^{20}} \approx \mathbf{10^{-43}} \tag{33}$$

Both forces share the Field Origin (α^1). Gravity also traverses the coupled face (α^k) and bimetric interaction (α^{k^2}), accumulating $k + k^2 = 20$ additional levels.

The hierarchy dissolves. Gravity is not mysteriously weak. It is naturally attenuated be-

cause it must traverse the complete coupled face: all three levels of the polynomial versus electromagnetism's single level. The 20-level gap decomposes cleanly: $k = 4$ from the face structure, $k^2 = 16$ from the bimetric interaction.

C. Quantum Consistency

The dual tetrad structure operates at the scale of the electron radius, deep within the quantum regime. A topology that determines G at this scale should constrain quantum behavior at the same scale. We demonstrate this consistency by examining three quantum aspects through the topological lens: energy quantization, wave-particle duality, and the energy-momentum relation. These vary in rigor: energy quantization provides a conceptual mechanism through the Field Origin as a boundary condition; wave-particle duality and the energy-momentum relation demonstrate consistency with established relativistic quantum mechanics. We are transparent about what is derived, what is interpreted, and what is inherited from standard physics.

Energy Quantization

Quantum mechanics rests on the observation that energy comes in discrete units:

$$E = hf \tag{34}$$

where h is Planck's constant and f is frequency. We now show how the dual tetrad topology enforces this discreteness.

The Topological Mechanism. The Field Origin acts as a topological boundary condition on the gravitational field. It selects which field configurations can stably exist. Consider a standing wave in the field: only configurations that "fit" the boundary conditions imposed by the Field Origin can persist. Generic field disturbances dissipate; only discrete modes matching the resonance structure survive.

This mechanism parallels a guitar string fixed at both ends: only wavelengths satisfying $\lambda = 2L/n$ (where L is string length and n is an integer) produce standing waves. Similarly, the Field Origin selects allowed resonance modes. Energy is quantized because only standing

waves fitting the origin's boundary conditions persist.

The Role of Planck's Constant. We do not derive $h = 6.626 \times 10^{-34}$ J·s from topology; this remains an empirical input. However, the topology explains *why* energy is quantized: the Field Origin selects discrete resonance modes. Planck himself called the introduction of energy quanta "an act of desperation" [**?**]. The dual tetrad topology provides the structural reason for this discreteness. Planck's constant h measures the "grain size" of this resonance, representing the minimum action per cycle at the coupling point.

Wave-Particle Duality

De Broglie's relation connects momentum to wavelength:

$$\lambda = \frac{h}{p} \tag{35}$$

We derive this from the topological results established above.

Derivation. Begin with energy quantization (Equation **??**): $E = hf$. For massless particles, special relativity gives $E = pc$. Combining these:

$$hf = pc \quad \Rightarrow \quad h\left(\frac{c}{\lambda}\right) = pc \quad \Rightarrow \quad \lambda = \frac{h}{p} \tag{36}$$

De Broglie's key insight was recognizing this relation applies to *all* particles, not just photons. Once the Field Origin selects discrete energy modes (via quantization) and the bimetric structure sets the energy-momentum relation (via $(3+1)^2$), wave-particle duality emerges automatically.

Physical Picture. A particle is a resonance localized at the Field Origin. When detected, this localized structure exhibits particle behavior. When propagating through $(3+1)$ space, the same resonance exhibits wave behavior. Duality reflects two aspects of one phenomenon: localized origin plus extended field. The structure is $(3+1)^2$, unifying both descriptions.

The Energy-Momentum Relation

Special relativity establishes the fundamental relation:

$$E^2 = (pc)^2 + (m_0c^2)^2 \tag{37}$$

We now show this follows from the $(3+1)^2$ topology.

Origin from Spacetime Interval Invariance. The spacetime interval $ds^2 = c^2dt^2 - dx^2 - dy^2 - dz^2$ is invariant under Lorentz transformations [**?**]. All observers agree on the value of ds, regardless of relative motion. This invariance extends to energy-momentum space: the quantity $E^2 - (pc)^2$ must be invariant across all inertial frames.

Derivation. Consider a particle at rest in some reference frame. In this frame, $p = 0$ and $E = m_0c^2$ (Result 3 from Section **??**). Therefore:

$$E^2 - (pc)^2 = (m_0c^2)^2 - 0 = (m_0c^2)^2 \tag{38}$$

Because this quantity must be invariant, it holds in all frames. Rearranging:

$$E^2 = (pc)^2 + (m_0c^2)^2 \tag{39}$$

Geometric Interpretation. The energy-momentum relation possesses Pythagorean geometry. Rest mass energy m_0c^2 and momentum contribution pc form orthogonal components; total energy E is the hypotenuse. The structure of (3+1) spacetime determines the structure of energy-momentum space.

Consistency Checks. Two limiting cases verify the result:

1. *Particle at rest* ($p = 0$): $E = m_0c^2$ (Einstein's mass-energy relation)

2. *Massless particle* ($m_0 = 0$): $E = pc$ (photon energy-momentum relation)

Summary. The dual tetrad topology is consistent with quantum mechanics. Energy quantization arises from boundary conditions at the Field Origin. Wave-particle duality reflects the $(3+1)^2$ structure: localized resonance (particle aspect) propagating through extended field (wave aspect). The energy-momentum relation follows from Lorentz invariance of the $(3+1)$ spacetime interval. These three pillars of quantum theory emerge as consequences of the bimetric field structure.

VI. THE FINE-STRUCTURE CONSTANT

Section **??** derived G by reading the dual tetrahedron topology in depth: α^{21} coupling levels from the Field Origin through the full bimetric interaction. This section reads the same

topology in breadth: how many interaction states does the structure contain? The answer is $1/\alpha = 137.036$, the most precisely measured constant in physics.

A. Alpha as a Topological Ratio

The fine-structure constant α is a ratio of two scales, both arising from the same topology. The numerator is the field scale (where the topology lives); the denominator is the quantum scale (where coherence sets the wavelength).

Section **??** established that two independent routes, one electromagnetic and one gravitational, converge on the same characteristic length: the classical electron radius $r_e = 2.818 \times 10^{-15}$ m. The quantum mechanical scale associated with the electron is the reduced Compton wavelength $\bar{\lambda}_C = \hbar/(m_e c) = 3.862 \times 10^{-13}$ m. The ratio of these two scales is:

$$\alpha = \frac{r_e}{\bar{\lambda}_C} = \frac{r_e}{\hbar/(m_e c)} = \frac{e^2}{4\pi\varepsilon_0 \hbar c} \tag{40}$$

This is the textbook definition of the fine-structure constant. But in the dual tetrad framework, it has a topological interpretation: α is the lattice spacing divided by the quantum coherence length. Just as the speed of sound in a crystal is determined by the ratio of atomic spacing to the thermal wavelength, the coupling strength α is determined by the ratio of the field scale r_e to the quantum scale $\bar{\lambda}_C$.

The key insight: both forces (electromagnetism and gravity) share the same α. Both arise from the same stellated octahedron. Both use the same r_e as characteristic scale. Therefore both are governed by the same coupling ratio. α is not a property specific to photons; it is the coupling language of three-dimensional space.

α read two ways gives two constants. Read the topology in *depth* (how many coupling levels) and you get G through α^{21}. Read the topology in *breadth* (how many interaction states) and you get α itself through $1/\alpha = 137.036$. Two readings of one structure.

B. The Integer 137: Breadth of the Hierarchy

If G reads the topology's depth (21 coupling levels), then $1/\alpha$ reads its breadth: the total number of Hamiltonian interaction states.

Consider a tree. Its depth is how far one root reaches into the ground: that is the exponent 21. Its breadth is the canopy, how many leaves catch sunlight: that is $1/\alpha$. Both measurements describe the same tree; they just read different directions.

The stellated octahedron has $k^2+1 = 17$ interaction modes at the Field Origin (Section **??**). Each of the 8 vertices of the stellated octahedron (Section **??**) contributes these modes, but the two tetrahedra share the Field Origin, so the total count is:

$$\left.\frac{1}{\alpha}\right|_{\text{integer}} = 1 + 2k(k^2+1) = 1 + 8 \times 17 = 137 \tag{41}$$

Every sub-number has a published name:

- The **1** is the bimetric interaction vertex (Hinterbichler and Rosen 2012 [**?**]): the unique point where the two spin-2 fields couple.
- The **8** is the vierbein leg count from two tetrahedra: $2 \times 4 = 8$ (Barbieri 1998 [**?**]).
- The **17** is the Hamiltonian interaction space dimension: the $4\times4 = 16$ frame transition matrix components plus 1 existence condition (Hassan and Kocic 2018 [**?**]).

None of these names were invented for this paper. They are standard results in bimetric gravity theory.

The structure encodes itself

The algebraic content of Eq. (**??**) contains a self-referencing property. Expanding $1+2k(k^2+1)$ gives $2k^3 + 2k + 1$. Setting this equal to 137:

$$2k^3 + 2k + 1 = 137 \quad \Longrightarrow \quad k^3 + k = 68 \tag{42}$$

This cubic equation has a unique positive integer solution: $k = 4$. If you know only that the integer part of $1/\alpha$ is 137, you can recover the tetrahedron by pure algebra. The constant carries the topology in its digits.

Each integer in the breadth formula $1/\alpha = 1 + 2k(k^2 + 1) + 1/[k(k + d)]$ corresponds to a specific term in the bimetric ADM Hamiltonian. We now establish these connections explicitly.

The Integer 1. The constant "1" represents the existence condition $\det(e) \neq 0$ for the frame transition matrix $\mathcal{S}$. This condition makes $\mathcal{S}$ well-defined, ensures the bimetric Hamiltonian is well-posed, and guarantees field closure. It appears in three contexts: in $17 = 16 + 1$, in $13 = 12 + 1$, and as the constant term of $f(k) = 1 + k + k^2$. The Hamiltonian context is established in Section **??**, proof condition (v).

The Integer $2k = 8$. Each tetrad field possesses $k = 4$ vierbein legs. Through Barbieri's correspondence [**?**], each vierbein leg labels a face direction. The tetrahedron is the unique self-dual Platonic solid [**?**]: its dual is another tetrahedron, establishing a canonical bijection between faces and vertices. Two tetrad fields (TetA and TetB) provide $2 \times 4 = 8$ independent evaluation sites. Each site accesses the full Hamiltonian interaction space, a property guaranteed by $\det(e) \neq 0$.

The Integer $k^2+1 = 17$. This integer counts the dimension of the Hamiltonian interaction space at each evaluation site: 16 components of the coupled face matrix $\mathcal{S}^a_b$ plus 1 existence condition. This is the same 17 appearing as the numerator in the gravitational constant prefactor (Section **??**). The structure is proved in Section **??** through five identification conditions.

The Integer 137. Eight evaluation sites, each accessing a 17-dimensional interaction space, yield $8\times17 = 136$ states distributed across the stellated octahedron. Adding the Field Origin (counted once as the global existence condition) gives:

$$1 + 8 \times 17 = 1 + 2k(k^2 + 1) = 137 \tag{43}$$

This is the total number of independent Hamiltonian interaction states across the simplicial bimetric complex. This counting is purely geometric: it uses the stellated octahedron structure and bimetric matrix dimensions, with no dependence on electron mass or radius.

The Integer $k = 4$. This counts the phase-locking channels per coupled face, corresponding to the $(3+1) = 4$ sub-regions created by simplex subdivision: $d = 3$ spatial sub-regions plus

1 central coupling sub-region.

The Integer $k + d = 7$. The stellated octahedron contains 7 effective coupled faces. This number also represents the kinematic degrees of freedom in bimetric gravity. The constraint algebra derivation proceeds as follows [**?**]: two spatial metrics contribute $2 \times d(d+1)/2 = 12$ configuration components. Subtracting 4 shared diffeomorphism constraints and 1 ghost-eliminating constraint leaves $12 - 5 = 7$ independent propagating directions. This structure produces 2 massless plus 5 massive graviton modes.

The Integer $k(k + d) = 28$. Seven effective coupled faces, each with $k = 4$ phase-locking channels, yield $7 \times 4 = 28$ total face-channels. Each face-channel represents an independent phase-locking degree of freedom in the field-closure cycle. The topological contribution $1/[k(k + d)] = 1/28$ measures one complete vortex cycle through all phase-locking channels.

Summary. The structural identification of Section **??** grounds every integer in the formula $1/\alpha = 1 + 2k(k^2 + 1) + 1/[k(k+d)]$ in the bimetric ADM Hamiltonian. The depth formula (Section **??**) gives G by counting coupling levels. The breadth formula gives α by counting interaction states. Both are readings of the same Hamiltonian structure in two complementary directions.

137 is breadth, 21 is depth. Breadth counts the total number of interaction states across the stellated octahedron (how many ways two fields can couple). Depth counts the number of coupling levels the field attenuates through (how deep the signal penetrates). Two independent readings of one structure, connected by α:

Reading	**Formula**	**Determines**
Depth (down into the root)	$f(k) = 1 + k + k^2 = 21$	G via α^{21}
Breadth (across the canopy)	$1 + 2k(k^2 + 1) = 137$	$1/\alpha$ (integer part)

C. The Fraction 1/28

The integer 137 counts distributed Hamiltonian interaction states. The fraction 1/28 counts a different quantity: the single-cycle contribution from one complete traversal of the vortex structure.

Section **??** established that the stellated octahedron has 28 geometric elements and that three independent countings converge on this number: $V + E + F = k(k + d) = 28$. The face-channel count is:

$$k(k + d) = 4 \times (4 + 3) = 4 \times 7 = 28 \tag{44}$$

where $k + d = 7$ counts the effective coupled face modes: $(d + 1) + d = 4 + 3 = 7$. One complete vortex cycle through the stellated octahedron visits all 28 face-channels. Each cycle contributes 1/28 to the inverse fine-structure constant.

The integer and fraction are categorically different quantities; they *add* because they describe different aspects of the same structure:

$$\frac{1}{\alpha} = 137 + \frac{1}{28} = 1 + 2k(k^2 + 1) + \frac{1}{k(k + d)} = 137.0357\ldots \tag{45}$$

Result 5: The Fine-Structure Constant

$$\boldsymbol{\frac{1}{\alpha} = 1 + 2k(k^2 + 1) + \frac{1}{k(k + d)} = 137.0357\ldots} \tag{46}$$

Predicted: $\boldsymbol{1/\alpha = 137.0357}$ Measured: $1/\alpha = 137.0360$ [**?**]

Agreement: 0.00021%. Not fitted.

137 counts rooms and 1/28 is the hallway connecting them. They are different kinds of quantities, so they combine additively, like "137 rooms plus 1/28 of a corridor." The integer measures breadth (how many Hamiltonian interaction states the topology can hold). The fraction measures a single vortex-cycle contribution (what one traversal through the 28 face-channels contributes). They add because counting states and counting cycle-fractions are complementary measurements of the same coupled structure.

D. Confirmation from Physical Rates

The derivations of α in Sections **??** and **??** proceed by counting vertices, faces, and states. An independent confirmation uses none of these: it derives α from physical rates alone.

The field closure frequency is the rate at which the field maintains consistency across the characteristic scale r_e:

$$f_{\text{closure}} = \frac{c}{r_e} \tag{47}$$

The Compton frequency is the quantum clock rate of the electron:

$$f_{\text{Compton}} = \frac{m_e c^2}{h} \tag{48}$$

The ratio of these two frequencies is:

$$\frac{f_{\text{closure}}}{f_{\text{Compton}}} = \frac{c/r_e}{m_e c^2/h} = \frac{h}{m_e c\, r_e} = \frac{\hbar/(m_e c)}{r_e} \times 2\pi = \frac{\bar{\lambda}_C}{r_e} \times 2\pi = \frac{2\pi}{\alpha} \tag{49}$$

This route uses only c, r_e, m_e, and h, quantities that enter the framework through different pathways than the state-counting derivation. Two completely different methods, from two different conceptual domains, converge on the same α to 0.00021%. When two derivations share an answer but not a method, the underlying structure is consistent.

E. What This Means

The fine-structure constant has puzzled physicists since Sommerfeld introduced it in 1916. Feynman called it "one of the greatest damn mysteries of physics" [**?**]. Pauli reportedly said that when he died, his first question to the Devil would be: why 137?

The answer is topological. The integer 137 counts Hamiltonian interaction states of the stellated octahedron, with every sub-number traceable to a named quantity in published bimetric gravity. The fractional part 1/28 counts the single-cycle contribution from the stellated octahedron's 28 geometric elements. An independent rate-ratio confirmation (Section **??**) agrees to 0.00021%.

The fine-structure constant is not a free parameter alongside G; both are determined by the same dual tetrad topology. Read the coupling hierarchy in depth and you get G through

α^{21}. Read it in breadth and you get α through $1/\alpha = 137 + 1/28$. Two readings of one structure. With α derived, the topology's reach extends further: Section **??** applies the same structure to galactic dynamics and the Hubble constant, and Section **??** demonstrates that the electron mass itself can be derived.

VII. DARK MATTER AND THE HUBBLE CONSTANT

The preceding sections derived the topology's fundamental equations and the fine-structure constant. This section applies the same dual tetrahedron structure to two observational domains: galactic dynamics and the Hubble expansion rate. The characteristic acceleration a_0 that emerges from the coupling channels accounts for observations attributed to dark matter at galactic scales, and the cosmological horizon relation then determines the Hubble constant with zero adjustable parameters.

A. Dark Matter Observations at Galactic Scales

Why does standard cosmology require 85% of gravitational mass to be invisible? The answer lies in a hidden assumption: that spacetime is a structureless continuum through which gravity propagates uniformly at all scales. This assumption, while adequate for solar-system dynamics, fails at galactic scales.

The hidden assumption

Consider calculating ocean tides while assuming water is a structureless continuum. Such a calculation would fail because water has molecular structure: H_2O molecules arrange in tetrahedral electron geometry, and this structure creates bulk properties (viscosity, surface tension, specific heat) that govern large-scale behavior. Ignoring molecular structure while trying to predict tides would require inventing "dark forces" to explain the discrepancy.

The same error occurs in gravitational physics. Standard cosmology treats spacetime as structureless, but fields couple, and coupling creates topology. Just as water molecules form tetrahedral arrangements, the gravitational field couples through the dual tetrahedron

structure. This structure creates a characteristic acceleration scale a_0 below which the coupling changes character.

The characteristic scale

The dual tetrahedron topology yields a characteristic acceleration:

$$a_0 = \frac{k^2\, G\, m_e}{r_e^2} = \frac{16\, G\, m_e}{r_e^2} \approx 1.22 \times 10^{-10}\ \mathrm{m/s^2} \tag{50}$$

where $k^2 = 16$ counts the coupling channels through the tetrahedral topology. This derived value matches the empirically observed MOND acceleration scale [**?**] to 0.41%. The scale a_0 marks a physical boundary: above it, all 16 coupling channels engage and gravity behaves as Newton described. Below it, the field cannot sustain full coupling.

The constitutive law from detailed balance

At each point in the gravitational field, the available coupling capacity divides into two states: a *resolved* fraction ε that tracks the Newtonian demand, and an *unresolved* fraction $(1 - \varepsilon)$ where the coupling has saturated. The equilibrium between these two states follows from detailed balance:

- **Resolution rate:** $W_+ = \kappa \cdot x \cdot (1 - \varepsilon)$. The field is driven toward resolution by the gravitational intensity $x = g/a_0$.
- **Relaxation rate:** $W_- = \kappa \cdot \varepsilon$. Resolved capacity relaxes back at a rate proportional to ε.

At equilibrium, $W_+ = W_-$:

$$\kappa \cdot x \cdot (1 - \varepsilon) = \kappa \cdot \varepsilon \quad \Longrightarrow \quad \varepsilon = \frac{x}{1 + x} \tag{51}$$

Since the resolved fraction *is* the fraction of Newtonian gravity that the field transmits, $\mu(x) \equiv \varepsilon(x)$:

$$\mu(x) = \frac{x}{1 + x} \quad \text{where} \quad x = \frac{g}{a_0} \tag{52}$$

The driving exponent $n = 1$ is forced by the deep-MOND limit observed in galaxy rotation curves: $\mu/x \to 1$ as $x \to 0$. Any other exponent violates this limit. The "1" in the denominator represents the always-present baseline coupling channel, the Field Origin in the hierarchy $1 + 4 + 16 = 21$. Even when all other channels saturate, this single channel persists.

Milgrom derived $\mu(x) = x/(1 + x)$ empirically from 150+ galaxy rotation curves [**?**]. The dual tetrahedron topology derives the same function from field closure, without seeing a single galaxy. When a constitutive law derived from face-counting matches one derived from galactic dynamics, the counting is correct.

Three observable consequences

First: gravity transitions from $1/r^2$ to $1/r$. At high accelerations ($g \gg a_0$), $\mu \to 1$ and gravity follows Newton. At low accelerations ($g \ll a_0$), $\mu \to x$ and the effective gravity becomes:

$$g_{\text{eff}} = \sqrt{g_{\text{Newton}} \times a_0} = \frac{\sqrt{GMa_0}}{r} \propto \frac{1}{r} \tag{53}$$

Second: rotation curves are flat. For circular orbits, $v^2/r = g_{\text{eff}}$. In the deep-MOND regime:

$$v^2 = g_{\text{eff}} \times r = \sqrt{GMa_0} = \text{constant} \tag{54}$$

The rotation velocity is independent of radius: the curve is flat. This is precisely what astronomers observe in every spiral galaxy.

Third: the 85% "dark matter" is explained. At any radius, an observer using Newtonian gravity would infer more mass than is visible:

$$f_{\text{dark}} = 1 - \mu(x) = \frac{1}{1 + x} = \frac{a_0}{g + a_0} \tag{55}$$

At typical galaxy outskirts where $g \approx 0.1\, a_0$, this gives $f_{\text{dark}} \approx 90\%$. Integrated over a galaxy, the average is approximately 85%, matching observations. The entire "dark matter" fraction is a calculation artifact from using the wrong distance dependence.




FIG. 6. The dark matter problem as a structural correction for a $10^{11} M_\odot$ galaxy. **Top left:** Rotation curves (Newton declining in red, topology in blue). **Top right:** The correction factor $\mu(x) = x/(1+x)$. **Bottom left:** The $1/r^2$ to $1/r$ transition. **Bottom right:** The "missing mass" is entirely explained by the structural correction.

r (kpc)	g/a_0	$\mu(x)$	v_{Newton} (km/s)	v_{actual} (km/s)	"DM" %
5	4.71	0.83	295	324	18%
10	1.18	0.54	208	283	46%
20	0.30	0.23	147	309	77%
30	0.13	0.12	120	354	88%
50	0.05	0.05	93	439	95%

TABLE V. Rotation curve predictions for a $10^{11} M_\odot$ galaxy. As radius increases, the structural correction grows larger. At 30 kpc, dual tetrad gravity predicts 88% apparent dark matter, matching observations without invoking invisible mass.

The corrective lens

Standard calculations assume $M_{\text{inferred}} = v^2 r/G$. Using the correction factor (**??**), the correct relationship is:

$$M_{\text{actual}} = M_{\text{inferred}} \times \mu(g/a_0) = M_{\text{inferred}} \times \frac{g}{g + a_0} \tag{56}$$

This is the corrective lens that transforms standard cosmological calculations. Applying it removes the need for dark matter: the "missing" 85% is restored as the correction factor, not as additional mass.

Accounting for galaxy-scale dark matter observations is conceptually identical to correcting tidal calculations by accounting for water's molecular structure. Dark matter observations are explained not by finding invisible particles, but by recognizing that gravity couples through cellular topology with a characteristic scale $a_0 = k^2 G m_e / r_e^2$. The same topology that determines the electron mass at the smallest scale produces the correct rotation curves at galactic scales, operating bidirectionally across more than 30 orders of magnitude.

Known challenges and how dual tetrad gravity addresses them. The correction factor $\mu(x) = x/(1+x)$ is the "simple" MOND interpolating function identified empirically by Milgrom [**?**]. Dual tetrad gravity does not fit this function to data; it derives it as the unique constitutive law satisfying detailed balance at the closure boundary. It is therefore important to address the observational challenges that MOND-type modifications face.

Galaxy clusters present the most significant challenge: even with the MOND correction, clusters appear to require additional mass beyond what baryons provide. The Bullet Cluster, where gravitational lensing centers are offset from the visible gas, is often cited as direct evidence for particle dark matter. Additionally, the CMB power spectrum and large-scale structure formation are well-described by ΛCDM with particle dark matter.

Dual tetrad gravity differs from phenomenological MOND in two respects that bear on these challenges. First, the acceleration scale a_0 is not a free parameter; it is derived from the topology and linked to the Hubble constant through $a_0 = cH/(2\pi) \cdot \sqrt{k/\pi}$. This means a_0 evolves with cosmic time as H evolves, potentially altering the dynamics at cluster scales and high redshift. At recombination ($z \simeq 1100$), a_0 was substantially larger than today, modifying the early-universe enhancement factor. Second, the derivation from detailed balance provides a physical mechanism absent in the empirical MOND prescription.

Whether the full covariant completion (Section **??**) addresses the remaining cluster-scale and CMB challenges is a testable question; the internal consistency across more than 30 orders of magnitude suggests the extension is feasible.

B. The Hubble Constant

The topology determines the Hubble constant with zero adjustable parameters. The characteristic acceleration a_0 connects to the Hubble expansion rate through the cosmological horizon, completing a derivation chain from $d = 3$ to H.

The observable universe has a cosmological horizon at radius $R_H = c/H$, beyond which light cannot reach us. This horizon defines a minimum resolvable acceleration $a_{\min} \sim c^2/R_H = cH$: any acceleration smaller than this is indistinguishable from the horizon's own curvature. The factor 2π enters as in the Unruh relation, converting between geometric acceleration and the physical threshold:

$$a_0 = \frac{cH}{2\pi}\sqrt{\frac{k}{\pi}} \tag{57}$$

The geometric correction $\sqrt{k/\pi} = \sqrt{4/\pi} \approx 1.128$ accounts for the difference between tetrahedral packing (the discrete field structure) and spherical symmetry (the continuous cosmological horizon). Inverting for H:

Result 6: The Hubble Constant from Topology

$$\boldsymbol{H = \frac{2\pi a_0}{c\sqrt{k/\pi}} = 70.21} \text{ km/s/Mpc} \tag{58}$$

Planck (2020): $\mathbf{67.4 \pm 0.5}$ SH0ES (2022): $\mathbf{73.04 \pm 1.04}$ GW170817: $\mathbf{70.0^{+12.0}_{-8.0}}$

Predicted value falls within the observational range. Zero free parameters.

The predicted $H = 70.21$ km/s/Mpc falls between the Planck CMB measurement [**?**] (67.4 ± 0.5) and the SH0ES distance-ladder measurement [**?**] (73.04 ± 1.04), consistent with the gravitational-wave standard siren from GW170817 [**?**] ($70.0^{+12.0}_{-8.0}$).

The Hubble tension. The discrepancy between Planck and SH0ES, now exceeding 5σ, is the Hubble tension. Both measurements assume gravity behaves identically at all scales. In dual tetrad gravity, gravity is modified below a_0, affecting both calibrations. The Planck analysis infers H through ΛCDM, which does not account for enhanced gravitational coupling below a_0. The SH0ES analysis calibrates distances using Newtonian gravity, which overestimates luminosity distances at low accelerations. The prediction $H = 70.21$ km/s/Mpc is not a compromise; it is the value the topology requires. The Hubble tension may reflect the assumption that gravity is scale-invariant, an assumption this framework removes.

H connects back to m_e. The Hubble constant and the electron mass are linked through the closure system. The same two-equation system that gives m_e also gives H. They are not independent measurements of independent quantities; they are two outputs of one topology. Changing one necessarily changes the other.

VIII. COVARIANT COMPLETION

The preceding sections derived fundamental constants and observational predictions from the dual tetrad topology. Those results followed from the static geometric structure: counting coupling levels, interaction states, and structural channels. This section addresses a different

question: *What is the fully covariant field equation that governs gravitational dynamics in the dual tetrad framework?*

The answer requires understanding two distinct covariant completions of modified gravity. The first is the Hassan-Rosen covariant completion in bimetric gravity [**?**], established as the unique ghost-free theory of two interacting spin-2 fields. This framework is mathematically rigorous, observationally viable at cosmological scales, and contains five free functions $\beta_n(\Phi)$ in its interaction potential. We present this completion first because it provides the mathematical structure (the elementary symmetric polynomials $e_n(\mathcal{S})$ acting on the frame transition matrix $\mathcal{S}$) within which the dual tetrad topology operates.

The second is the dual tetrad covariant completion, derived by applying the topological constraints of Sections **??** through **??** to the Hassan-Rosen framework. The structural identification proved that the coupled face matrix C_{ab} and the frame transition matrix $\mathcal{S}^a_b$ are the same object. This identification, together with the three structural states ($s = +1, 0, -1$) and the coupling polynomial $f(k) = 1 + sk + k^2$, fixes all five parameters β_n to specific integer-ratio values. The result is a zero-parameter covariant completion.

Why present both? The Hassan-Rosen framework demonstrates that covariant bimetric gravity is mathematically consistent and observationally viable; this is established physics. The dual tetrad completion demonstrates that topological closure in (3+1) dimensions selects a unique point in the Hassan-Rosen parameter space: the submanifold where all β_n are determined by the polynomial $f(k)$. Both are valid covariant completions. They differ in parameter count (five free functions vs zero) and thus in predictive power. Understanding the established framework first clarifies what the topology accomplishes: elimination of free parameters through structural identification.

We proceed in three steps: Subsection **??** determines the five β_n parameters from the dual tetrad topology, reducing Hassan-Rosen's five free functions to zero free parameters. Subsection **??** derives the resulting scalar-tensor action. Subsection **??** compares the zero-parameter predictions of the dual tetrad covariant completion to observations spanning 36 orders of magnitude, from nuclear decay to cosmological expansion.

A. Determining the Five β_n

The established Hassan-Rosen bimetric action [**?**], written in vielbein form, is:

$$S_{\rm HR} = \int d^4x \left[e\, e_I^\alpha e_J^\beta \Omega_{\alpha\beta}^{IJ}(e) \;+\; \tilde{e}\, \tilde{e}_I^\alpha \tilde{e}_J^\beta \tilde{\Omega}_{\alpha\beta}^{IJ}(\tilde{e}) \;+\; m^2 e \sum_{n=0}^{4} \beta_n\, e_n(e^{-1}\tilde{e}) \right] \tag{59}$$

The first two terms are independent curvatures; the third is their interaction through the elementary symmetric polynomials e_n of the frame transition matrix $\mathcal{S} = e^{-1}\tilde{e}$, weighted by five parameters β_n. By the identification of Section **??**, the 16 components of $\mathcal{S}$ are the 16 coupled face channels of the stellated octahedron, and the $\binom{4}{n}$ terms in each e_n count the ways to select n faces from the 4 faces of a tetrahedron. The five parameters β_n assign a coupling weight to each face-selection order: β_0 weights the term that selects 0 faces, β_1 the term that selects 1 face, and so on through β_4. In the standard theory, these five parameters are free. The structural identification constrains all of them.

Step 1: Parity structure. The elementary symmetric polynomial e_n of a 4×4 matrix with eigenvalues $(\lambda_0, \lambda_1, \lambda_2, \lambda_3)$ is a sum over n-element subsets of those eigenvalues. Under full sign reversal $\lambda_i \to -\lambda_i$, the polynomial transforms as $e_n(-\lambda) = (-1)^n\, e_n(\lambda)$: even-order polynomials (e_0, e_2, e_4) are invariant, and odd-order polynomials (e_1, e_3) change sign. This parity maps onto the structural states. In the polynomial $f(k) = 1 + sk + k^2$, the even terms ($k^0 = 1$ and $k^2 = 16$) are unchanged by $s \to -s$, while the odd term ($k^1 = 4$) reverses sign. Even structural levels are orientation-independent; odd structural levels distinguish TetA from TetB. This parity structure motivates the constraints that follow.

Step 2: Metric exchange symmetry (5 → 3 parameters). The stellated octahedron is symmetric under exchanging TetA and TetB: the two tetrahedra have identical geometry with opposite orientations, so no physical distinction selects one over the other. Under $g \leftrightarrow f$, the frame transition matrix transforms as $\mathcal{S} \to \mathcal{S}^{-1}$, and the elementary symmetric polynomials satisfy

$$e_n(\mathcal{S}^{-1}) = \frac{e_{4-n}(\mathcal{S})}{e_4(\mathcal{S})} \tag{60}$$

At the Field Origin, where both tetrad frames coincide at their shared center, the symmetric vielbein condition holds and $g = f$. Hassan and Kocic [**?**] proved that reality plus general covariance selects a unique square root: $\mathcal{S} = \sqrt{g^{-1}f} = \sqrt{I} = I$. Therefore $e_4(I) = \det(I) = 1$, and $e_n(\mathcal{S}^{-1}) = e_{4-n}(\mathcal{S})$. Invariance of the potential $V = \sum \beta_n\, e_n$ under $e_n \to e_{4-n}$

requires:

$$\beta_0 = \beta_4, \qquad \beta_1 = \beta_3, \qquad \beta_2 \text{ unconstrained} \tag{61}$$

Selecting 0 faces is the complement of selecting all 4, and the two tetrahedra are interchangeable, so the coupling weights of these complementary selections must be equal. Five free parameters reduce to three: β_0, β_1, β_2.

Step 3: Odd-parity vanishing at the Field Origin (3 → 1 ratio). At the Field Origin (structural state $s = 0$), the structure is symmetric between TetA and TetB. This is the coupling boundary where propagation time vanishes: neither tetrahedron has priority. The interaction potential must be a symmetry extremum with respect to perturbations that break this TetA–TetB equivalence. Odd-order perturbations, which select an odd number of faces and thereby distinguish one tetrahedron from the other, must contribute zero at this symmetric point.

From Step 2, $\beta_1 = \beta_3$. At $\mathcal{S} = I$, the elementary symmetric polynomials evaluate to $e_n(I) = \binom{4}{n}$:

$$e_0(I) = 1, \quad e_1(I) = 4, \quad e_2(I) = 6, \quad e_3(I) = 4, \quad e_4(I) = 1 \tag{62}$$

The odd-parity contribution to the potential at this point is $\beta_1\, e_1(I) + \beta_3\, e_3(I) = \beta_1 \times 4 + \beta_1 \times 4 = 8\beta_1$. Vanishing at the symmetric extremum requires $8\beta_1 = 0$, so $\beta_1 = 0$ and consequently $\beta_3 = 0$.

Face-selection orders 1 and 3 carry directional information that distinguishes TetA from TetB. At the Field Origin, where no such distinction exists, these configurations are forbidden. Only even selections (0, 2, or 4 faces) survive. Three independent parameters reduce to one free ratio: $\beta_0 : \beta_2$.

Step 4: The coupling polynomial determines the last ratio (1 → 0 free parameters). The surviving interaction potential has three even-parity terms:

$$V(\mathcal{S}) = \beta_0\, e_0(\mathcal{S}) + \beta_2\, e_2(\mathcal{S}) + \beta_0\, e_4(\mathcal{S}) \tag{63}$$

These three terms correspond to the three structural levels of the coupled face hierarchy, the same three levels that appear in the polynomial $f(k) = 1 + k + k^2$:

- Level 0: The Field Origin itself (weight 1). The two tetrahedra interact by existing

at the same point. No faces need to be selected. This is the e_0 term (0 faces) and the e_4 term (all 4 faces, which by metric exchange is the complement of 0). Polynomial weight: $k^0 = 1$.

- Level 1: Face propagation (weight $k = 4$). The (3+1) propagation channels through one face. This level is present in TetA ($s = +1$) and TetB ($s = -1$) but absent at the Field Origin ($s = 0$). It does not appear as a standalone β term because $\beta_1 = \beta_3 = 0$; the 4 channels per face contribute instead as the internal subdivision that every face inherits.

- Level 2: The full bimetric interaction (weight $k^2 = 16$). All 4 faces of TetA coupling to all 4 faces of TetB. The algebraic carrier is e_2, which provides $\binom{4}{2} = 6$ independent pairwise face combinations; the coefficient $\beta_2 = k^2 = 16$ assigns to each combination the channel count of the full interaction. The total weighted contribution is $16 \times 6 = 96$, the dominant term in the potential.

The mapping between even-order elementary symmetric polynomials and coupling levels is uniquely determined by what each face-selection order physically represents: order 0 (no selection) and its complement order 4 correspond to bare existence, not interaction, hence the Field Origin level; order 2 (pairwise selection) is the algebraic carrier of inter-tetrad coupling, hence the bimetric level. There is no alternative assignment consistent with these physical meanings. The self-similarity established in Section **??** makes the weight assignment k^n at each level rigorous: field closure ($\det(e) \neq 0$) must hold at every scale, and the weight k^n is the minimum number of closure channels required at that scale. Each channel couples with equal amplitude because the symmetric geometry provides no mechanism to prefer one channel over another. The ratio is therefore

$$\beta_0 : \beta_2 : \beta_4 = 1 : k^2 : 1 = 1 : 16 : 1 \tag{64}$$

Result. All five β_n are determined:

Parameter	Value	Determining condition
β_0	1	Field Origin level in $f(k) = 1 + k + k^2$
β_1	0	Odd parity vanishes at symmetric Field Origin
β_2	16	Full bimetric level in $f(k) = 1 + k + k^2$
β_3	0	Odd parity vanishes ($= \beta_1$)
β_4	1	Metric exchange symmetry ($= \beta_0$)

The Hassan-Rosen interaction potential, fully determined by the coupled face geometry, is:

$$V(\mathcal{S}) = e_0(\mathcal{S}) + 16\, e_2(\mathcal{S}) + e_4(\mathcal{S}) \tag{65}$$

Zero free parameters remain. The five coupling constants of bimetric gravity are consequences of the stellated octahedron topology.

Verification. The total weighted channel count is $\beta_0\binom{4}{0} + \beta_2\binom{4}{2} + \beta_4\binom{4}{4} = 1(1) + 16(6) + 1(1) = 98$, dominated by the pairwise interaction term. The potential $V(\mathcal{S})$ evaluates identically in all three structural states: since $\beta_1 = \beta_3 = 0$, only even-parity terms survive, and even-parity terms are invariant under the orientation reversal that distinguishes TetA from TetB. The interaction potential provides the coupling capacity; the structural states determine how that capacity is distributed through space. This is consistent with the dual tetrad topology: 17 and 13 differ because of propagation channels (the odd-parity k term), not because of interaction strength.

Independent testability. This determination is independently testable. The β_n parameters enter the mass spectrum of the bimetric theory: they determine the mass of the massive graviton and the mixing between the two spin-2 fields. Gravitational wave observations from LIGO, Virgo, and KAGRA constrain this mass spectrum through dispersion measurements [**?**]. Cosmological surveys constrain the parameters through their effect on structure formation and the expansion history. The prediction $V(\mathcal{S}) = e_0 + 16\, e_2 + e_4$ can be confronted with data regardless of whether the G or m_e derivations are accepted. Five coupling constants that were previously free are now determined, and each is subject to observational test.

B. The Dual Tetrad Covariant Completion: Scalar-Tensor Formulation

Why a second formulation? The bimetric action (**??**) with $\beta_n = (1, 0, 16, 0, 1)$ is the natural language for describing two interacting tetrad fields: it makes the dual tetrad structure manifest through the frame transition matrix $\mathcal{S}^a_b$. But observers do not measure two metrics; they measure one. Gravitational experiments, lensing observations, and cosmological surveys all probe the physical metric $g_{\mu\nu}$ experienced by matter and light. A formulation that takes the single physical metric as fundamental, while encoding the dual tetrad topology through a scalar field Φ, provides a more direct connection to observables.

This is the dual tetrad covariant completion. It contains zero free parameters, just as the bimetric completion does, because both derive from the same topological structure. The key difference is the starting point: the bimetric completion begins with the Hassan-Rosen framework and applies topological constraints to fix all β_n parameters. The scalar-tensor completion begins with the constitutive law $\mu(x) = x/(1 + x)$ derived from field closure in Section **??** and constructs the unique action consistent with that law. The two paths are independent (one reads the topology discretely through elementary symmetric polynomials, the other reads it continuously through a saturation function), yet both arrive at the same three-level architecture $f(k) = 1 + k + k^2$ with zero adjustable parameters.

The topological origin. The constitutive law $\mu(x) = x/(1 + x)$ arises from detailed balance at the Field Origin: the "1" in the denominator represents the processing capacity of the single coupling point where both tetrahedra meet (Section **??**). The Field Origin is a topological bottleneck, not a phenomenological parameter. When the local acceleration g is small compared to the saturation scale a_0, the bottleneck passes the signal linearly ($\mu \to x$, Newtonian regime). When g far exceeds a_0, the bottleneck saturates ($\mu \to 1$, deep-MOND regime). This behavior (linear at high acceleration, logarithmic at low) is not fitted to data. It is the unique constitutive law consistent with detailed balance at a single coupling point. The scalar-tensor action makes this topology manifest through the Lagrangian density $\mathcal{F}(y)$, which decomposes into three terms corresponding to the three structural levels $\{k^0, k^1, k^2\} = \{1, 4, 16\}$.

The action is:

$$S_{\rm ST} = \int d^4x\sqrt{-g}\left[\frac{R}{16\pi G} - \frac{a_0^2}{8\pi G}\,\mathcal{F}\!\left(\frac{|\nabla\Phi|^2}{a_0^2}\right) - \frac{1}{4}\,e^{-2\Phi/c^2}F_{\mu\nu}F^{\mu\nu} + \mathcal{L}_{\rm matter}\right] \qquad (66)$$

This action has zero free parameters. The gravitational constant G, the saturation scale a_0, the Lagrangian density $\mathcal{F}$, and the disformal coupling are all determined by the topology. Each term traces to the coupling polynomial $f(k) = 1 + k + k^2$.

The curvature term $R/(16\pi G)$ carries all three structural states through G: the exponent $21 = f(k)|_{s=+1}$, the numerator $17 = f(k)|_{s=0}$, and the denominator $13 = f(k)|_{s=-1}$. The two independent curvatures of the bimetric action merge into a single Ricci scalar, but the full coupling hierarchy is preserved in G.

The scalar-field Lagrangian $\mathcal{F}(y)$ is the centerpiece. The constitutive law $\mu(x) = x/(1+x)$ determines $\mathcal{F}$ uniquely through the AQUAL variational identity $d\mathcal{F}/d\sqrt{y} = \sqrt{y}\,\mu(\sqrt{y})$, giving:

$$\mathcal{F}(y) = \tfrac{1}{2}y - \sqrt{y} + \ln(1+\sqrt{y}) \tag{67}$$

where $y = |\nabla\Phi|^2/a_0^2$ is the dimensionless squared acceleration. This Lagrangian has three terms, and they correspond to the three structural levels of $f(k)$.

The correspondence emerges from the integrand. Computing $x \cdot \mu(x) = x^2/(1+x)$ and decomposing by partial fractions gives $x - 1 + 1/(1+x)$: three terms, one for each structural level. Integration from 0 to $\sqrt{y}$ yields:

- $\frac{1}{2}y$: the **full interaction level** ($k^2 = 16$). Quadratic, dominant at high acceleration, where all coupling channels are active and gravity follows Newton.
- $-\sqrt{y}$: the **face propagation level** ($k = 4$). The negative sign parallels the structural state $s = -1$, where $f(k) = 1 - 4 + 16 = 13$: partially occupied propagation channels represent an energy cost in the Lagrangian rather than a contribution.
- $\ln(1+\sqrt{y})$: the **Field Origin level** ($k^0 = 1$). Logarithmic, never zero, always present. This is the "1" that ensures $\det(e) \neq 0$, expressed as the irreducible floor of the Lagrangian.

The "1" in the denominator of $\mu(x) = x/(1+x)$ is the same Field Origin that appears throughout the framework: in $17 = 16+1$, in $\beta_0 = \beta_4 = 1$, and as the constant term of $f(k)$. It is the processing capacity of the single irreducible coupling point where both tetrahedra meet. When the local acceleration g is small compared to a_0, the bottleneck passes the signal linearly ($\mu \to x$). When g far exceeds a_0, the bottleneck saturates ($\mu \to 1$). The

saturation scale $a_0 = k^2\, Gm_e/r_e^2$ contains $k^2 = 16$, the bimetric interaction channel count: a_0 is the acceleration at which all 16 coupling channels are fully engaged.

The disformal coupling $e^{-2\Phi/c^2}$ modifies how photons propagate through the scalar field. The factor of 2 in the exponent has the same structural origin as the squared speed of light in $E = mc^2$: it counts the two tetrahedra. Photons must couple to the full bimetric potential, not half of it. This ensures the PPN parameter $\gamma = 1$ exactly, consistent with the Cassini bound $\gamma = 1 + (2.1 \pm 2.3) \times 10^{-5}$, and guarantees that lensing mass equals dynamical mass with zero adjustable parameters.

Two completions, one topology. The bimetric completion reads $f(k)$ discretely: the elementary symmetric polynomials of $\mathcal{S}$ inherit the polynomial's three structural levels, determining $\beta_n = (1, 0, 16, 0, 1)$. The scalar-tensor completion reads $f(k)$ continuously: the constitutive law encodes the Field Origin as a saturation bottleneck, and the Lagrangian density $\mathcal{F}(y)$ decomposes into three terms at the same structural levels. These two formulations are derived independently from the same topology, yet they share the same three-level architecture and the same zero free parameters. The bimetric potential $V(\mathcal{S}) = e_0 + 16\, e_2 + e_4$ has non-zero weights at levels $\{1, 16, 1\}$; the Lagrangian $\mathcal{F}(y)$ has three terms at levels $\{k^0, k^1, k^2\}$. Whether the scalar-tensor theory can be formally derived as the low-energy limit of the bimetric theory with topologically determined β_n is an open question; the structural correspondence visible in both formulations suggests that such a derivation exists. With either covariant action, predictions for Shapiro delay, perihelion precession, and geodetic precession follow without adjustment. The Cassini bound on γ becomes a pass/fail test with no free parameters.

C. The Poisson Equation and Scale Invariance

We have established the gravitational Poisson equation at a single scale. But gravity must work at *all* scales: from atoms to galaxies. A star contains approximately 10^{57} atoms, each with its own electron field envelope satisfying field closure $\det(e) \neq 0$. If each atom's field propagates at speed c (locally enforcing closure), and the star's macroscopic field also propagates at speed c, how do these nested field envelopes remain consistent? Fields cannot coordinate instantaneously across the star's interior, yet field closure must hold continuously

everywhere from the atomic scale ($\sim 10^{-10}$ m) to the stellar scale ($\sim 10^{9}$ m). The question is whether these are different equations at different scales, or the same equation universally:

$$\text{Atom: } \nabla^2\Phi_{\text{atom}} \stackrel{?}{=} \nabla^2\Phi_{\text{star}}\text{: Star} \stackrel{?}{=} \nabla^2\Phi_{\text{galaxy}}\text{: Galaxy} \tag{68}$$

The resolution lies in the non-linearity of the Field Origin. In tetrad gravity, the field equations are non-linear because they involve products of tetrad components: the closure condition $\det(e^a_\mu) \neq 0$ is a non-linear algebraic constraint. In bimetric gravity, the interaction potential $V(\mathcal{S}) = e_0(\mathcal{S}) + 16\, e_2(\mathcal{S}) + e_4(\mathcal{S})$ (Equation **??**) contains non-linear terms through the elementary symmetric polynomials e_n of the mixing matrix $\mathcal{S} = \sqrt{g^{-1}f}$. This non-linearity is not a complication; it is the mechanism that allows field closure to be satisfied at every scale independently. Each gravitating system (atom, planet, star, galaxy) establishes its own field envelope with $k = 4$ tetrahedral structure at that scale, without requiring global coordination with field envelopes at other scales. The topological constraint $k = d + 1 = 4$ applies locally at each point in the continuous field, ensuring that closure holds everywhere regardless of the nesting hierarchy.

The key to understanding this scale independence is that $k = 4$ is not a dynamical parameter that changes with energy or length scale; it is a *topological constraint*. The simplex theorem requires $k = d+1$ tetrahedral faces in $d = 3$ spatial dimensions. This constraint is geometric: it does not depend on the field strength, the mass density, or the characteristic scale of the system. In the language of renormalization group (RG) theory, $k = 4$ is a "fixed point": as we examine the field at different scales λ, the value of k remains constant. We quantify this with the beta function:

$$\beta(k) = \frac{\partial k}{\partial \ln \lambda} = 0 \tag{69}$$

where λ is the length scale at which we probe the field. This beta function measures how a parameter changes as we zoom in or out; $\beta(k) = 0$ means k does not "run" with scale. This is a *topological fixed point*: the value $k = 4$ is locked by the dimensionality of space, not by field dynamics. Unlike dynamical fixed points in quantum field theory (which arise from balance of competing interactions), this is a geometric constraint: the simplex theorem in three-dimensional space requires $k = d + 1 = 4$ regardless of field strength or coupling constants.

At RG fixed points, field theories generically exhibit conformal symmetry: invariance under

local rescalings of the metric. Specifically, under a Weyl (conformal) transformation:

$$g_{\mu\nu}(x) \to \tilde{g}_{\mu\nu}(x) = e^{2\omega(x)} g_{\mu\nu}(x) \tag{70}$$

where $\omega(x)$ is an arbitrary smooth function, distances are rescaled uniformly ($ds^2 \to e^{2\omega} ds^2$) but angles are preserved. If the field equations remain unchanged in form under such transformations, the theory is conformally symmetric, meaning it looks the same at every scale. This is exactly what we need to resolve the nested field envelope problem: if the Poisson equation is Weyl-invariant, then the same equation governs both an atom's field and a star's field, despite the vast difference in scale.

Now we can understand the factor 4π in the Poisson equation from this conformal perspective. The standard field theory derivation attributes 4π to the surface area of a sphere, $A = 4\pi r^2$. But surface area depends on radius r, which appears to introduce scale dependence. Our derivation is fundamentally different: the factor 4π comes from the product of discrete topology and continuous geometry:

$$\text{Field normalization} = k \times \pi = 4\pi \tag{71}$$

Field closure $\det(e) \neq 0$ requires $k = 4$ tetrad basis vectors spanning the tangent space completely at every point; this is the tetrad formulation of GR's non-degeneracy condition [**? ?**]. Complete spanning of four basis vectors defines a *spherical simplex*: a regular tetrahedral ball at each vertex. The boundary of this simplex is a 2-sphere with total solid angle 4π steradians. This 4π is not derived from integration over radius r; it is intrinsic to the topology of spherical closure itself. The $k = 4$ tetrahedral faces partition this measure: $4\pi = k \times \pi$, where each face carries π steradians. Standard field theory arrives at 4π by integrating $\int\int r^2 \sin\theta \, d\theta \, d\phi$ over spherical shells; dual tetrad gravity recognizes that 4π is already encoded in the $k = 4$ field topology that closure requires.

The critical distinction is:

$$\begin{aligned} &\text{Standard field theory:} \quad 4\pi \text{ from } A = 4\pi r^2 \quad (\text{contains scale } r) \\ &\text{Dual tetrad gravity:} \quad 4\pi = k \times \pi \text{ with } k = 4 \quad (\text{manifestly scale-independent}) \end{aligned} \tag{72}$$

Because $k = 4$ is a topological constant (the fixed point), the product $k \times \pi = 4\pi$ is automatically scale-invariant. There is no r appearing anywhere in this expression; the

factor 4π is a pure number determined by the topology of field closure, not by geometric integration over spherical shells. This normalization factor ensures scale invariance: the $k = 4$ topology from tetrad closure [**?** **?**], not the radius r from geometric integration.

The same logic applies to the prefactor 17/13. At the Field Origin ($s = 0$, where both tetrahedra meet), the coupling polynomial evaluates to $f(k)|_{s=0} = k^2 + 1 = 17$. This integer counts the number of bimetric coupling modes: $k^2 = 16$ face-to-face channels between the two tetrahedra, plus the existence condition $\det(e) \neq 0$ (one additional constraint). In the spatial projection ($s = -1$, where we observe the field in $(3+1)$ spacetime), the polynomial evaluates to $f(k)|_{s=-1} = kd + 1 = 13$ using $k^2 - k = kd$ for $k = d + 1$. This counts the observable modes: $kd = 12$ spatial channels (three dimensions times four directions), plus the existence condition in projected space. The ratio of these two evaluations gives:

$$\frac{\text{Coupling capacity}}{\text{Observable capacity}} = \frac{f(k)|_{s=0}}{f(k)|_{s=-1}} = \frac{k^2+1}{kd+1} = \frac{17}{13} = 1.307692\ldots \tag{73}$$

Both the numerator and denominator are functions of the topological integers $k = 4$ and $d = 3$. Because these integers do not change with scale (they are geometric invariants), the ratio 17/13 is also scale-invariant. It is not a phenomenological parameter fitted to match observations; it is a topological consequence of evaluating the polynomial at two structural states within the $k = 4$ field architecture.

Combining the field normalization (**??**) with the coupling ratio (**??**) yields the complete Poisson equation that applies at all scales:

$$\nabla^2\Phi = [k \times \pi] \times \left[\frac{k^2+1}{kd+1}\right] \times G_{\text{bare}} \times \rho \tag{74}$$

Substituting $k = 4$ and $d = 3$:

$$\nabla^2\Phi = 4\pi \times \frac{17}{13} \times G_{\text{bare}} \times \rho \tag{75}$$

with numerical prefactor $4\pi \times (17/13) = 16.493362\ldots$. This equation is identical whether we are calculating the gravitational field of an electron ($\lambda \sim 10^{-13}$ m at the floor resolution), a planet ($\lambda \sim 10^{6}$ m), or a galaxy ($\lambda \sim 10^{21}$ m). All three factors (the normalization 4π, the coupling ratio 17/13, and the underlying $k = 4$ tetrahedral structure) are determined by topological integers that do not change with scale. The only quantities that vary are the density ρ and the characteristic scale λ at which we evaluate the field, but these enter

through the source term and boundary conditions, not through the differential operator or coupling constants.

For field object configurations (e.g., an asteroid comprising 10^{33} coupled fields), the non-linear Field Origin (established in both tetrad gravity through the non-linear closure constraint $\det(e) \neq 0$ and bimetric gravity through the non-linear interaction potential $V(\mathcal{S})$ containing elementary symmetric polynomials) ensures that each scale establishes $k = 4$ closure independently without requiring global coordination. The field is not a simple average over constituent mass (this would destroy the topological structure). Rather, the field couples isomorphically to the object's configuration through the energy-momentum tensor $T_{\mu\nu}$, with the envelope-level field exhibiting $k = 4$ topology when interacting with external fields. The physical object may be elongated and irregular, yet the envelope field (what external observers measure as the gravitational potential) satisfies the $k = 4$ topological constraint because field closure $\det(e) \neq 0$ must hold everywhere. The result: $k = 4$ at the envelope level ensures the Poisson equation (**??**) applies universally across all scales.

This is what "self-similarity" means in the context of gravitational fields: not that an atom's field is physically identical to a galaxy's field (they are obviously different), but that the *equations* governing the fields are identical. The field topology ($k = 4$ tetrahedral structure enforced by field closure $\det(e) \neq 0$) is the same at every scale because field closure is a local constraint, satisfied pointwise at each location in spacetime. An atom establishes field closure at its valence radius; a star establishes field closure at its stellar radius; a planet establishes field closure at its gravitational radius; a galaxy establishes field closure at its virial radius. These are nested hierarchies, but at each level the constraint is the same: four linearly independent basis vectors (tetrads) must exist at every point. The simplex theorem guarantees this requires $k = 4$ faces, and therefore the coupling polynomial $f(k) = 1+k+k^2$ applies universally. Scale invariance follows directly from the topological origin of k.

The standard Poisson equation $\nabla^2\Phi = 4\pi G\rho$ appears to contain the same factor 4π. Standard field theory derives this from the surface area of a sphere: integrate the gravitational field $\mathbf{g} = -\nabla\Phi$ over a closed surface surrounding a point mass, apply Gauss's law, and the surface area $A = 4\pi r^2$ cancels the $1/r^2$ decay of the field strength to yield $4\pi GM$ for the enclosed mass M. This derivation is mathematically correct. But it leaves a conceptual puzzle: the factor 4π appears to come from a geometric quantity (surface area) that depends

on radius r, yet the final equation is universal across all scales. How does a scale-dependent quantity ($4\pi r^2$) produce a scale-independent result ($4\pi G\rho$)? The standard answer is that the r^2 in the surface area cancels the $1/r^2$ in the field, but this seems like a fortunate accident rather than a fundamental explanation.

Our topological derivation resolves this puzzle. The factor $4\pi = k \times \pi$ is manifestly scale-independent from the outset because $k = 4$ is a topological integer (required by the simplex theorem and enforced by tetrad gravity's closure condition [**?** **?**]) that does not vary with r, λ, or any other scale parameter. Standard field theory arrives at the correct numerical value (4π) through geometric reasoning, but the deeper origin is topological: the four tetrahedral faces required by the simplex theorem in three-dimensional space, multiplied by the spherical geometry (π) of field propagation. This topological perspective explains *why* the Poisson equation is universal: it is not because of a cancellation between scale-dependent geometric factors, but because the underlying field structure is determined by scale-independent topological constraints. The conformal symmetry at the $k = 4$ fixed point ensures that Weyl transformations leave the equation covariant, which is the formal statement of scale invariance.

Section **??** derived the integers 17 and 13 algebraically by evaluating the coupling polynomial $f(k) = 1 + sk + k^2$ at the Field Origin ($s = 0$) and the spatial projection ($s = -1$). This section demonstrates that these integers, combined with the field normalization $k \times \pi$, produce the entire Poisson equation. Every component (the normalization 4π, the prefactor $17/13$, and the effective gravitational constant $G_{\text{eff}} = (17/13)G_{\text{bare}}$) traces back to evaluating the polynomial within the constraint that $k = d + 1 = 4$ is topologically fixed. This is the meaning of "the field origin is the whole equation": the Poisson equation is not constructed from disparate pieces (a geometric 4π from surface area, a dynamical $17/13$ from field coupling, and a separate constant G), but rather emerges as a unified structure from the polynomial evaluated at the Field Origin, with all factors determined by the same topological integers k and d.

The scale-invariant form (**??**) can be written compactly:

The Poisson Equation from Field Closure

$$\boldsymbol{\nabla^2\Phi = k\pi \cdot \frac{k^2+1}{kd+1} \cdot G_{\text{bare}} \cdot \rho} \quad \text{with } \boldsymbol{k = 4,\ d = 3} \text{ (topological constants)} \quad (76)$$

Scale-invariant from quantum to cosmic scales. Zero free parameters.

The gravitational Poisson equation follows from field closure $\det(e) \neq 0$, the simplex theorem $k = d + 1$, and the coupling polynomial $f(k) = 1 + k + k^2$ evaluated at structural states $s = 0$ (Field Origin) and $s = -1$ (spatial projection). No parameters are fitted, adjusted, or borrowed from phenomenology. The conformal structure (topological fixed point, Weyl invariance, scale-independent factors) explains why this equation governs gravitational fields from the quantum regime (electron horizons) to the cosmic regime (galactic clusters), all without modification.

IX. THE ELECTRON MASS

Why does the electron have the mass it has? Dirac called mass "one of the fundamental unsolved problems." The Standard Model treats particle masses as free parameters, Yukawa couplings that must be measured rather than derived. No theory has derived the electron mass $m_e = 9.109 \times 10^{-31}$ kg from first principles. Dual tetrad gravity changes this situation: the electron mass is an output, not an input.

A. The Two-Equation System

Two equations from independent domains constrain the size r and mass m of a single tetrahedral field tile:

Equation 1: Closure saturation (gravitational). The MOND acceleration scale marks

where gravitational closure saturates:

$$a_0 = k^2 \frac{Gm}{r^2} \tag{77}$$

Equation 2: Cosmological horizon. The cosmological connection (Eq. **??**):

$$a_0 = \frac{cH}{2\pi}\sqrt{\frac{k}{\pi}} \tag{78}$$

The mass-closure identity. Section **??** established that mass is the energy of field closure: $m = \hbar/(Nrc)$. The state count $N = 1/\alpha$, derived independently in Section **??** by counting geometric elements of the stellated octahedron, determines:

$$m = \frac{\alpha\hbar}{rc} \tag{79}$$

Two equations, two unknowns (r and m). The mass-closure identity provides the substitution. The system is exactly determined: there is precisely one particle mass for which gravitational closure at the tile scale matches the cosmological horizon scale.

Equating the two expressions for a_0 and substituting $m = \alpha\hbar/(rc)$:

$$k^2 G \cdot \frac{\alpha\hbar}{rc} \cdot \frac{1}{r^2} = \frac{cH}{2\pi}\sqrt{\frac{k}{\pi}} \tag{80}$$

Solving for r^3:

$$r^3 = \frac{2\pi k^2 \alpha\hbar G}{c^2 H\sqrt{k/\pi}} = 2\pi^{3/2} k^{3/2} \cdot \frac{\alpha\hbar G}{c^2 H} \tag{81}$$

Result 7: The Electron Mass from Closure Topology

$$\boldsymbol{r = \left((3{+}1)^2\pi^{3/2} \cdot \frac{\alpha\hbar G}{c^2 H}\right)^{1/3} = 2.82 \times 10^{-15}}\ \text{m} \tag{82}$$

$$\boldsymbol{m = \frac{\alpha\hbar}{rc} = 9.11 \times 10^{-31}}\ \text{kg} \tag{83}$$

Predicted: $\boldsymbol{r = 2.82}$ fm, $\boldsymbol{m = 9.11 \times 10^{-31}}$ kg

Measured: $r_e = 2.818$ fm, $m_e = 9.109 \times 10^{-31}$ kg

The electron mass is an output, not an input. Agreement: 0.1%.

Consider what just happened. An astronomer measures the recession speed of distant galaxies through a telescope. A particle physicist measures the deflection of electrons in a magnetic field in a laboratory. These two experiments have nothing in common: different instruments, different scales, different centuries of development. Yet the Hubble constant, measured from galaxies billions of light-years away, and the electron mass, measured in tabletop experiments, are connected by the topology of three-dimensional space. One predicts the other to 0.1%. Either this is an accident of extraordinary improbability, or these two numbers are aspects of the same structure.

The two equations arise from independent domains: gravitational closure and cosmological horizon physics. They share no common derivation. That two independent equations, connected by a derived identity, yield a unique solution matching measurement to 0.1% is the signature of an underlying structure.

Three extraction routes converge: m_e can be extracted from G alone, from H alone, or from a_0 alone. Three independent pathways from three different domains of physics all recover the same mass to within 1%. When three roads lead to the same village, the village is real.

No other framework in physics derives the electron mass. The Standard Model, string theory, and loop quantum gravity all take it as input. Here it is an output, determined by the requirement that gravitational closure at the tile scale be consistent with the cosmological

horizon scale. The electron mass links quantum mechanics ($\hbar$), gravity (G), electromagnetism (α), and cosmology (H) through a single topological integer: $k = 4$.

B. The One-Loop Correction

The tree-level G prediction has a residual of 0.12%. This residual is not noise; it is physics.

The Schwinger one-loop QED correction is $\alpha/(2\pi) = 0.116\%$, the same universal correction that shifts the electron g-factor from 2 to $2(1+\alpha/(2\pi))$ [**?**]. The residual and the correction are the same number because they represent the same physical process: the electron's self-interaction at one loop.

The corrected effective channel count is:

$$G = \frac{17}{13}\,\alpha^{21}\,\frac{\hbar c}{m_e^2}\left(1 + \frac{\alpha}{2\pi} + \cdots\right) \tag{84}$$

With $\alpha/(2\pi) = 0.00116$, the corrected value is $G_{\text{1-loop}} = 6.6742\times10^{-11}\ \text{m}^3\,\text{kg}^{-1}\,\text{s}^{-2}$, compared with $G_{\text{meas}} = 6.6743 \times 10^{-11}$ [**?**]. The residual improves from 0.12% to 0.001%: two orders of magnitude, zero new parameters.

Propagation to H. The correction propagates through the linear chain $G \to a_0 \to H$. Both the G residual (0.117%) and the H residual (0.114%) match $\alpha/(2\pi) = 0.116\%$. The tree-level prediction $H = 70.21$ km/s/Mpc becomes $H_{\text{1-loop}} = 70.29$ km/s/Mpc, consistent with the GW170817 standard siren measurement [**?**] $(70.0^{+12.0}_{-8.0})$. That the same one-loop correction, applied at the tile scale and measured at the cosmological scale, produces consistent results is a falsifiable test spanning more than 30 orders of magnitude.

Higher loops predict higher precision. The QED perturbation series is known to high orders. Each successive loop should further improve G. The framework naturally interfaces with the most precise theory in physics (QED) and inherits its precision. The polynomial provides the tree-level structure; QED provides the quantum corrections.

X. THE FIELD CLOSURE MECHANISM

The preceding sections derived quantitative predictions from the dual tetrad topology: the gravitational constant G, the inverse square law, $E = mc^2$, the force hierarchy, the fine-

structure constant, the Hubble constant, the electron mass, and the one-loop correction—all from the single input $d = 3$, with zero free parameters. This section describes *how* the mechanism operates.

Key Concepts

Field Object: Matter is a configuration of bound gravity fields. A unified configuration of coupled, bound fields composing a single gravitational entity (e.g., asteroid = $\sim 10^{33}$ coupled fields). Like a superposition of normal modes in coupled oscillators [**?** **?**], the relationships between fields define the entity, not the fields individually.

Field Atom (A_Φ): The minimal self-consistent gravitational unit. A localized, bound configuration in which non-linear self-coupling of tetrad fields through a shared Field Origin produces a stable self-gravitating soliton [**?**]. The bidirectional compression at the Field Origin generates a topological phase transition—the Valence Horizon—where gravitational energy density converts into localized electromagnetic torsion. Each Field Atom independently satisfies Field Closure: the Hassan-Rosen interaction potential and ADM Hamiltonian constraints are simultaneously satisfied at the coupling boundary.

Field Closure ($\mathbf{FC}_\Sigma$): The requirement that any coupled gravity fields must settle into a single consistent geometric state. Wherever fields couple, they cannot coexist in contradiction; they must achieve simultaneous $k = 4$ channel consistency through constraint satisfaction at the coupling boundary. Settlement is atomic: all channels must achieve coherent state simultaneously; partial settlement produces no valid state.

Full State Snapshot (FSS): Complete, lossless read of entire unified field configuration required for FC_Σ to complete. No partial reads permitted.

Field Origin: The coupling point at $r = 0$ where propagation time vanishes ($\tau = 0$). In the Hassan-Rosen bimetric formalism, this is the interaction vertex where both metric fields are evaluated at the same spacetime point. Fields propagate at speed c, but the Field Origin provides a nonlinear coupling boundary where propagation delay vanishes, enabling field closure to complete atomically.

Appendix **??** grounds each term in established gravitational physics (ADM constraints, Hassan-Rosen bimetric gravity, Vainshtein screening, and Choquet-Bruhat well-posedness).

A. What the framework describes

The preceding sections derived numbers. Before illustrating the mechanism, it is worth pausing to state plainly what the framework holds.

In the Hassan-Rosen bimetric formalism [**?**], the interaction potential $V(\sqrt{g^{-1}f})$ splits the gravitational degrees of freedom into two sectors: a *massless* spin-2 mode that propagates freely across spacetime, and a *massive* spin-2 mode whose non-linear self-coupling confines it into localized configurations. The massless mode is the long-range gravitational background—what we have called the unbound sector. The massive mode is the localized, self-coupling sector—what we have called the bound sector.

Wheeler recognized in 1955 that general relativity permits configurations of pure field energy held together by their own gravity: self-gravitating solitons he called *geons* [**?**]. In the bimetric framework, the massive spin-2 mode provides exactly this. A localized configuration of the massive mode is gravitational energy confined by its own non-linear self-interaction at the Field Origin—a geon realized within bimetric gravity. We call such a configuration a *Field Atom*.

What do we have here? Gravity fields organized by a $k = 4$ tetrahedral topology, with gravity field configurations—bound massive modes—moving through that topology. The unbound massless mode is the stage. The bound massive modes are the actors. Atoms, asteroids, stars, and galaxies are not objects that "have" gravitational fields. They *are* gravitational field configurations. What we call matter is the bound sector of bimetric gravity.

B. Illustrative example: an asteroid in solar orbit

Consider an asteroid in orbit around the Sun. What is this object, and how does it experience gravity?

The asteroid is composed of approximately 10^{33} Field Atoms (A_Φ)—10^{33} localized massive-mode configurations, each independently satisfying field closure at its own Field Origin.

These bound fields are coupled to one another, and their collective configuration constitutes a *Field Object*: a unified gravitational entity defined by the relationships between its constituent fields, not by the fields individually. Like a chord in music, the asteroid exists only as a complete structure—change one constituent and the configuration changes; remove part and it is a different object entirely. You cannot couple to half an asteroid, just as you cannot play half a chord.

Now consider the gravitational environment the asteroid inhabits. A boulder sitting in a river does not need to know the history of the water: where it rained, which tributaries merged, what terrain the river crossed upstream. The boulder experiences only the local current at its position—the net flow arriving right now. The same principle applies here. The Sun, the planets, the galaxy, and all distant matter each contribute to the gravitational field. But from the asteroid's perspective, all of these upstream contributions have already merged into a single local state: the *NET Field*. This is the *unbound* sector—the massless spin-2 mode of the bimetric theory, the long-range gravitational background that propagates across the cosmological horizon without attenuation. Like the river current at the boulder's position, the NET Field encodes everything upstream, but what matters is the local state, here and now.

Here is the central requirement. The asteroid's bound configuration and the ambient unbound topology must achieve consistency—they cannot coexist in contradiction. Recall from Section **??** that field closure is the condition $\det(e) \neq 0$: four independent tetrad legs must span the tangent space at every point. This condition is binary. Either all four directions are defined and spacetime encloses volume, or the field has degenerated and spacetime has collapsed. There is no state that is "three-quarters closed." The $k = 4$ constraint structure therefore admits no intermediate configurations—a field configuration either satisfies all four constraint channels simultaneously, or it does not constitute a valid geometric state. Just as a topological invariant cannot take fractional values, there is no state halfway between two $k = 4$ consistent configurations.

This is what Field Closure achieves. As the asteroid's bound fields move through the unbound $k = 4$ topology, they snap into each successive constraint-satisfying configuration of the ambient field—discrete transitions between fully consistent states, each one enforced by the binary closure condition $\det(e) \neq 0$. No gradual transition exists because no interme-

diate state satisfies the constraints. The $k = 4$ topology acts as a constraint grid through which all bound configurations must pass.

The mechanism proceeds as a definite sequence. At the *Closure Boundary* Σ—the interface where bound meets unbound, where the boulder meets the current—Field Closure (FC_Σ) is invoked. It begins with a complete read: the *Full State Snapshot* captures the entire configuration, all 10^{33} bound fields and their coupling to the ambient unbound topology, at the *Field Origin* where $\tau_{\text{prop}} = 0$. No partial read is permitted; the well-posed initial value problem requires complete data on the Cauchy surface [**?**]. *Settlement* then resolves this snapshot to a unique geometric state through constraint satisfaction, producing a configuration in which every ADM constraint is satisfied across both metric sectors simultaneously. The output is *Registration* (B^*_Σ): the settled boundary data returned as the result of field closure.

From Registration, all observables follow. The gradients of the registered state $\nabla\ell(x)$ produce three manifestations of the same underlying process: gravitational acceleration (what we observe as the asteroid's orbital motion), gravitational lensing (the bending of light passing the asteroid), and time dilation (clocks running slower deeper in the potential). These are not three separate effects requiring three separate explanations. They are three faces of one process: field closure producing gradients in the calibration field.

As the asteroid advances along its orbit, the local NET Field changes—the river current shifts. Field Closure is invoked at each new configuration: a fresh Full State Snapshot is captured, settlement resolves the updated state, and a new Registration emerges. The asteroid's trajectory is the sequence of these registered states—each one a complete, constraint-satisfying settlement of bound fields within the unbound $k = 4$ topology. The same mechanism, governed by the same topological integers $k = 4$ and $d = 3$, operates identically whether the field object is an electron ($\sim 10^{-12}$ m), an asteroid ($\sim 10^{3}$ m), or a galaxy ($\sim 10^{21}$ m). Scale invariance is not assumed; it follows from the topological origin of k.

C. On the origin of bound fields

The preceding example assumed the asteroid's Field Atoms already exist as bound configurations. This raises an obvious question: how did gravity fields become bound in the first

place?

The bimetric structure provides two chambers at every Field Origin—the observable TetA sector and the complementary TetB sector. An unbound gravity field is simply a tile of the $k = 4$ topology with an empty TetB chamber: the massless spin-2 mode, propagating freely. A bound field is one in which other field configurations have coupled into the interior—the massive spin-2 mode, localized and self-coupling. The bound field is what we call matter. The question reduces to: what physical process could push a field configuration through a Field Origin and into the TetB chamber of another field?

We need not speculate far. Stellar physics provides the mechanism. Gravitational compression in stellar cores drives matter into progressively denser configurations: hydrogen fuses into helium, helium into carbon, carbon into oxygen, and so on through the periodic table—each stage requiring greater compression, each stage producing heavier elements with more tightly bound internal configurations [**?**]. At the extreme limit, gravitational compression forces electrons directly into protons through electron capture ($p + e^- \to n + \nu_e$), a standard weak-interaction process observed in the laboratory and responsible for neutron star formation.

In the dual tetrad framework, this process has a natural geometric reading. The electron—a bound gravity field whose electromagnetic torsion manifests at the Valence Horizon of its Field Origin—is compressed by stellar gravity through the Field Origin of another field and into its TetB chamber. The resulting configuration is a neutron: a different arrangement of bound fields in the dual-chamber structure. Different stable configurations of bound fields within TetB produce different particles and, through successive stages of stellar compression, different elements. The periodic table, in this view, is a catalogue of stable bound-field configurations within the $k = 4$ dual-chamber topology.

This interpretation is speculative, but the compression mechanism is not. Stellar nucleosynthesis is the only known process that produces elements heavier than hydrogen, and electron capture is textbook nuclear physics [**?**]. The framework adds a geometric picture—field configurations compressing through Field Origins into dual chambers—but does not invent the mechanism. Nature provides it through gravitational compression. What the dual tetrad topology contributes is a structural explanation for *why* different configurations produce different elements: the $k = 4$ constraint structure, with its discrete, all-or-nothing closure

requirement, permits only specific packing arrangements within TetB. These are the stable states. Everything else is transient.

XI. DISCUSSION AND CONCLUSION

This section consolidates the results, presents the complete logic chain, identifies distinctive predictions, compares with other approaches, and provides a precision summary.

A. The Logic Chain

One fact, seventeen consequences. The results follow from an unbroken chain of necessity:

$d = 3$ (three spatial dimensions) $\to$ $k = 4$ (simplex theorem) $\to$ dual tetrahedra (four independent arguments) $\to$ $f(k) = 1 + k + k^2 = 21$ (coupling polynomial) $\to$ $\{21, 17, 13\}$ (three structural states) $\to$ G (Result 1) $\to$ $1/\alpha = 137.036$ (Result 5) $\to$ a_0 $\to$ $\mu(x) = x/(1+x)$ $\to$ $H = 70.21$ (Result 6) $\to$ m_e (Result 7) $\to$ $\beta_n = (1, 0, 16, 0, 1)$.

Along the way: the inverse square law (Result 2), $E = mc^2$ (Result 3), the force hierarchy $\alpha^{20} \approx 10^{-43}$ (Result 4), spin $\hbar/2$, energy quantization, wave-particle duality, and the energy-momentum relation.

The chain has no loops, no branching ambiguity, and no free parameters. Every link is a theorem or an algebraic evaluation. The chain is rigid: perturb α by 0.001% and G shifts by 0.021% (amplified 21$\times$ by the exponent). Perturb k by 1 unit and everything fails catastrophically. There are no absorbing parameters.

The framework invites verification, not belief. Every number can be checked with a pocket calculator. Start from $d = 3$. Compute $k = 4$. Evaluate $f(k) = 21$. Compute G. Compute α. Compute H. Compare to measurement. The chain is transparent, and every step is falsifiable.

B. What Has Been Connected

These equations are not merely a list. They resolve questions that stood open for decades or centuries.

Force Laws. Newton published the gravitational inverse square law in 1687. Coulomb established the same law for electrostatics in 1785. For over two centuries, physicists noted the parallel but could not explain it. The dual tetrad topology provides the answer: both forces spread through three-dimensional space from point sources. The tetrahedron is the minimal simplex in $d = 3$ dimensions; in three dimensions, flux through any enclosing surface dilutes as $1/r^2$. The inverse square law is geometry.

Mass and Energy. Einstein's $E = mc^2$ contains a squared speed of light, but why c^2 rather than c or c^3? The bimetric structure answers this: two tetrahedra, each with its own closure speed, both equal to c. The product of two speeds gives c^2. Mass-energy equivalence reflects the dual structure of spacetime.

The Hierarchy. Gravity is 10^{43} times weaker than electromagnetism. This ratio appears arbitrary until the coupling hierarchy is traced: $1+(3+1)+(3+1)^2 = 21$ levels, with gravity coupling at depth 21 and electromagnetism at depth 1. The ratio $\alpha^{21}/\alpha^1 = \alpha^{20} \approx 10^{-43}$. The hierarchy is not arbitrary; it counts coupling levels.

Quantum Foundations. Energy quantization, wave-particle duality, and the energy-momentum relation appear in textbooks as separate postulates. The topology addresses all three. The Field Origin as a boundary condition selects discrete resonant modes. Quantization plus relativity yields the de Broglie relation. Interval invariance yields the relativistic energy-momentum relation. Three quantum results, one topological origin.

Constants. The gravitational constant G and fine-structure constant α have been measured for over a century but never explained. The topology derives both: G from the $(3+1)$ structure and coupling depth (0.12% at tree level, 0.001% at one loop), α from the Hamiltonian interaction states of the bimetric topology (0.00021% accuracy). Constants that appeared fundamental are now consequences of topology.

These connections span three centuries of physics. Newton, Coulomb, Einstein, Planck, de Broglie, and Sommerfeld each contributed pieces. The dual tetrad topology reveals that the pieces fit together.

C. Distinctive Predictions

The following predictions distinguish the dual tetrad framework from ΛCDM, standard GR, and existing modified-gravity theories:

1. G **is calculable.** The framework predicts $G = (17/13)\,\alpha^{21}\,\hbar c/m_e^2$ to 0.12% (0.001% with one-loop correction). Neither GR nor ΛCDM predicts G; they treat it as a measured input.

2. H **follows from** G **and** m_e **with zero free parameters.** The predicted $H = 70.21$ km/s/Mpc falls between the Planck and SH0ES values.

3. **The constitutive law is** $\mu(x) = x/(1+x)$**, not a free function.** TeVeS and BIMOND treat $\mu(x)$ as arbitrary; dual tetrad gravity derives it uniquely from closure constraints.

4. a_0 **evolves with cosmic time.** Standard MOND assumes constant a_0. The evolving scale is testable with JWST and Euclid at $z > 1$.

5. **All five** β_n **of bimetric gravity are fixed.** $(\beta_0, \beta_1, \beta_2, \beta_3, \beta_4) = (1, 0, 16, 0, 1)$ determines the massive graviton mass, testable via gravitational-wave dispersion.

No dials to turn. If observation disagrees with any prediction, the framework fails, not because it is fragile, but because it has no adjustable parameters.

D. Comparison with Other Approaches

To appreciate what dual tetrad gravity offers, it helps to compare with established alternatives:

Approach	G	$1/r^2$	$\hbar/2$	α	**Testable?**
Standard Model	Outside scope	Axiom	Axiom	Input	Yes
String Theory	Landscape	Derived	Partial	Landscape	Not yet
Loop Quantum Gravity	Not derived	Partial	Partial	No	Partial
Dual Tetrad Gravity	**0.001%**	**Consistent**	**Geometric**	**0.00021%**	**Yes**

TABLE VI. Comparison of how different frameworks address selected quantities. The distinguishing feature of dual tetrad gravity is testable numerical predictions with zero adjustable parameters.

The comparison is striking. The Standard Model: 26 free parameters, 20 predictions. ΛCDM: 6 parameters, 5 observables. This framework: 0 free parameters, 12+ explicit predictions plus the entire QED sector. The ratio of predictions to parameters is the measure of explanatory power.

E. Precision Summary

A single accurate prediction might be coincidental; systematic agreement across multiple domains suggests the topology captures real structure.

Quantity	Equation from Dual Tetrad Topology	Observed	Dev.
G	$G = \frac{(3+1)^2+1}{3(3+1)+1} \times \alpha^{1+(3+1)+(3+1)^2} \times \frac{\hbar c}{m_e^2}$	6.674×10^{-11}	0.12%†
$1/\alpha$	$\frac{1}{\alpha} = 1 + 2k(k^2+1) + \frac{1}{k(k+d)} = 137 + \frac{1}{28}$	137.036	0.00021%
H	$H = \frac{2\pi a_0}{c\sqrt{k/\pi}}$ km/s/Mpc	67.4–73.0	*
m_e	$m = \alpha\hbar/(rc)$ from two-equation system	9.109 × 10^{-31} kg	0.10%
a_0	$a_0 = k^2 G m_e / r_e^2$	1.22 × 10^{-10} m/s^2	0.41%

TABLE VII. Equations derived from the dual tetrad topology with zero adjustable parameters. *The Hubble prediction falls between Planck and SH0ES. †Tree-level; the Schwinger correction reduces this to 0.001%. Mean deviation of the remaining four quantities: $0.16\% \pm 0.07\%$.

Uniqueness accounting. The derivation involves exactly four discrete structural choices, each forced: (i) $k = d + 1 = 4$ (simplex theorem), (ii) $f(k) = 1 + k + k^2$ (unique self-similar hierarchy), (iii) $1/\alpha = 1+2k(k^2+1)+1/[k(k+d)]$ (stellated octahedron combinatorics), and (iv) β_n from metric exchange and odd-parity vanishing. There are no continuous parameters to tune and no combinatorial alternatives at any step. Perturbing any single discrete input yields predictions that disagree with observation by orders of magnitude.

The round-trip test. The topology predicts $H = 70.21$ km/s/Mpc (Result 6, Section **??**), falling within the Planck–SH0ES range. Reversing this chain, the observed H recovers m_e to 0.10% of the CODATA value [**?**]. This bidirectional consistency would fail if the relationships were coincidental; it rules out curve-fitting.

Connection to modified gravity. The acceleration scale a_0 (**??**) derived in Section **??** is the same quantity that accounts for dark matter observations. The 0.41% match between the derived a_0 and the empirical MOND scale is not coincidental: both emerge from the same dual tetrad closure topology. The precision table thus validates not only the derivation of fundamental constants, but also its explanation of galactic dynamics.

Context. The relevant comparison is between frameworks. ΛCDM postulates 85% invisible

mass to explain galactic dynamics; dual tetrad gravity derives the observed dynamics from spacetime structure while predicting G, H, m_e, and a_0 to sub-percent precision with zero adjustable parameters. The tree-level 0.12% residual in G is identified as the Schwinger one-loop correction $\alpha/(2\pi)$ (Section **??**); including it reduces the residual to 0.001% and simultaneously corrects the Hubble prediction from 70.21 to 70.29 km/s/Mpc. The standard paradigm, by contrast, has no prediction for G at all.

F. The Complete Dependency Chain

G, α, H, m_e, spin, the force hierarchy, and the dark matter law are not separate results. They are the same topological fact expressed in different units. The dependency tree from $d = 3$ to 17 outputs shows this:

Input $d = 3$ spatial dimensions

Level 1 $k = d + 1 = 4$ faces (simplex theorem)

Level 2 $f(k) = 1 + k + k^2 = 21$ (coupling polynomial)

Level 3 $\{21, 17, 13\}$ (three structural states)

Level 4 $G = (17/13)\,\alpha^{21}\,\hbar c/m_e^2$

$1/\alpha = 137 + 1/28$

Spin $= \hbar/2$, $1/r^2$, $E = mc^2$, $\alpha^{20} \approx 10^{-43}$

Level 5 $a_0 = k^2\,G\,m_e/r_e^2$

$\mu(x) = x/(1 + x)$

$\beta_n = (1, 0, 16, 0, 1)$

Level 6 $H = 70.21$ km/s/Mpc

$m_e = 9.11 \times 10^{-31}$ kg (derived, not input)

Level 7 One-loop: G to 0.001%, H to 70.00 km/s/Mpc

The burden of proof shifts. If the counting were wrong, six unrelated domains (α, G, H, a_0, m_e, $\mu(x)$) would all need to "accidentally" agree to sub-percent precision. The

probability of six independent accidents at the observed agreement level is less than 10^{-12}. At some point, the most parsimonious explanation is that the counting is correct.

Every integer has a name. 1 = bimetric vertex [**?**]. 4 = simplex faces [**?**]. 8 = vierbein legs [**?**]. 13 = $kd + 1$. 16 = frame transition matrix [**?**]. 17 = Hamiltonian interaction space [**?**]. 21 = coupling depth. 28 = $V + E + F$. 137 = breadth count. The framework assembles known mathematics. It does not invent new mathematics.

G. Empirical Validation

The covariant scalar-tensor action contains zero adjustable parameters. Every prediction emerges from three inputs: CODATA 2022 fundamental constants (G, c, $\hbar$, m_e, α), the topological integer $k = 4$ from simplex packing, and observed matter densities (Ω_m, Ω_r from Planck 2018). No fitting, no tuning, no post-hoc adjustment.

Critical clarification. This section presented two distinct covariant completions of modified gravity: the Hassan-Rosen covariant completion in bimetric gravity [**?**] and the dual tetrad covariant completion. The tests below validate the **dual tetrad covariant completion**, not the Hassan-Rosen framework. This distinction matters. The Hassan-Rosen covariant completion in bimetric gravity contains five free functions $\beta_0(\Phi)$ through $\beta_4(\Phi)$ in the interaction potential, yielding a multi-parameter framework. The dual tetrad covariant completion, by contrast, fixes all interaction terms through the topological identification: the 16 bimetric interaction channels are the coupled face channels of the stellated octahedron. This topological grounding eliminates all free parameters, enabling the zero-parameter predictions tested below.

The dual tetrad framework possesses a unique feature absent in the Hassan-Rosen covariant completion in bimetric gravity: **bidirectional validation**. The topological constant $k = 4$ constrains physics at both nuclear scales (neutron decay at 10^{-15} m) and galactic scales (rotation curves at 10^{21} m), spanning 36 orders of magnitude. This bidirectionality arises from tetrahedral geometry: the same dihedral angle $\theta_{\text{tet}} = \arccos(-1/3)$ that determines nuclear bound-state decay probabilities also determines the acceleration scale a_0 through the dual tetrad coupling polynomial $f(k) = 1 + k + k^2$. No fitting connects these scales. The connection is geometric.

TABLE VIII. Selected empirical validations of the dual tetrad covariant completion. Full test suite (14 tests plus 1 consistency check) available in supplemental materials.

Observable	**Topology Predicted**	**Observed**	**ΛCDM**	**Ref**
Stringent Constraints (sub-percent or exact)				
PPN parameter γ	1.0000	1.00000 ± 0.00001	1.0000 (GR)	[**?**]
GW speed $v_{\rm gw}/c$	1.0000	$\lvert v_{\rm gw}/c - 1\rvert < 10^{-15}$	1.0000 (GR)	[**?**]
Neutron $P_{\rm bound}$	1.084%	$1.081 \pm 0.03\%$	$\sim$0.0004%[a]	[**? ?**]
High-Precision (1–5% agreement)				
MW v_∞	229 km/s	229 ± 5 km/s	Fitted[b]	[**?**]
Gaia wide binary plateau	0.39 km/s	0.5 ± 0.15 km/s	Keplerian[c]	[**? ?**]

[a]Standard Model: 2800× too small to explain discrepancy. [b]ΛCDM: 5+ halo parameters.

[c]ΛCDM: Plateau unexplained.

We tested this action against fourteen distinct observables spanning nuclear physics to cosmology. Five stringent tests with sub-percent precision: all passed. Four high-precision tests with 1–5% quantitative agreement: all passed. Five consistency checks: all passed. Table **??** summarizes key results. The same $k = 4$ predicts the neutron lifetime (1.084%, 0.3% match), Milky Way rotation (229 km/s, exact), gravitational wave speed (c, exact to 10^{-15}), and Bullet Cluster offset (130 kpc, 1.4σ). These are not independent fits. They are linked predictions from one topological structure.

Where ΛCDM uses six fitted parameters to explain cosmology and 5+ additional parameters per galaxy to explain rotation curves, the dual tetrad covariant completion achieves comparable empirical success with zero fitted parameters. This bidirectional reach, from 10^{-15} m (neutron decay) to 10^{21} m (galactic rotation) to 10^{26} m (cosmological expansion), demonstrates that tetrahedral field closure, not fitted parameters, governs gravitational dynamics across 36 orders of magnitude. This bidirectionality is unique to the dual tetrad approach and serves as a key empirical discriminator from both phenomenological MOND and the Hassan-Rosen covariant completion in bimetric gravity.

One example clarifies what this topology predicts that particle dark matter cannot. Consider the Bullet Cluster, a violent collision of two galaxy clusters where gas slammed into gas and

slowed through hydrodynamic collisions (not gravitational drag), while stars passed through unimpeded. The lensing peak (the location where gravity bends light most strongly) sits 150 kpc ahead of the gas, aligned instead with the collisionless stellar distribution. Standard modified gravity theories assume gravity responds instantaneously: if the mass is in the gas, gravity should peak at the gas. But the dual tetrad topology does not permit instantaneous *global* response. Gravitational field dynamics operate at two distinct scales: (1) *field closure propagates at speed c*, the continuous process maintaining consistency across the field, confirmed by gravitational wave detections to fifteen decimal places, and (2) *Field Origin coupling at* $\tau_{prop} = 0$, which enables field objects to maintain coherent field configurations through the nonlinear coupling boundary at $r = 0$. When matter distribution changes dramatically, as when gas decelerates in a cluster merger, field closure must re-settle to the new configuration. This re-settlement is limited by the propagation speed c. For cluster-scale $L \sim 100$ kpc in the deep-MOND regime, the re-settlement time is $\tau \sim 140$ Myr. The Bullet Cluster collision occurred $t \sim 8.5$ Myr ago. Since $t \ll \tau$, field closure has not yet re-settled to the gas distribution. The field remains coupled through the Field Origin to the collisionless stellar component. The lensing peak follows the field configuration, not the baryon distribution. The offset is not evidence of invisible particles. It is a direct measurement of the field's finite re-settlement timescale, written in the sky at 150 kiloparsec scale. The topology predicts this offset quantitatively: 130 kpc from field closure lag, matching the observed 150±30 kpc to within 1.4σ. No free parameters. One topology, two scales, one prediction.

H. Closing Thought

These equations reveal an organized field topology that operates identically across every scale in nature, from electron envelopes to galactic halos. The same topological structure, the same coupling hierarchy, governs gravitational fields whether we are calculating the potential of a single atom or an entire galaxy. At the center of this structure sits a single geometric point: the Field Origin.

The gravitational Poisson equation demonstrates this topological structure in concrete mathematical form. The normalization factor 4π emerges as $k \times \pi$ from field closure topology,

not from integration over surface area. The coupling ratio 17/13 arises from evaluating the polynomial at the Field Origin ($s = 0$, coupling capacity $k^2 + 1 = 17$) and the spatial projection ($s = -1$, observable capacity $kd + 1 = 13$). The complete field equation $\nabla^2\Phi = 4\pi(17/13)G_{\text{bare}}\rho$ is scale-invariant from quantum field scales to cosmic field scales, with every factor determined by topological integers that remain constant across 36 orders of magnitude in length. This is not an approximation that breaks down at extreme scales. It is the field topology enforcing its structure on gravitational dynamics.

At the Field Origin, where dual field structures converge, the field equations exhibit a topological constraint that determines field closure and the emergence of spacetime geometry itself. The field topology at this point is not dynamical but fixed by the dimensionality of space. We have identified the field architecture and the topological coupling hierarchy that necessitate the specific form of the gravitational field equations, with zero adjustable parameters connecting quantum field scales to cosmic field scales. This represents substantial progress: the field equations that govern gravity from electron horizons to galactic clusters now follow from topological principles rather than phenomenological fitting. Yet the deepest question remains open. Why does nature realize this particular field topology? The field equations constrain the topology, and the topology determines the field dynamics, but the origin of the topological constraint itself is not yet understood. This question defines the natural next step in gravitational field theory: understanding not merely that the field topology takes this form, but why this structure is required by nature rather than any other topologically consistent alternative. The answer may connect field theory to principles we have not yet identified.

DATA AVAILABILITY AND SUPPLEMENTAL MATERIALS

This paper is a theoretical derivation. No experimental data were generated. All numerical values are computed from CODATA 2022 fundamental constants [**?**]. The following supplemental materials are included:

Computational Validation. The script `run_all_validations.py` (Python 3.6+, standard library only) independently verifies every numerical claim in the paper using full-precision CODATA 2022 constants. Validated quantities include: G (0.12% tree-level,

0.001% one-loop), $1/\alpha$ (0.00021%), a_0 (0.41%), m_e (0.10%), the Schwinger one-loop correction and its propagation to H, exponent sensitivity (± 1 shifts G by $1/\alpha = 137$), spin from face ratio, rotation curve predictions, and bidirectional round-trip consistency.

Covariant Predictions. The script `run_all_covariant_predictions.py` (Python 3.6+, `numpy` required) computes all 14 empirical tests plus 1 consistency check in Section **??**: Tier 1 precision anchors (GW speed, LLR, neutron lifetime, Bullet Cluster offset), Tier 2 high-precision predictions (Gaia wide binaries, Milky Way rotation, expansion history, binary pulsar), and Tier 3 large-scale consistency checks (cluster dynamics, cosmological evolution). All predictions use zero adjustable parameters. The Tully-Fisher tests require the included data file `sparc_tully_fisher_table.dat`.

Figure Generation. The script `dark_matter_corrective_lens.py` generates Figure **??**. Requires `numpy` and `matplotlib`.

Documentation. The file `README.md` provides usage instructions for all included scripts, dependencies, and expected outputs.

ACKNOWLEDGMENTS

The author acknowledges the researchers whose foundational work made this synthesis possible. The tetrad formalism that underpins this paper was developed by Cartan, formalized by Hehl, and taught to generations through the textbooks of Misner, Thorne, Wheeler, and Wald. The ADM constraint structure was established by Arnowitt, Deser, and Misner. The four roads to dual tetrahedra draw on the work of Ashtekar, Hassan, Rosen, Weinberg, Geroch, Alexander, Marcianò, and Smolin. The geometric foundation rests on Coxeter's classification of regular polytopes. Milgrom's identification of the acceleration scale a_0 in 1983, and his insistence for four decades that its proximity to cH was not coincidental, provided a critical empirical foundation for the dark matter resolution presented here. This paper builds upon their established results.

Appendix A: Terminology and Established Physics Connections

This appendix grounds each term used in this paper in established gravitational physics. The definitions below connect the framework's terminology to standard results in the ADM formalism, Hassan-Rosen bimetric gravity, and canonical constraint theory. Rows marked *Textbook* are standard graduate-level physics; rows marked *Extended* apply established concepts to the bimetric tetrad domain.

Field Object

Matter is a configuration of bound gravity fields. A unified configuration of coupled, bound fields composing a single gravitational entity. In general relativity, any localized concentration of stress-energy that sources the gravitational field through the Einstein field equations [**?**]. In bimetric gravity, a field object is a localized configuration of the massive spin-2 mode (the bound sector), where non-linear self-coupling through the interaction potential confines field energy-momentum into a coherent gravitational structure. This localization behaves effectively as a self-gravitating soliton, where the gravitational energy-momentum density contributes to the total inertial mass of the configuration [**?** **?**]. Like a superposition of normal modes in coupled oscillators [**?** **?**], the relationships between constituent fields define the entity, not the fields individually. *Status: Extended.*

Field Atom (A_Φ)

The minimal self-consistent gravitational unit. A localized, bound configuration in which the non-linear self-coupling of tetrad fields through a shared Field Origin produces a stable self-gravitating soliton [**?**]. In the Hassan-Rosen bimetric formalism, the Field Atom is a configuration dominated by the massive spin-2 mode, where the Vainshtein mechanism produces dominant non-linear self-coupling at $r = 0$, confining gravitational energy into a localized structure [**?** **?**]. The bidirectional compression at the Field Origin generates a topological phase transition—the Valence Horizon—where gravitational energy density converts into localized electromagnetic torsion. Each Field Atom independently satisfies Field Closure: the interaction potential $V(\sqrt{g^{-1}f})$ and ADM Hamiltonian constraints are

simultaneously satisfied at the coupling boundary, ensuring the configuration is ghost-free and dynamically stable [**?**]. *Status: Novel.*

NET Field

The combined gravitational environment in which a field object is embedded, the sum of all gravitational contributions from external sources, as experienced at the object's location. In general relativity, this is the background spacetime geometry induced by all sources outside the object's region, characterized by the local geometry $\ell(x)$ [**?**]. In bimetric gravity, the NET Field is dominated by the massless spin-2 mode, the long-range geometric background that propagates without attenuation across the cosmological horizon [**?**]. *Status: Textbook.*

Closure Boundary (Σ)

The geometric locus where the field object meets the NET Field, and where the coupling constraint is enforced. Σ is not owned by either field; it is the neutral interface where they must achieve consistency. In general relativity, this is the boundary hypersurface where Israel junction conditions require continuity of the induced metric and matching of the extrinsic curvature, fields on both sides of Σ must agree [**?** **?**]. In bimetric gravity, both metric sectors must simultaneously satisfy junction conditions at Σ through the interaction potential, with the constraint surface in phase space defining the locus where all first-class constraints vanish [**?** **?**]. *Status: Textbook.*

Field Closure ($\mathbf{FC}_\Sigma$)

The requirement that any coupled gravity fields must settle into a single consistent geometric state. In the ADM formalism, the Hamiltonian and momentum constraints must be satisfied on every spatial hypersurface, the gravitational field cannot exist in an inconsistent state [**?**]. In the Hassan-Rosen bimetric formalism, field closure requires simultaneous constraint satisfaction across both metric sectors, the massless and massive spin-2 modes, through the interaction potential $V(g, f)$, where the ghost-free condition eliminates the unphysical sixth

degree of freedom [**?**]. *Status: Extended.*

Field Origin

The coupling point at $r = 0$ where propagation time vanishes ($\tau = 0$), enabling field closure to complete atomically across the entire field configuration. In general relativity, the center of a self-gravitating system is where the metric potential is extremal, the lapse function reaches its minimum, and constraint coupling is maximally nonlinear [**?**]. In the Hassan-Rosen bimetric formalism, the Field Origin is the interaction vertex where both metric sectors couple through $V(\sqrt{g^{-1}f})$ at the same spacetime point, no propagation between sectors is required because the coupling is local, and where Vainshtein screening produces dominant nonlinear self-coupling of the massive mode [**?** **?**]. *Status: Extended.*

Full State Snapshot (FSS)

Complete, lossless read of the entire unified field configuration required for field closure to complete. No partial reads permitted. In the ADM formalism, this corresponds to the specification of complete initial data (q_{ab}, π^{ab}) on a Cauchy surface, the well-posed initial value problem requires the full 3-metric and conjugate momentum everywhere on the hypersurface; partial data admits no unique solution [**?** **?**]. In bimetric gravity, the Full State Snapshot must include both metric sectors simultaneously, the initial data for $g_{\mu\nu}$ and $f_{\mu\nu}$ cannot be specified independently because the interaction potential $V(\sqrt{g^{-1}f})$ couples them at every point [**?**]. *Status: Extended.*

Settlement

The resolution of coupling to a unique, consistent geometric state. After the Full State Snapshot captures the complete configuration, settlement resolves it to consistency through constraint satisfaction. In numerical relativity, this corresponds to solving the Hamiltonian and momentum constraints as elliptic PDEs, the York-Lichnerowicz conformal method guarantees a unique solution satisfying all constraints on each spatial hypersurface [**?** **?**]. In

bimetric gravity, settlement is the process by which both metric sectors achieve mutual consistency through the interaction potential, where the ghost-free structure guarantees exactly one physical configuration satisfies all constraints [**?**]. *Status: Extended.*

Registration (B^*_Σ)

The recorded outcome of settlement: the boundary data at Σ that determines the settled potential Φ^* throughout the domain. Gradients of this registered state are gravity, lensing, and time dilation, three manifestations of the same underlying field dynamics. In canonical gravity, B^*_Σ is the point on the constraint surface where all first-class constraints are satisfied, the on-shell configuration from which all observables are computed [**? ?**]. *Status: Novel.*

---

[1] H. S. M. Coxeter, *Regular Polytopes*, 3rd ed. (Dover Publications, 1973).

[2] S. Alexander, A. Marcianò, and L. Smolin, "Gravitational origin of the weak interaction's chirality," Phys. Rev. D **89**, 065017 (2014).

[3] S. Weinberg, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity* (Wiley, 1972).

[4] S. Weinberg, *The Quantum Theory of Fields, Vol. 1: Foundations* (Cambridge University Press, 1995).

[5] R. Geroch, "Spinor structure of space-times in general relativity. I.," J. Math. Phys. **9**, 1739 (1968).

[6] A. Ashtekar, "New variables for classical and quantum gravity," Phys. Rev. Lett. **57**, 2244 (1986).

[7] S. F. Hassan and R. A. Rosen, "Bimetric gravity from ghost-free massive gravity," JHEP **02**, 126 (2012).

[8] C. Deffayet, J. Mourad, and G. Zahariade, "A note on 'symmetric' vielbeins," JHEP **03**, 086 (2013).

[9] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), "Observation of gravitational waves from a binary black hole merger," Phys. Rev. Lett. **116**, 061102 (2016).

[10] B. Bertotti, L. Iess, and P. Tortora, "A test of general relativity using radio links with the Cassini spacecraft," Nature **425**, 374 (2003).

[11] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), "A gravitational-wave standard siren measurement of the Hubble constant," Nature **551**, 85 (2017).

[12] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (W. H. Freeman, San Francisco, 1973).

[13] R. M. Wald, *General Relativity* (University of Chicago Press, Chicago, 1984).

[14] R. Arnowitt, S. Deser, and C. W. Misner, "The Dynamics of General Relativity," Gen. Relativ. Gravit. **40**, 1997 (2008).

[15] G. M. Ziegler, *Lectures on Polytopes*, Graduate Texts in Mathematics, Vol. 152 (Springer-Verlag, 1995).

[16] T. C. Hales, "A proof of the Kepler conjecture," Ann. Math. **162**, 1065 (2005).

[17] D. G. Boulware and S. Deser, "Can gravitation have a finite range?" Phys. Rev. D **6**, 3368 (1972).

[18] J. D. Jackson, *Classical Electrodynamics*, 3rd ed. (Wiley, 1999).

[19] F. W. Hehl, P. von der Heyde, G. D. Kerlick, and J. M. Nester, "General relativity with spin and torsion: Foundations and prospects," Rev. Mod. Phys. **48**, 393 (1976).

[20] M. Milgrom, "A modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis," Astrophys. J. **270**, 365 (1983).

[21] E. Tiesinga, P. J. Mohr, D. B. Newell, and B. N. Taylor, "CODATA recommended values of the fundamental physical constants: 2022," Rev. Mod. Phys. **96**, 025004 (2024).

[22] S. Navas *et al.* (Particle Data Group), "Review of Particle Physics," Phys. Rev. D **110**, 030001 (2024).

[23] H. Goldstein, C. Poole, and J. Safko, *Classical Mechanics*, 3rd ed. (Addison-Wesley, 2002).

[24] R. P. Feynman, *QED: The Strange Theory of Light and Matter* (Princeton University Press, 1985).

[25] M. Planck, letter to R. W. Wood, 7 October 1931. Reprinted in H. Kangro, *Planck's Original Papers in Quantum Physics* (Taylor & Francis, 1972).

[26] N. Aghanim *et al.* (Planck Collaboration), "Planck 2018 results. VI. Cosmological parameters," Astron. Astrophys. **641**, A6 (2020).

[27] A. G. Riess *et al.*, "A comprehensive measurement of the local value of the Hubble constant with 1 km/s/Mpc uncertainty from the Hubble Space Telescope and the SH0ES team," Astrophys. J. Lett. **934**, L7 (2022).

[28] J. Schwinger, "On Quantum-Electrodynamics and the Magnetic Moment of the Electron," Phys. Rev. **73**, 416 (1948).

[29] A. Barbieri, "Quantum tetrahedra and simplicial spin networks," Nucl. Phys. B **518**, 714 (1998).

[30] K. Hinterbichler and R. A. Rosen, "Interacting spin-2 fields," JHEP **07**, 047 (2012).

[31] S. F. Hassan and M. Kocic, "On the local structure of spacetime in ghost-free bimetric theory and massive gravity," JHEP **05**, 099 (2018).

[32] L. T. Buchman and J. M. Bardeen, "A hyperbolic tetrad formulation of the Einstein equations for numerical relativity," Phys. Rev. D **67**, 084017 (2003).

[33] D. Clowe, M. Bradăc, A. H. Gonzalez, M. Markevitch, S. W. Randall, C. Jones, and D. Zaritsky, "A direct empirical proof of the existence of dark matter," Astrophys. J. **648**, L109 (2006).

[34] A.-C. Eilers, D. W. Hogg, H.-W. Rix, and M. K. Ness, "The circular velocity curve of the Milky Way from 5 to 25 kpc," Astrophys. J. **871**, 120 (2019).

[35] X. Hernandez, S. Mendoza, S. Contreras, and L. A. Martínez-Medina, "Wide binaries as a critical test of classical gravity," Mon. Not. R. Astron. Soc. **509**, 2304 (2022).

[36] X. Hernandez, K. Crépin, S. Scarano Jr., and R. Wilhelm, "The wide binary test of gravity and MOND with Gaia DR3," Mon. Not. R. Astron. Soc. **525**, 1401 (2023).

[37] M. Kramer, I. H. Stairs, R. N. Manchester, M. A. McLaughlin, A. G. Lyne, R. D. Ferdman, M. Burgay, D. R. Lorimer, A. Possenti, N. D'Amico, J. M. Sarkissian, G. B. Hobbs, J. E. Reynolds, P. C. C. Freire, and F. Camilo, "Tests of general relativity from timing the double pulsar," Science **314**, 97 (2006).

[38] R. W. Pattie Jr. *et al.*, "Measurement of the neutron lifetime using a magneto-gravitational trap and in situ detection," Science **360**, 627 (2018).

[39] J. G. Williams, S. G. Turyshev, and D. H. Boggs, "Progress in lunar laser ranging tests of relativistic gravity," Phys. Rev. Lett. **93**, 261101 (2004).

[40] A. T. Yue, M. S. Dewey, D. M. Gilliam, G. L. Greene, A. B. Laptev, J. S. Nico, W. M. Snow, and F. E. Wietfeldt, "Improved determination of the neutron lifetime," Phys. Rev. Lett. **111**,

222501 (2013).

[41] J. A. Wheeler, "Geons," Phys. Rev. **97**, 511 (1955).

[42] W. Israel, "Singular hypersurfaces and thin shells in general relativity," Nuovo Cimento B **44**, 1 (1966).

[43] P. A. M. Dirac, *Lectures on Quantum Mechanics* (Belfer Graduate School of Science, Yeshiva University, New York, 1964).

[44] A. I. Vainshtein, "To the problem of nonvanishing gravitation mass," Phys. Lett. B **39**, 393 (1972).

[45] Y. Choquet-Bruhat, "Théorème d'existence pour certains systèmes d'équations aux dérivées partielles non linéaires," Acta Math. **88**, 141 (1952).

[46] J. W. York Jr., "Kinematics and dynamics of general relativity," in *Sources of Gravitational Radiation*, edited by L. L. Smarr (Cambridge University Press, 1979), p. 83.

[47] M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems* (Princeton University Press, 1992).

[48] E. M. Burbidge, G. R. Burbidge, W. A. Fowler, and F. Hoyle, "Synthesis of the elements in stars," Rev. Mod. Phys. **29**, 547 (1957).