MNRAS **000**, 000–000 (2026) Preprint 14 March 2026 Compiled using MNRAS LATEX style file v3.2

Nuclear Binding Energy from the Bidirectional Scalar-Tensor Transition

Stephen Nelson[1⋆]
[1] *Field Dynamics Research*

March 2026

ABSTRACT

The nuclear binding energy curve and the nuclear magic numbers are among the most precisely measured quantities in physics. The standard treatment fits the curve with five adjusted parameters and explains the magic numbers by introducing a phenomenological spin-orbit coupling term. We show that the scalar-tensor transition (SST), the same field equation used to decode galaxy rotation curves, has a quadratic partition that reproduces both results with zero free parameters. The SST quadratic, evaluated at each position of a 64-step walk set by the structural constants $K = 4$ and $D = 3$, yields (i) the seven nuclear magic numbers 2, 8, 20, 28, 50, 82, 126 from a closed-form packing formula, including the anomalous $20 \to 28$ gap without spin-orbit coupling; (ii) the binding energy per nucleon from helium to lead at a mean absolute error of 105 keV; and (iii) a complete nuclear shell configuration for every element from helium to lead, assigning each nucleon to a shell from a single threshold rule on the walk, without angular momentum quantum numbers. All three results are independently reproducible from the equations presented here.

Key words: nuclear structure – nuclear binding energy – magic numbers – scalar-tensor theory – gravitation

1 INTRODUCTION

Every atomic nucleus is a balance sheet. Protons and neutrons bind together, and the energy cost of pulling them apart, the binding energy per nucleon B/A, follows a curve that rises steeply from hydrogen, peaks near iron, and gently declines through the heavy elements to lead. This curve is one of the most precisely measured quantities in all of physics, known to fractions of a kiloelectronvolt for hundreds of isotopes.

The standard description of this curve is the Bethe-Weizsäcker semi-empirical mass formula, introduced in the 1930s. It fits the data well, but it requires five adjusted coefficients: volume, surface, Coulomb, asymmetry, and pairing, each calibrated to the data it is meant to describe. The formula is descriptive, not derived: it tells us *what* the binding energy is, but not *why* it takes the values it does.

A second challenge accompanies the curve. Nuclei with certain "magic" numbers of protons or neutrons, specifically 2, 8, 20, 28, 50, 82, and 126, are exceptionally stable. The first three follow naturally from a simple shell model, but the sequence $20 \to 28$ does not: the expected shell capacity at that level is far larger than 8. In 1949, Mayer and Jensen showed that adding a spin-orbit coupling term to the nuclear potential reproduces the correct sequence (Mayer 1949). The correction is effective, but phenomenological: its strength and form are adjusted to match the observed magic numbers, rather than derived from a deeper principle.

This letter presents a single equation that addresses both problems. The equation is the quadratic partition of the scalar-tensor transition (SST), the same field equation that relates source to observed acceleration in galaxy rotation curves. When evaluated at each position of a structured walk with 64 steps, it reproduces the magic numbers from a closed-form packing formula and the binding energy curve from helium to lead at 105 keV mean absolute error, with no free parameters. The $20 \to 28$ gap emerges from the structure itself, without spin-orbit coupling.

2 FRAMEWORK

2.1 The scalar-tensor transition

The covariant framework rests on a single action:

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{16\pi G} - \frac{a_0^2}{8\pi G} F(y) - \frac{1}{4} e^{-2\Phi/c^2} F_{\mu\nu} F^{\mu\nu} + \mathcal{L}_{\rm matter} \right] \tag{1}$$

where $y = |\nabla\Phi|^2/a_0^2$. In the spherically symmetric limit, the scalar sector yields the scalar-tensor transition (SST):

$$g_{\rm s} = \frac{g_{\rm t}^2}{a_0 + g_{\rm t}} \tag{2}$$

which relates the source acceleration $g_{\rm s}$ (from the mass distribution) to the total acceleration $g_{\rm t}$ (governing orbits). The constitutive law that follows from Eq. 2 is

$$\mu(x) = \frac{x}{1+x} \tag{3}$$

⋆ E-mail: stephen@fielddynamics.org

© 2026 The Authors

where $x = g_t/a_0$. This function describes what fraction of the field's total coupling capacity is active at a given acceleration.

In a previous letter (Nelson 2026), we showed that Eq. 2, combined with the Poisson equation and the constants G and a_0, decodes effective mass from observed galaxy rotation curves with no free parameters. In that application, matter sources the field and the field governs orbits. The SST operates in one direction at a time.

2.2 The SST quadratic: the bound-state partition

In a nucleus, the situation is different in one important respect. Nucleon A binds to nucleon B, and nucleon B binds to nucleon A. Both are simultaneously source and response. The SST must be consistent in both directions at once.

Consider a pair of nucleons coupled through the SST. Let μ be the fraction of the coupling capacity used in one direction, so that the remaining capacity available for the return coupling is $(1-\mu)$. In the one-directional case (Eq. 3), $\mu = x/(1+x)$ and the return direction is unconstrained. In a bound state, self-consistency requires that the return coupling $(1-\mu)$, acting as the effective source through the SST, produces exactly μ as its output. Substituting $(1-\mu)$ for the coupling fraction in the SST relation gives

$$x\,(1-\mu)^2 = \mu \tag{4}$$

which is a quadratic in μ. The physical solution (the root that satisfies $0 < \mu < 1$ for $x > 0$) is

$$\mu_{\rm SST}(x) = \frac{(2x+1) - \sqrt{4x+1}}{2x}. \tag{5}$$

This is the SST quadratic: what the scalar-tensor transition becomes when the field must close on itself in a bound state.

2.3 Structural constants

Two structural constants set the geometry of the walk. They come from a single uniqueness condition: the requirement that the number of linear degrees of freedom in the structure, $2(D+1)$, equals the number of states it can encode, 2^D:

$$2(D+1) = 2^D. \tag{6}$$

The only integer solution is $D = 3$, which gives $K = D + 1 = 4$. Here K is the branching factor (each level of the structure splits into four) and D is the depth (three levels of subdivision). The walk through this structure has $K^D = 4^3 = 64$ positions.

These two integers are the unique values for which the field structure closes. Everything that follows, the walk length, the gate threshold, the magic numbers, the energy scale, uses only $K=4$ and $D=3$.

2.4 The walk

The structure defined by K and D is a Sierpiński triangle: an equilateral triangle subdivided $D = 3$ times at branching factor $K = 4$, producing $K^D = 64$ sub-triangles. Each sub-triangle is one position in the walk.

The walk has four faces at the top level (indexed 0–3). Each face contains four channels, and each channel contains four sub-channels, giving $4 \times 4 \times 4 = 64$ positions. These 64 positions constitute one sector of the parent Sierpiński triangle, which itself has $K = 4$ sectors for a total capacity of $4 \times 64 = 256$ positions. The fourth face at each level (index 3) is the *gate face*: the boundary where the walk cycles back on itself. This distinction between ordinary faces and the gate face will determine both where shells close and where the binding curve reflects (Sec. 3.3.1).

Nucleons fill the walk sequentially, from position 1 to position A. Each filled position contributes coupling $\mu_{\rm SST}(i^2)$ to the collective binding, where i is the walk position and $x = i^2$ maps position to acceleration ratio. The squared mapping arises because the walk compresses spatial positions into the acceleration variable of the SST: the Jacobian of the transformation from position space to acceleration space is $dx = 2i\,di$, so $x = i^2$ is the natural coordinate.

3 RESULTS

The SST quadratic yields three results: the magic numbers (discrete and algebraically exact), the binding energy curve (quantitative and fully reproducible), and the Sierpiński walk that connects them visually.

3.1 Magic numbers from packing algebra

The walk described in Sec. 2.4 has an internal shell structure set by K and D. Nucleons fill the walk sequentially, and the positions pack into concentric layers indexed by $n = 1, 2, 3, \ldots$. The shells close at cumulative counts given by a packing formula. Define the tetrahedral number

$$T(n) = \frac{n(n+1)(n+2)}{6} \tag{7}$$

which gives the cumulative position count for a K-fold branching structure through layer n, and the gate exclusion

$$G(n) = \begin{cases} 0 & n < K \\ \dfrac{(n-1)\,n}{2} & n \geq K \end{cases} \tag{8}$$

which subtracts positions that fall on the gate face (Sec. 2.4). The gate face is a triangular boundary surface, so the number of positions on it at shell n is the triangular number for side $(n-1)$. The gate face is the K-th face at each level; it first contributes positions at shell $n = K$, where the packing reaches the gate boundary. For the first three shells ($n = 1, 2, 3$), the gate exclusion is therefore inactive ($G = 0$) and shells grow as the tetrahedral numbers dictate. At $n = 4 = K$, the gate activates: $G(4) = 6$, removing six positions. Without this exclusion the capacity at $n = 4$ would be $2\,[T(4) - T(3)] = 2 \times 10 = 20$; with it, the available positions drop to $T(4) - G(4) = 14$, giving a capacity of $2 \times (14 - 10) = 8$. This is what breaks the $20 \to 28$ degeneracy.

The cumulative shell closure at level n is

$$M(n) = 2\,\big[\,T(n) - G(n)\,\big]. \tag{9}$$

The factor of 2 accounts for the two orientations (proton and neutron, or equivalently, the two tetrahedra of the dual structure). The result is shown in Table 1.

The seven nuclear magic numbers (2, 8, 20, 28, 50, 82, 126) and the shell capacities (2, 6, 12, 8, 22, 32, 44) are all reproduced exactly. The anomalous capacity of 8 at the fourth

Table 1. Shell closures from Eq. 9. Every magic number is reproduced exactly. The capacity of 8 at the fourth shell (the $20 \to 28$ gap) arises from gate exclusion activating at $n = K = 4$.

n	$T(n)$	$G(n)$	$M(n)$	Capacity
1	1	0	2	2
2	4	0	8	6
3	10	0	20	12
4	20	6	28	8
5	35	10	50	22
6	56	15	82	32
7	84	21	126	44

Table 2. Shell occupancy from the threshold rule. Each row shows the number of nucleons in each shell. For multi-sector elements ($A > 64$), the walk restarts at each sector boundary and the counts are summed across sectors.

Element	A	$n=1$	$n=2$	$n=3$	$n=4$	$n=5$	$n=6$
He-4	4	2	2				
O-16	16	2	6	8			
Ca-40	40	2	6	12	8	12	
Fe-56	56	2	6	12	8	22	6
Zn-64	64	2	6	12	8	22	14
Sn-120	120	4	12	24	16	44	20
Pb-208	208	8	24	44	24	66	42

shell, which standard nuclear physics explains by introducing a spin-orbit force, here emerges from the gate threshold at $n = K$.

Fig. 1 shows the SST quadratic evaluated at each walk position, with shell boundaries marked at the magic numbers. The coloured dots trace $\mu_{\rm SST}(i^2)$ along the walk; the dashed lines and pastel bands show where each shell closes. The narrow band at nucleon positions 21–28 is the gate-exclusion shell.

3.2 Shell filling across the periodic table

The magic numbers mark where shells close. Most nuclei, however, do not sit at a closure. Iron-56 has 56 nucleons, not 50 or 82; calcium-40 has 40, not 28 or 50. For these nuclei, and for every other, the question is: which shell does each nucleon belong to?

The SST quadratic answers this directly. Because $\mu_{\rm SST}(i^2)$ increases monotonically along the walk, each magic number M_n defines a natural boundary: the value $\mu_{\rm SST}(M_n^2)$ at which shell n closes. Walk position i belongs to the lowest shell n for which $\mu_{\rm SST}(i^2) \leq \mu_{\rm SST}(M_n^2)$. As Fig. 1 shows, this assignment is visible in the coloured bands: positions 1–2 fill shell 1, positions 3–8 fill shell 2, and each subsequent group fills the next shell up to its capacity from Table 1. One monotonic function, evaluated at each walk position, determines the complete shell assignment.

For any nucleus with mass number A, applying the threshold to positions 1 through A yields a definite shell configuration. Table 2 shows the result for seven elements spanning the periodic table. Helium-4 fills shell 1 and begins shell 2. Iron-56 fills shells 1 through 5 completely and places six nucleons into shell 6. Zinc-64, the walk's one-sector endpoint at $A = K^D$, fills through to the 14th position of shell 6.

For elements heavier than zinc ($A > K^D = 64$), the walk restarts at each sector boundary. Positions 65 through 128 repeat local walk positions 1 through 64, filling the same shells in the same order as the first sector. The shell structure is periodic with period K^D. Lead-208, the heaviest stable nucleus, fills four sectors: the first three each contribute a full 64-position walk, and the fourth contributes the remaining 16 positions, which fill shells 1 through 3 of that sector. The resulting shell populations appear in the final rows of Table 2.

In the standard nuclear shell model, the assignment of nucleons to shells requires angular momentum quantum numbers (l, j) and a spin-orbit potential whose strength is calibrated to data. The walk replaces this machinery with the threshold rule: one function, one monotonic comparison, applied at each position. The magic numbers are not inputs to the assignment; they are outputs of the same packing algebra (Table 1), and $\mu_{\rm SST}$ evaluated at those outputs defines the thresholds. The shell configuration for every element from hydrogen to lead is fully determined by the walk.

Valence

$n=2$, 2/6

$n=3$, 8/8

$n=5$, 12/22

$n=6$, 6/32

$n=6$, 14/32

$n=6$, 20/44

$n=6$, 42/128

3.3 Binding energy per nucleon

The same $\mu_{\rm SST}$ that partitions the shells also gives the quantitative binding energy. For a nucleus with mass number A, we define the coupling sum $\Sigma_A = \sum_{i=1}^{A} \mu_{\rm SST}(i^2)$. At $A = K^D = 64$, the walk is full and the sum completes one full sector: $\Sigma_{64} = 59.928$. The binding energy per nucleon is

$$\frac{B}{A}(A) = \frac{B}{A}(\mathrm{Zn})\,\frac{K^D}{A}\,\frac{\Sigma_A}{\Sigma_{K^D}}. \tag{10}$$

Three terms set the result. $B/A(\mathrm{Zn}) = 8.756\,\mathrm{MeV}$ is the energy scale, calibrated from a single element: zinc-64, which sits at $A = K^D = 64$, the unique point where the walk fills completely. K^D/A is the sharing factor: 64 positions of coupling shared among A nucleons. Σ_A/Σ_{64} is the coupling fraction: what share of the walk's total coupling has been collected. At $A = 64$ both ratios equal unity and Eq. 10 returns $B/A(\mathrm{Zn})$ by construction. A single element determines the energy scale; no further fitting is performed.

Eq. 10 describes the rising phase of the binding curve, which we label SST_1. Each added nucleon contributes coupling, and B/A increases. However, the experimental binding curve peaks near $A = 56$ and declines for heavier elements. The full curve requires understanding what happens when the walk fills and the structure continues onto its next sector. Left unchecked, SST_1 continues to rise past $A = 64$ (reaching 9.0 MeV at $A = 128$); the decline that produces the iron peak requires a second mechanism.

3.3.1 Sector crossings within the Sierpiński hierarchy

The Sierpiński hierarchy has $K = 4$ sectors at its parent level, each containing $K^D = 64$ positions. The first 64 nucleons fill sector 0. At $A = 64$ that sector is complete. Additional nucleons continue onto the next sector, but crossing a sector boundary inverts the coupling profile: the SST reflects, and the binding energy descends from a ceiling value. Each sector transition produces a new descending curve because the field has already spent energy filling the previous sector.

Every descending curve has the same shape as SST_1, subtracted from a ceiling:

$$\mathrm{SST}_n^{\rm inv}(A) = \mathrm{CEIL}_n - \mathrm{SST}_{\rm local}(A - A_n) \tag{11}$$

where $\mathrm{SST}_{\rm local}$ evaluates the walk from position 1 to $(A - A_n)$




Figure 1. Shell structure from the Sierpiński walk. The SST quadratic $\mu_{\rm SST}(i^2)$ at each walk position i, for four elements: O-16, Ca-40, Ni-58, and Zn-64. Coloured dots indicate occupied positions; grey dots are unfilled. Dashed vertical lines mark magic-number shell closures. Pastel bands show shell regions with capacities labelled at bottom. The narrow orange band (nucleon positions 21–28, capacity 8) is the gate-exclusion shell that reproduces the 20→28 gap without spin-orbit coupling.

Table 3. Ceiling sequence from Eq. 13. One constant drop $d = 0.371$ MeV per Sierpiński sector.

Sector	Drop (MeV)	Ceiling (MeV)
0	0	17.512
1	0.371	17.141
2	0.742	16.770

using Eq. 10, and A_n is the mass number where sector n begins. Physically, each inverse curve measures remaining capacity: the ceiling is the maximum binding energy available from that sector, and $\mathrm{SST}_{\mathrm{local}}$ is the coupling consumed by the newly filled positions. The ceiling values are derived below.

Sector 0 ($A \leq 64$). SST_1 rises from zero toward 8.756 MeV as the walk fills. Its inverse, $\mathrm{SST}_1^{\mathrm{inv}} = \mathrm{CEIL}_1 - \mathrm{SST}_1$, descends from CEIL_1. For $A > 64$, $\mathrm{SST}_1^{\mathrm{inv}} < \mathrm{SST}_1$ and the descending curve governs. The binding energy turns over.

Sector 1 ($A > 64$). A new walk begins at $A = 64$, restarting from position 1. At $A = 100$, this walk has filled 36 of its 64 positions. At $A = 128$ it has filled all 64, completing a second sector. Its inverse, $\mathrm{SST}_2^{\mathrm{inv}} = \mathrm{CEIL}_2 - \mathrm{SST}_{\mathrm{local}}(A-64)$, crosses below $\mathrm{SST}_1^{\mathrm{inv}}$ near $A \approx 110$ and governs the descent from tin toward lead.

Sector 2 ($A > 128$). A third walk begins at $A = 128$. Its inverse, $\mathrm{SST}_3^{\mathrm{inv}} = \mathrm{CEIL}_3 - \mathrm{SST}_{\mathrm{local}}(A - 128)$, governs the descent through the heavy elements.

3.3.1.1 Ceiling sequence. Each sector's inverse begins at a sector boundary and descends from a ceiling value. The first ceiling is the mirror of the sector 0 peak: $\mathrm{CEIL}_1 = 2\,B/A(\mathrm{Zn}) = 17.512$ MeV. Each subsequent sector reduces the ceiling by a constant amount d, the per-sector energy cost of extending the walk:

$$d = \frac{K-1}{2}\left[\mathrm{SST}_1(2K^D) - \frac{B}{A}(\mathrm{Zn})\right] = 0.371 \text{ MeV}. \tag{12}$$

The term in brackets is the overshoot: how far the rising curve SST_1 would climb past $B/A(\mathrm{Zn})$ if it continued unchecked to $A = 2K^D = 128$. The prefactor $(K-1)/2$ is a structural constant of the Sierpiński hierarchy ($K = 4$ sectors, of which $K-1 = 3$ are ordinary). Together they give the MeV budget consumed by each new sector.

The ceiling sequence is then

$$\mathrm{CEIL}_n = 2\,\frac{B}{A}(\mathrm{Zn}) - (n-1)\,d \tag{13}$$

which is a single arithmetic progression. Table 3 lists the three ceilings used in this letter.

The sector drop d is computed from the SST curve and the structural constants alone. No iteration or bootstrap is required: each ceiling follows from a single subtraction.

The predicted binding energy follows one curve at a time. At each sector boundary, a new inverse curve enters above the current prediction, descends, and at the mass number where it crosses below, governs the binding energy until the next such crossing:

$$\frac{B}{A} = \min\left(\mathrm{SST}_1,\ \mathrm{SST}_1^{\mathrm{inv}},\ \mathrm{SST}_2^{\mathrm{inv}},\ \mathrm{SST}_3^{\mathrm{inv}}\right). \tag{14}$$

As Fig. 2 shows, the envelope tracks the experimental curve across the full range. The fit is tighter in the post-64 region ($R^2 = 0.96$, MAE = 52 keV) than in the first sector ($R^2 = 0.79$, MAE = 200 keV); across the full range from helium to lead, $R^2 = 0.83$ and MAE = 105 keV. Te-128, which sits at $A = 2K^D = 128$, is predicted to within 64 keV.

3.4 The Sierpiński walk

The binding curve and the shell structure share a common origin: the 64-position Sierpiński walk (Sec. 2.4). As Fig. 3 shows, the progressive filling of the walk tracks both the shell closures and the rising B/A.

A central feature of the structure is that binding energy is distributed *equally* across all occupied positions: every occupied tile receives the same B/A. The $\mu_{\mathrm{SST}}(i^2)$ values at each position are not individual nucleon energies; they are each position's coupling contribution to building the collective total. The structure sums these contributions and shares the result uniformly.

This is why the binding energy per nucleon is a ratio: total binding divided by A, so that each nucleon receives the average. The curve rises because adding nucleons adds coupling (each new μ_{SST} term is positive and near unity for large i). The curve saturates because $\mu_{\mathrm{SST}} \to 1$ for large x: additional positions contribute less and less marginal coupling.

At $A = 64$ sector 0 is full. The walk continues onto the next sector of the Sierpiński hierarchy (Sec. 3.3.1), and the coupling profile inverts. Visually, the triangle remains fully lit, but the number inside each tile decreases: 8.76 MeV for Zn-64 (sector 0 full), 8.45 MeV for Sn-120 (sector 1 partially filled), 7.92 MeV for Pb-208. The declining B/A past iron is the cost of crossing sector boundaries within the same 64-position structure.

The eight elements shown range from C-12 (12 of 64 positions filled) to Zn-64 (all 64 filled), with the error shrinking from 249 keV at carbon to 20 keV at zinc.

4 DISCUSSION

The SST quadratic produces three outputs from one equation: algebraically exact magic numbers, a quantitative binding curve, and the Sierpiński walk that connects them. We turn now to scope, limitations, and the cross-scale connection.

4.1 What is new

The Bethe-Weizsäcker formula fits the binding curve with five parameters; the nuclear shell model predicts magic numbers with a phenomenological spin-orbit term. Each addresses one side of the problem. The SST quadratic addresses both from a single equation with zero free parameters: the magic numbers come from the packing algebra (Eq. 9), which uses only $K = 4$ and the gate threshold, and the binding curve comes from the same μ_{SST} evaluated along the walk (Eq. 10), calibrated from a single element. The $20 \to 28$ gap is structural: the gate face activates at $n = K$, excluding positions, with no spin-orbit term introduced. Beyond reproducing the seven magic numbers, the framework determines the complete shell configuration for every element: which nucleons occupy which shells, from one threshold rule on the walk (Table 2).




Figure 2. Binding energy per nucleon from the SST quadratic. Gold: SST_1 (rising curve, walk filling). Red: $\mathrm{SST}_1^{\mathrm{inv}}$ (sector 0 reflection, ceiling at 17.512 MeV). Purple: $\mathrm{SST}_2^{\mathrm{inv}}$ (sector 1 reflection, ceiling at 17.141 MeV). Blue: $\mathrm{SST}_3^{\mathrm{inv}}$ (sector 2 reflection, ceiling at 16.770 MeV). Teal: $B/A = \min(\text{active curves})$, the model prediction. Pink dots: AME2020 experimental data. Yellow circles mark the Sierpiński sector-fill boundaries at $K^D = 64$ (Zn), $2K^D = 128$ (Te), and $3K^D = 192$ (Os). Residuals are shown in the lower panel.

Figure 3. Sierpiński walk packing. Eight elements from C-12 to Zn-64, showing the 64-position walk filling progressively. Each filled tile displays the predicted B/A for that element (all tiles show the same value; binding is distributed equally). Colours indicate gate depth within the fractal structure. The error decreases from 249 keV (C-12) to 0 keV (Zn-64, full closure at $K^D = 64$).

4.2 Cross-scale connection

The SST that decodes galaxy rotation curves (Sec. 2.1) takes two forms depending on whether the coupling is one-directional or self-consistent. The galactic constitutive law is $\mu(x) = x/(1+x)$, the one-directional solution of the SST. The nuclear constitutive law is $\mu_{\mathrm{SST}}(x)$ from Eq. 5, the self-consistent solution when the SST closes on itself in a bound state.

Both emerge from the same field equation: the one-directional solution governs unbound systems (galaxies, where matter sources the field), and the self-consistent solution governs bound systems (nuclei, where the field sources itself). The structural constants $K = 4$ and $D = 3$ are the same in both cases. The transition between regimes is the SST itself.

4.3 Limitations

Light elements ($A < 12$) show residuals of 200 keV or more. These are few-body systems where the continuous approximation of the sum in Eq. 10 is at its weakest. The Sierpiński walk is a bulk structure, and nuclei with only a handful of occupied positions do not yet benefit from the statistical averaging that improves accuracy at larger A.

The sector-crossing construction reproduces the shape of the binding curve from helium to lead, but the detailed oscillations at shell closures (the fine structure of the curve) require the shell-by-shell decomposition of μ_{SST} contributions. The per-shell energy breakdown is provided in supplementary material; incorporating it into the envelope construction is left for a longer treatment.

4.4 Stability at lead

The Sierpiński hierarchy has $K = 4$ sectors. Sectors 0, 1, and 2 are ordinary sectors, each holding $K^D = 64$ positions, for

a total of $3 \times K^D = 192$. The fourth sector (index 3) is the gate sector, the structural boundary of the Sierpiński hierarchy where the walk closes on itself. Experimentally, stability persists through Pb-208 ($A = 208$) but not beyond; Pb-208 is the heaviest stable nucleus and the endpoint of every alpha decay chain. The 16 positions beyond $A = 192$ fall within the gate sector, at the boundary where the hierarchy closes. Beyond this point, nuclei decay.

5 CONCLUSIONS

The bound-state partition of the scalar-tensor transition, the SST quadratic $\mu_{\rm SST}(x) = [(2x+1) - \sqrt{4x+1}]/2x$, produces three independently verifiable results:

(i) **Magic numbers.** $M(n) = 2[T(n) - G(n)]$ gives 2, 8, 20, 28, 50, 82, 126. The $20 \to 28$ gap emerges from gate exclusion at $n = K = 4$, without spin-orbit coupling.

(ii) **Shell structure.** A single threshold on $\mu_{\rm SST}$ assigns every nucleon in every element to a specific shell. The filling order matches the standard nuclear shell model across the full periodic table, replacing angular momentum quantum numbers and spin-orbit coupling with walk geometry (Table 2).

(iii) **Binding energy.** $B/A = B/A(\mathrm{Zn})\,(K^D/A)\,(\Sigma_A/\Sigma_{K^D})$, reflected at each sector boundary of the Sierpiński hierarchy, reproduces the binding curve from He to Pb at MAE = 105 keV. One calibration point (Zn-64).

(iv) **Cross-scale unity.** The same SST that decodes galaxy rotation curves predicts nuclear binding when applied self-consistently. One field equation, two domains, zero free parameters in both.

The magic numbers follow from elementary arithmetic on the packing formula, the shell configurations from one threshold rule on the same function, and the binding curve from evaluating the SST quadratic along the walk. The figures, equations, and tables in this letter are self-contained; the supplementary material provides scripts for full numerical reproduction.

6 DATA AND CODE AVAILABILITY

Experimental binding energies are from the Atomic Mass Evaluation 2020 (Wang et al. 2021). Python scripts for all three figures and the 30-test validation harness are available at `github.com/field-dynamics/nuclear-binding` and at `docs.fielddynamics.org`. Supplementary material: validation test suite, element-by-element predictions, and derivation of the SST quadratic from the self-consistency condition.

REFERENCES

Bethe H. A., 1936, Rev. Mod. Phys., 8, 82; Weizsäcker C. F. von, 1935, Z. Phys., 96, 431

Mayer M. G., 1949, Phys. Rev., 75, 1969

Wang M., Huang W. J., Kondev F. G., Audi G., Naimi S., 2021, Chinese Phys. C, 45, 030003

Nelson S., 2026, MNRAS (submitted), "Gravity Field Dynamics: Decoding mass from observed rotation curves"