

The slopes of the two tangents of inflection are given by the expression

$$q \pm \left(\frac{-2p}{3} \right)^{3/2}.$$

If the slope of one inflection tangent is zero, then the slope of the other is $2q$.

MATHEMATICAL NOTES

EDITED BY E. F. BECKENBACH, University of California
and Institute for Numerical Analysis of the National Bureau of Standards

Material for this department should be sent directly to E. F. Beckenbach, University of California, Los Angeles 24, California.

A GENERALIZED INTEGRAL TEST FOR CONVERGENCE OF SERIES

MORGAN WARD, California Institute of Technology

The following useful generalization of the familiar Maclaurin-Cauchy integral test for convergence of real series deserves to be better known. It is apparently due to G. H. Hardy,* who made a redundant hypothesis on $f(t)$. The integrals may be taken either in the sense of Riemann or in the sense of Lebesgue.

THEOREM. *Let $f(t)$ be a complex-valued function of the real variable in the interval $1 \leq t < \infty$, such that $f'(t)$ exists and is integrable to $f(t)$ over any finite interval $1 \leq t \leq T$. Then if $\int_1^\infty f'(t)dt$ is absolutely convergent, the series $\sum_1^\infty f(n)$ and the integral $\int_1^\infty f(t)dt$ converge and diverge together.*

Proof: By Abel's partial summation formula, we have

$$\sum_{r=1}^n a_r b_r = A_n B_n - \sum_{r=1}^{n-1} A_r (b_{r+1} - b_r),$$

where $A_r = a_1 + a_2 + \cdots + a_r$, ($r = 1, 2, \cdots, n$).

Let $s_n = \sum_{r=1}^n f(r)$. Then on taking $a_r = 1$ and $b_r = f(r)$ in the summation formula, we find that

$$s_n = nf(n) - \sum_{r=1}^{n-1} r(f(r+1) - f(r)).$$

Now if $[t]$ denotes as usual the greatest integer in t , then

* G. H. Hardy: Proc. London Math. Soc. (2), vol. 9, 1910, pp. 126-144.

$$r(f(r+1) - f(r)) = \int_r^{r+1} [t]f'(t)dt.$$

Also

$$nf(n) - 1 \cdot f(1) = \int_1^n \frac{d}{dt} (tf(t))dt,$$

or

$$nf(n) = f(1) + \int_1^n f(t)dt + \int_1^n tf'(t)dt.$$

On substituting these expressions into the formula for s_n , simplifying and transposing, we obtain the formula

$$s_n - \int_1^n f(t)dt = f(1) + \int_1^n (t - [t])f'(t)dt.$$

Now $|(t - [t])f'(t)| < |f'(t)|$. Hence the infinite integral $\int_1^\infty (t - [t])f'(t)dt$ is convergent, and

$$(1) \quad \lim_{n \rightarrow \infty} \left(s_n - \int_1^n f(t)dt \right) \text{ exists.}$$

Now assume that the integral $\int_1^\infty f(t)dt$ is convergent. Then $\lim_{n \rightarrow \infty} \int_1^n f(t)dt$ exists. Hence by (1), $\lim_{n \rightarrow \infty} s_n$ exists; that is, the series $\sum_1^\infty f(n)$ is convergent.

The converse result is a little more troublesome. Assume that $\sum_1^\infty f(n)$ converges. Then

$$(2) \quad \lim_{n \rightarrow \infty} f(n) = 0,$$

and by (1),

$$(3) \quad \lim_{n \rightarrow \infty} \int_1^n f(t)dt \text{ exists.}$$

Now $f(T) = f(1) + \int_1^T f'(t)dt$. But since $\int_1^\infty f'(t)dt$ converges, $\lim_{T \rightarrow \infty} \int_1^T f'(t)dt$ exists. Hence $\lim_{T \rightarrow \infty} f(T)$ exists, so that by (2),

$$(4) \quad \lim_{t \rightarrow \infty} f(t) = 0.$$

Now

$$\begin{aligned} \left| \int_1^T f(t)dt - \int_1^{[T]} f(t)dt \right| &= \left| \int_{[T]}^T f(t)dt \right| \leq \max_{[T] \leq t \leq T} |f(t)| (T - [T]) \\ &< \max_{t \geq [T]} |f(t)|. \end{aligned}$$

Hence by (4)

$$\lim_{T \rightarrow \infty} \left(\int_1^T f(t) dt - \int_1^{[T]} f(t) dt \right) = 0.$$

But

$$\lim_{T \rightarrow \infty} \int_1^{[T]} f(t) dt$$

exists by (3). Hence $\lim_{T \rightarrow \infty} \int_1^T f(t) dt$ exists; that is $\int_1^\infty f(t) dt$ is convergent.

As an example, suppose that $f(t) = t^{-1} e^{-i\mu \log t}$, μ real. Then $f'(t) = O(1/t^2)$ and the conditions of the theorem are met. But

$$\int_1^T f(t) dt = \frac{i}{\mu} (e^{-i\mu \log T} - 1).$$

Hence $\int_1^\infty f(t) dt$ diverges. Therefore $\sum_1^\infty 1/n^{1+i\mu}$ diverges.

Again, suppose that $f(t) = e^{i\alpha \log t}/t^\beta$, where α and θ are real, and $\text{Re } \beta > \alpha > 0$, $\theta \neq 0$. Then $f'(t)$ is continuous and of order $t^{-1-\mu}$, where $\mu = \text{Re } \beta - \alpha$, in the range $1 \leq t < \infty$. Hence the conditions of the theorem are met. Now the infinite integral $\int_1^\infty f(t) dt$ is easily seen to converge on making the change of variable $s = t^\alpha$. Hence the infinite series $\sum_1^\infty e^{i\alpha \log n}/n^\beta$ converges. In particular then, if β is real, we see that the two real series

$$\sum_1^\infty \frac{\cos n^\alpha \theta}{n^\beta} \quad \text{and} \quad \sum_1^\infty \frac{\sin n^\alpha \theta}{n^\beta}$$

both converge if $\beta > \alpha > 0$ and $\theta \neq 0$.

The ordinary integral test is included as a special case if we use Lebesgue integrals; for if $f(t)$ is real, continuous and tends to zero steadily, $f'(t)$ exists almost everywhere and $f(t) = \int_1^t f'(s) ds + f(1)$. Since $|f'(t)| = -f'(t)$, the hypotheses of the theorem are evidently satisfied.

GEOMETRY OF THE SQUARE ROOT OF THREE

C. S. OGILVY, Trinity College, Hartford, Conn.

That the diagonal of a square is incommensurable with its side and the quotient is representable by the continued fraction

$$1 + \frac{1}{2 + \frac{1}{2 + \frac{1}{2 + \dots}}}$$

is easy to prove geometrically. The corresponding fact that the altitude of an equilateral triangle and half its side are incommensurable and the quotient representable by