

A CERTAIN CLASS OF POLYNOMIALS.*

BY MORGAN WARD.

Contents.

	Page
1. Definition of a Regular Sequence of Polynomials	43
2. Fundamental Properties of Regular Sequences	43
3. Harmonic Sequences	44
4. Condition a Harmonic Sequence be Regular	45
5, 6. Properties of the Simplest Regular Sequence	45, 47
7. Definition and Properties of Cyclic Sequences	47
8. Additional Properties of Cyclic Sequences	49

1. The remarkable properties which a sequence of polynomials

$$Y_0(x), Y_1(x), Y_2(x), \dots$$

of degrees $0, 1, 2, \dots$ in x possesses when the two functional equations

$$\frac{dY_n(x)}{dx} = Y_{n-1}(x), \quad Y_n(-x-1) = (-1)^n Y_n(x)$$

are satisfied have been systematically developed by N. Nielsen.† It is of some interest to consider sequences of polynomials satisfying the more general equations

$$(1) \quad \frac{dY_n(x)}{dx} = Y_{n-1}(x),$$

$$(2) \quad Y_n(ax+b) = \tau_n Y_n(x), \quad (n = 0, 1, 2, \dots)$$

where a, b are any complex numbers. Such a sequence we call a *regular* sequence. The main properties of regular sequences are as follows:

2. If a is not a root of unity, there is only one regular sequence; namely

$$Y_n(x) = \frac{c}{n!} \left(x + \frac{b}{a-1} \right)^n, \quad \tau_n = a^n \quad (n = 0, 1, 2, \dots).$$

If however, a is a root of unity, say a primitive π th root, there exist an infinite number of regular sequences. Let

$$(k_0 + k_1 t + k_2 t^2 + \dots) e^{xt} = K_0(x) + K_1(x) t + K_2(x) t^2 + \dots$$

* Received February 4, 1929.

† *Traité Élémentaire des Nombres de Bernoulli*, Paris 1923.

be the generating function of any such sequence. Then if $r\pi \leq n < (r+1)\pi$, k_n may be expressed as a linear function of $k_0, k_\pi, k_{2\pi}, \dots, k_{r\pi}$

$$k_n = \sum_{s=0}^r k_{s\pi} H'_{n-s\pi},$$

where H'_0, H'_1, H'_2, \dots depend only on a and b and are independent of the particular regular sequence $[K_n(x)]$ we have selected. Moreover, if a is a given π th root of unity, all possible solutions (1) and (2) are obtained by giving $k_0, k_\pi, k_{2\pi}, \dots$ the proper values.

3. We now proceed to the proof of these results. A sequence of polynomials

$$H_0(x), H_1(x), H_2(x), \dots$$

of degrees $0, 1, 2, \dots$ in x which satisfies (1) is said to be *harmonic*.* The following properties of harmonic sequences are easily proved.*

(i) If $[H_n(x)]$ is a harmonic sequence, there exists a sequence of constants

$$[h_n]: h_0, h_1, h_2, \dots, h_n, \dots,$$

such that for all values of n ,

$$H_n(x) = \frac{h_0 x^n}{n!} + \frac{h_1 x^{n-1}}{(n-1)!} + \frac{h_2 x^{n-2}}{(n-2)!} + \dots + h_n, \quad H_n(0) = h_n.$$

We denote such a sequence by $[H_n(x), h_n]$.

(ii) If†

$$\begin{aligned} H(x, t) &= H_0(x) + H_1(x)t + H_2(x)t^2 + \dots, \\ h(t) &= h_0 + h_1 t + h_2 t^2 + \dots \end{aligned}$$

are the generating functions of the sequences $[H_n(x)], [h_n]$

$$H(x, t) = e^{xt} h(t).$$

(iii) If b is any constant,

$$H_n(x+b) = H_n(b) + \frac{xH_{n-1}(b)}{1!} + \frac{x^2H_{n-2}(b)}{2!} + \dots + \frac{x^nH_0(b)}{n!}.$$

(iv) Let $[K_n(x), k_n]$ be a second harmonic sequence. Then there exists a unique ordinary sequence $[\alpha_n]$ such that for all values of n

$$K_n(x) = \alpha_0 H_n(x) + \alpha_1 H_{n-1}(x) + \dots + \alpha_n H_0(x).$$

* Nielsen, l. c., chap. III, section XI.

† This property of harmonic sequences is substantially due to Appell: *Annales de l'École Normale*, (2) 10 (1880), 119-120.

Moreover $\Delta_r(1) = 1$, $\Delta_r(0) = 1/r$, and $\Delta_r(a)$ vanishes if a is any primitive π th root of unity ($2 \leq \pi \leq r$).

6. From (8) and (i) it follows that

$$(9) \quad H_n(x) = \sum_{r=0}^n \frac{h_0}{r!(n-r)!} \lambda^{n-r} x^r = \frac{h_0}{n!} (x + \lambda)^n.$$

If (9) holds,

$$\frac{dH_n(x)}{dx} = H_{n-1}(x), \quad H_n(ax+b) = \frac{h_0}{n!} \left(ax+b + \frac{b}{a-1}\right)^n = a^n H_n(x),$$

so that (9) is a regular sequence whether or not a is a root of unity. On collecting these results, we have

THEOREM 3. A sequence $[H_n(x)]$ satisfying the two functional equations

$$(1) \quad \frac{dY_n(x)}{dx} = Y_{n-1}(x),$$

$$(2) \quad Y_n(ax+b) = a^n Y_n(x), \quad (a \neq 0, 1),$$

is always given by

$$(8) \quad h_n = \frac{h_0 \lambda^n}{n!},$$

$$(9) \quad H_n(x) = \frac{h_0 (x + \lambda)^n}{n!},$$

where h_0 is an arbitrary constant, and

$$(7) \quad \lambda = \frac{b}{a-1}.$$

Moreover if a is not a root of unity, this solution of (1) and (2) is unique.

7. The first result stated in section 2 is now proved. The regular sequences for which a is a root of unity are of much greater interest. It will be convenient in studying them to define $H_n(x)$ to mean the polynomial $\frac{(x+\lambda)^n}{n!}$ which gives the simplest regular sequence and to define a cyclic sequence of order π to mean any solution of (1) and (2) for which a is a primitive π th root of unity. Then

$$(10) \quad a^r = 1$$

when and only when $r \equiv 0 \pmod{\pi}$.

THEOREM 4. If $[K_n(x), k_n]$ is a cyclic sequence of order π then $K_n(x)$ may be uniquely represented in the form

$$(11) \quad K_n(x) = \alpha_0 H_n(x) + \alpha_\pi H_{n\pi}(x) + \dots + \alpha_{r\pi} H_{n-r\pi}(x)$$

where $\alpha_0, \alpha_\pi, \dots, \alpha_{r\pi}$ are constants, and $r\pi \leq n < (r+1)\pi$. Moreover every sequence of the form (11) is a cyclic sequence of order π .

For since $[K_n(x)]$ and $[H_n(x)]$ are both harmonic sequences, there exists by (iv) a unique ordinary sequence $[\alpha_n]$ such that

$$K_n(x) = \alpha_0 H_n(x) + \alpha_1 H_{n-1}(x) + \dots + \alpha_n H_0(x).$$

Hence the first part of the theorem will be proved if we can show $\alpha_r = 0$, $r \not\equiv 0 \pmod{\pi}$. Now

$$K_n(ax+b) = \sum_{r=0}^n H_{n-r}(ax+b) = \sum_{r=0}^n \alpha_r a^{n-r} H_{n-r}(x),$$

$$K_n(ax+b) = a^n K_n(x) = \sum_{r=0}^n \alpha_r a^n H_{n-r}(x).$$

Write $x+\lambda = y$ so that $H_{n-r}(x) = \frac{y^{n-r}}{(n-r)!}$. Then we have identically in y

$$\sum_{r=0}^n \frac{\alpha_r a^{n-r}}{(n-r)!} y^{n-r} = \sum_{r=0}^n \frac{\alpha_r a^n y^{n-r}}{(n-r)!}$$

so that by equating coefficients of corresponding powers of y ,

$$\alpha_r (a^r - 1) = 0 \quad (r = 0, 1, 2, \dots, n).$$

(11) now follows from (10). The last part of the theorem is obvious from (9) and (10).

If we write in (11) first $x = -\lambda$ and then $x = 0$ we obtain

THEOREM 5. For any cyclic sequence $[K_n(x)]$ of order π ,

$$\begin{aligned} K_n(-\lambda) &= 0, & n &\not\equiv 0 \pmod{\pi}, \\ &= \alpha_n, & n &\equiv 0 \pmod{\pi}, \end{aligned}$$

$$(12) \quad a^{-n} K_n(b) = K_n(0) = k_n = \frac{\alpha_0 \lambda^n}{n!} + \frac{\alpha_\pi \lambda^{n-\pi}}{(n-\pi)!} + \dots + \frac{\alpha_{r\pi} \lambda^{n-r\pi}}{(n-r\pi)!},$$

where $r\pi \leq n < (r+1)\pi$ and $n = 0, 1, 2, \dots$.

We have from (12)

$$\begin{aligned} k_0 &= \alpha_0, \\ k_\pi &= \frac{\alpha_0 \lambda^\pi}{\pi!} + \alpha_\pi, \\ k_{2\pi} &= \frac{\alpha_0 \lambda^{2\pi}}{(2\pi)!} + \frac{\alpha_\pi \lambda^\pi}{\pi!} + \alpha_{2\pi}, \\ k_{3\pi} &= \frac{\alpha_0 \lambda^{3\pi}}{(3\pi)!} + \frac{\alpha_\pi \lambda^{2\pi}}{(2\pi)!} + \frac{\alpha_{2\pi} \lambda^\pi}{\pi!} + \alpha_{3\pi}, \\ &\dots \end{aligned}$$

Hence $\alpha_0, \alpha_\pi, \alpha_{2\pi}, \dots$ are uniquely determined in terms of $k_0, k_\pi, k_{2\pi}, \dots$. Since (12) gives k_n for all values of n in terms of $\alpha_0, \alpha_\pi, \alpha_{2\pi}, \dots$ we have proved

THEOREM 6. *If $[K_n(x), k_n]$ is any cyclic sequence of order π and if the values of $k_0, k_\pi, k_{2\pi}, \dots$ are given, then all the other k_n are uniquely determined.*

8. We shall now give the explicit expression for k_n as a function of $k_0, k_\pi, k_{2\pi}, \dots$ proving the last result in section 2. It is convenient to borrow a definition from the calculus of generating functions.

Given any sequence of constants

$$[c_n]: \quad c_0, c_1, c_2, \dots \quad (c_0 \neq 0),$$

the sequence

$$[c'_n]: \quad c'_0, c'_1, c'_2, \dots$$

determined by the equations:

$$\begin{aligned} c'_0 c_0 &= 1 \\ c_0 c'_n + c_1 c'_{n-1} + c_2 c'_{n-2} + \dots + c_n c'_0 &= 0 \quad (n = 1, 2, 3, \dots) \end{aligned}$$

is called the *inverse* of the sequence $[c_n]$. It is clear that the generating functions $c(t)$, $c'(t)$ of the two sequences are reciprocals of each other. Assume that $k_0 \neq 0$, and consider the expression

$$A(z^\pi) = \frac{k_0 + k_\pi z^\pi + k_{2\pi} z^{2\pi} + \dots}{1 + \frac{\lambda^\pi}{\pi!} z^\pi + \frac{\lambda^{2\pi}}{(2\pi)!} z^{2\pi} + \dots} = u_0 + u_\pi z^\pi + u_{2\pi} z^{2\pi} + \dots$$

On clearing of fractions, we have the following set of equations to determine $u_0, u_\pi, u_{2\pi}, \dots$,

$$\begin{aligned} k_0 &= u_0, \\ k_\pi &= \frac{u_0 \lambda^\pi}{\pi!} + u_\pi, \\ k_{2\pi} &= \frac{u_0 \lambda^{2\pi}}{(2\pi)!} + \frac{u_\pi \lambda^\pi}{\pi!} + u_{2\pi}, \\ &\dots \end{aligned}$$

Since these equations become identical with the equations (12) on replacing u by α we have proved

THEOREM 7. *The generating function of the sequence $\alpha_0, \alpha_\pi, \alpha_{2\pi}, \dots$ in Theorem 5 is given by*

$$A(z^\pi) = \frac{K(z^\pi)}{H(z^\pi)}$$

where

$$K(z^\pi) = \sum_{n=0}^{\infty} k_{n\pi} z^{n\pi},$$

$$H(z^\pi) = \sum_{n=0}^{\infty} \frac{1}{(n\pi)!} \lambda^{n\pi} z^{n\pi} = \frac{1}{\pi} \{e^{a\lambda z} + e^{a^2\lambda z} + \dots + e^{a^\pi\lambda z}\}.$$

The reciprocal of $H(z^\pi)$ generates the sequence inverse to $\left[\frac{\lambda^{n\pi}}{(n\pi)!}\right]$ since by (8)

$$h_{n\pi} = \frac{\lambda^{n\pi}}{(n\pi)!}$$

we write

$$(13) \quad \frac{1}{H(z^\pi)} = H'(z^\pi) = h'_0 + h'_\pi z^\pi + h'_{2\pi} z^{2\pi} + \dots$$

If $\pi = 2$, $a = b = -1$, the case Nielsen has studied*,

$$H'(z^2) = \operatorname{sech} \frac{z}{2} = E_0 + \frac{E_2 z^2}{2^2 2!} + \frac{E_4 z^4}{2^4 4!} + \dots$$

where E_0, E_2, E_4, \dots are Euler's numbers.

Moreover it is clear that

$$(14) \quad \alpha_{n\pi} = k_{n\pi} h'_0 + k_{(n-1)\pi} h'_\pi + \dots + k_0 h'_{n\pi}.$$

Let $[H''_n(x), h''_n]$ denote the cyclic sequence defined by

$$H''_n(x) = \frac{h'_0(x+\lambda)^n}{n!} + \frac{h'_\pi(x+\lambda)^{n-\pi}}{(n-\pi)!} + \frac{h'_{2\pi}(x+\lambda)^{n-2\pi}}{(n-2\pi)!} + \dots \\ \dots + \frac{h'_{r\pi}(x+\lambda)^{n-r\pi}}{(n-r\pi)!},$$

where as usual $r\pi \leq n \leq (r+1)\pi$. Then by (12), if $[K_n(x), k_n]$ is any cyclic sequence

$$k_n = \sum_{s=0}^r \frac{\alpha_{s\pi} \lambda^{n-s\pi}}{(n-s\pi)!} = \sum_{t=0}^r \sum_{s=t}^r \frac{k_{t\pi} h'_{(s-t)\pi} \lambda^{n-s\pi}}{(n-s\pi)!}$$

on substituting from (14) for $\alpha_{s\pi}$ and changing the order of summation. Now

$$\sum_{s=t}^{s=r} \frac{h'_{(s-t)\pi} \lambda^{n-s\pi}}{(n-s\pi)!} = \sum_{u=0}^{u=r-t} \frac{h'_{u\pi} \lambda^{n-t\pi-u\pi}}{(n-t\pi-u\pi)!} = H''_{n-t\pi}(0) = h''_{n-t\pi}.$$

Furthermore if

$$H'(t^\pi) e^{tx} = H'_0(x) + H'_1(x)t + H'_2(x)t^2 + \dots$$

* Nielsen, l. c., chapter VI.

Then

$$H'_n(x) = \frac{h'_0 x^n}{n!} + \frac{h'_\pi x^{n-\pi}}{(n-\pi)!} + \dots + \frac{h'_{r\pi} x^{n-r\pi}}{(n-r\pi)!} = H''_n(x-\lambda).$$

Hence we have proved

THEOREM 8. *If $[K_n(x), k_n]$ is any cyclic sequence of order π*

$$k_n = \sum_{s=0}^r k_{s\pi} H'_{n-s\pi}(\lambda),$$

where $r\pi \leq n < (r+1)\pi$, $\lambda = \frac{b}{a-1}$ and the generating function of the polynomials $H'_n(x)$ is

$$\pi e^{tx} [e^{at} + e^{a^2 t} + \dots + e^{a^{\pi} t}]^{-1}.$$

CALIFORNIA INSTITUTE, PASADENA, CALIFORNIA.

October 29, 1928.