

1 **Investigating behavior of the potato psyllid *Bactericera cockerelli* (Šulc) (Hemiptera:**
2 **Triozidae) on three potato genotypes with putative resistance to “*Candidatus Liberibacter***
3 ***solacearum*”**

4 Austin N. Fife^{1,2,5}, Arash Rashed³, Richard G. Novy⁴, and Erik J. Wenninger¹

5 **Abstract** The potato/tomato psyllid *Bactericera cockerelli* (Šulc) (Hemiptera: Triozidae)
6 transmits “*Candidatus Liberibacter solanacearum*” (Lso), the causal agent of zebra chip
7 disease (ZC) in potato. ZC creates large economic losses when disease incidence is high. No
8 commercial potato variety has been found resistant to the pathogen or the disease
9 symptoms. We observed host acceptance behaviors using no-choice assays on breeding
10 clones derived from *Solanum chacoense* Bitter with putative resistance to Lso and/or ZC.
11 We also compared oviposition and egg fertility for psyllids held on these genotypes.
12 Probing frequency and female walking duration were highest on ‘Russet Burbank,’
13 suggesting greater activity on this variety, than on the three resistant genotypes. The
14 number of eggs did not differ among genotypes but declined on all genotypes during the
15 last period of observation (18-20 days after confinement with a male). Egg fertility did not
16 differ among genotypes for the first three observation periods (16-18 days after
17 confinement with a male) but more eggs were fertile on Russet Burbank than on 10LB or
18 3LB during the last observation period (18-20 days after confinement with a male). We
19 discuss the modality of resistance in context of these genotypes.

20 **Key Words** *Bactericera cockerelli*, *Candidatus Liberibacter solanacearum*, behavior, host plant
21 resistance

1 University of Idaho - Kimberly Research & Extension Center, 3806 N 3600 E, Kimberly, ID, 83341, USA

2 Current address: The University of Florida - North Florida Research and Education Center,
155 Research Road, Quincy, FL, 32351, USA

3 University of Idaho, 875 Perimeter Dr, Moscow, ID, 83844, USA

4 United States Department of Agriculture, Agricultural Research Service, 1693 S 2700 W,
Aberdeen, ID, 83210, USA

5 Corresponding author email: afife@ufl.edu

Formatted: Font: 12 pt, Font color: Auto

Deleted:

Deleted:

Deleted: 5

Formatted: Superscript

Formatted: Superscript

Deleted: 26/03/2019¶

Deleted: Z

Commented [WE(1): I prefer to avoid stating explicitly that it is the causal agent since Koch's postulate cannot be verified given that the bacterium has not been cultured. Most likely it is the causal agent, but it is safer to say that it is "associated with ZC.

Deleted: Russet Burbank

Commented [WE(2): Reviewers generally want you to avoid making vague statements like this. It's better to summarize the conclusion rather than saying we discuss it. I feel like we did that with the abstract in your thesis.

Commented [WE(3): Typically you would choose key words that do not show up in your title. Pick terms that would lead someone to your paper when they are doing a key word search when looking for articles. Maybe *Solanum tuberosum*, tomato psyllid

Formatted: Font: 12 pt

Commented [WE(4): Is “THE University of Florida” the phrasing that they insist upon like THE Ohio State University? That is not a thing with UI.

Commented [WE(5): Check that Rich's address is correct.

Formatted: Superscript

Deleted: 2 Corresponding author email: afife@ufl.edu

Deleted: 3

Deleted: The

Deleted: 4

Deleted: The

Deleted: University of Idaho - Aberdeen Research and Extension Center

29 Introduction

30 The potato/tomato psyllid, *Bactericera cockerelli* (Šulc) (Hemiptera: Trioizidae), is a small
31 sternorrhynchan insect pest of solanaceous crops such as potato, tomato, cape gooseberry,
32 tobacco, pepper, eggplant and tamarillo (Knowlton and Thomas 1934, Wallis 1955, Martin
33 2008, 2009, Aguilar et al. 2013). First discovered in Colorado (Šulc 1909), potato psyllids
34 have a history closely tied to potato growing regions in North America to and to potato
35 diseases (Richards and Blood 1973). The geographical distribution of *B. cockerelli* ranges
36 from southern Canada to Central America, throughout the Western United States
37 (Munyanenza et al. 2007, Rehman et al. 2010, Butler and Trumble 2012) and a recent
38 introduction to New Zealand (Martin 2008, Liefing et al. 2009, Teulon et al. 2009).

39 Interest in potato psyllids grew during the 1920s due to the apparent association of this
40 insect with a condition affecting solanaceous plants known as 'psyllid yellows' (Richards
41 1928, Eyer and Crawford 1933, Richards and Blood 1973). More recently, potato psyllids
42 have been identified as vectors of "*Candidatus Liberibacter solanacearum*" (Lso)
43 (Rhizobiaceae: Alphaproteobacteria) (Cicero et al. 2016, Goolsby et al. 2007, Munyanenza et
44 al. 2007, Liefing et al. 2009). Lso is an uncultured gram-negative α -proteobacterium
45 (Liefing et al. 2009) that infects solanaceous plants. Lso is transmitted to the plant's
46 phloem by the psyllid's saliva while feeding (Cooper and Bamberg 2014).

47 Symptoms of Lso infection in potato include stunting, swollen axillary buds, aerial tubers,
48 leaf purpling, chlorosis and reduced yield (Munyanenza et al. 2007, 2008). Infection also
49 alters tuber sugars and phenolics, resulting in brown stripes that char and blacken when
50 fried (Navarre et al. 2009, Alvarado et al. 2012, Buchman et al. 2012). This condition is
51 known as 'zebra chip' disease (ZC) (Hansen et al. 2008, Liefing et al. 2009, Lin et al. 2009,
52 Crosslin et al. 2011). ZC-affected tubers are unmarketable, which results in large economic
53 losses for growers (Rosson et al. 2006, Munyanenza et al. 2007). Yield reduction from Lso
54 infection has ranged from 43% to 93% in some cases (Munyanenza et al. 2008, 2011).

55 Lso and ZC symptoms were first described in 1994 in Mexico (Secor and Rivera-Varas
56 2004, Munyanenza et al. 2009) and first detected in the United States in 2000 (Secor and
57 Rivera-Varas 2004). Lso and ZC were first detected in the Pacific Northwest (PNW) states
58 of Idaho, Washington and Oregon in 2011 (Crosslin et al. 2012, Murphy et al. 2012). Since
59 2011, Lso and ZC continue to threaten potato production in the PNW, increasing
60 production costs for growers (Guenthner et al. 2012, Greenway 2014, Wenninger et al.
61 2017, Greenway and Rondon 2018).

62 Current management of ZC targets the potato psyllid vector, usually relying on multiple
63 applications of insecticides (Guenthner et al. 2012, Greenway 2014, Echegaray and Rondon
64 2017). In 2018, around half of Eastern Idaho growers' insecticide expenditures were
65 related to ZC control (Greenway and Rondon 2018). Chemicals such as abamectin,
66 imidacloprid, spiromesifen, thiamethoxam and dinotefuran (Goolsby, Adamczyk, et al.
67 2007, Vega-Gutiérrez et al. 2008, Gharalari et al. 2009, Guenthner et al. 2012) are
68 commonly used but, some psyllid populations are starting to develop resistance to common
69 neonicotinoids and abamectin (Liu and Trumble 2004, Hernández-Bautista et al. 2013,
70 Prager et al. 2013, Chávez et al. 2015). The difficulty and large expense of psyllid control

Commented [WE(6)]: Shouldn't this document be double spaced? Check instructions for authors

Formatted: Font: 12 pt, Font color: Auto

Formatted: Font: 12 pt, Font color: Auto

Formatted: First Paragraph

Deleted: ¶

Commented [WE(7)]: Never ever ever begin a sentence with an abbreviated genus

Deleted: .

Deleted: 's

Deleted: geographical distribution

Deleted: w

Deleted: Publications regarding the psyllid initially emerged from 1926-1928, due

Deleted: ¶

P

Deleted: recently

Deleted: ;

Deleted: which

Deleted: ebra chip

Deleted: was

Commented [WE(8)]: Eschewing the Oxford comma makes me cringe, but if you are going to do it throughout, then you have to be consistent.

Deleted: Traditional

Deleted: relies

Deleted: pest

Deleted: ¶

89 emphasizes the need for alternative and improved pest management strategies such as
90 host plant resistance.

91 Host plant resistance is a valuable part of integrated pest management (Kogan 1988, Butler
92 and Trumble 2012a, Munyaneza 2012b, Diaz-Montano et al. 2013). Even a small amount of
93 resistance/tolerance of a plant to a vector or its pathogen can reduce damage below action
94 thresholds and reduce pesticide applications (Kennedy et al. 1987). Host plant resistance
95 increases pesticide efficiency and helps to delay insecticide resistance (Gharalari et al.
96 2009). Currently no commercial potato varieties have been found with acceptable
97 resistance to Lso (Munyaneza et al. 2011, Anderson et al. 2012).

98 Potatoes which have been bred with closely related plants such as *Solanum chacoense*
99 Bitter (Rashidi et al. 2017) and *Solanum berthaultii* Hawkes (Butler et al. 2011) have shown
100 less Lso infection and/or ZC symptoms than other genotypes tested. By determining how
101 these genotypes resist or tolerate either Lso or the psyllid vector itself (Kennedy et al.
102 1987, Putten et al. 2001, Butler et al. 2011), we can decide which traits should be bred or
103 cloned into commercial cultivars to develop resistant potato cultivars (Kaloshian 2004,
104 Casteel et al. 2006, 2007).

105 We examined psyllid host acceptance behaviors as well as oviposition and egg fertility on
106 three potato breeding clones derived from *Solanum chacoense*: 'A07781-10LB' ('10LB'),
107 'A07781-3LB', ('3LB') and 'A07781-4LB' ('4LB') (Rashidi et al. 2017). These genotypes
108 exhibit high tolerance and low susceptibility to Lso (Rashidi et al. 2017), possibly due to
109 resistance or tolerance to the psyllid vector. Russet Burbank was used as a susceptible
110 control (Munyaneza et al. 2011). Our results will help clarify potato psyllid interactions on
111 these genotypes, which will help plant breeders to develop Lso-resistant potatoes
112 (Kennedy et al. 1987).

113 Materials and Methods

114 Plant Characteristics and Living Conditions

115 Potato clones were provided by the USDA-ARS, Small Grains and Potato Germplasm
116 Research Unit Aberdeen, ID, USA. We used three sibling clones derived from *Solanum*
117 *chacoense* Bitter with putative tolerance/resistance to Lso: A07781-3LB, A07781-4LB, and
118 A07781-10LB (Rashidi et al. 2017). 'Russet Burbank' was used because it is susceptible to
119 Lso (Munyaneza et al. 2011) and because of its large impact on potato production in the
120 Pacific Northwest (NASS Northwest Regional Field Office 2017). The selected potatoes
121 were grown in a greenhouse maintained between 25-32°C, 32% RH, with a photoperiod of
122 16:8 (L:D). Plants were grown in pots of approximately 8.5 cm length × 8.5 cm width × 9.5
123 cm height, with a soil mixed in ratios of 4:4:4:1 peat moss: compost: coconut coir: perlite.
124 Fertilizer was not used on experimental plants to avoid nitrogen increases which may alter
125 insect feeding behaviors (Pfeiffer and Burts 1983, 1984). We used plants in their vegetative
126 growth stage (growth stage II) (Dwelle et al. 2003).

Commented [WE(9)]: This paragraph needs to be related to psyllids and ZC, otherwise it sort of comes out of left field. Even just moving the last sentence of this paragraph to the start and adding something to the effect that resistance is ultimately the most sustainable disease management approach.

Commented [WE(10)]: Here and elsewhere, don't put a space between the year and the letter

Deleted:

Deleted:

Deleted: '

Deleted: .

Deleted: /

Deleted: to

Deleted: create

Deleted: es

Deleted: and

Commented [WE(11)]: I'm not sure how to reconcile

Deleted: /

Deleted: -

Deleted:

Formatted: Font: 12 pt, Font color: Auto

Formatted: Font: 12 pt, Font color: Auto

139 **Insect Characteristics and Living Conditions**

Formatted: Font: 12 pt, Font color: Auto

140 A Lso-positive potato psyllid colony was reared in the same greenhouse conditions as
141 described above to avoid phenological asynchrony (Hodkinson et al. 2015). Psyllids were
142 allowed free access to both Russet Burbank potatoes and 'Yellow Pear' tomatoes (*Solanum*
143 *lycopersicum* L.). Colony plants were fertilized once weekly with approximately 17 g of
144 24:8:16 NPK fertilizer per gallon of water (MiracleGro All Purpose Plant Food, Scotts
145 Company, Marysville, OH). Plants were replaced as needed.

146 **Lso Detection**

Formatted: Font: 12 pt, Font color: Auto

147 Idaho harbors four haplotypes of the potato psyllid: Northwestern, Western, Central and
148 Southwestern and two haplotypes of Lso: Lso A and Lso B (Dahan et al. 2017, Wenninger et
149 al. 2017). Our lab colony was determined to be comprised of 'Central' psyllids infected with
150 Lso 'B' via the methods described in Swisher and Crosslin (2014 a). A sample of forty
151 psyllids taken from the colony were transferred to individual microcentrifuge tubes filled
152 with 70% ethanol. Lso incidence was tested at the Aberdeen Research and Extension
153 Center (Aberdeen, ID, USA). DNA extraction was based on the methods described by
154 Marzachi et al. (1998). Individual psyllids were ground by a homogenizer (Omni
155 International Inc., Kennesaw, GA), macerating each psyllid for 1 minute at high speed and
156 an additional minute at medium speed in 500 of Cetyl Trimethylammonium Bromide 2%
157 (Alpha Teknova, Inc., Hollister, CA, Cat. No. C2190) (Composition: 2% CTAB, 100mM Tris-
158 HCl, pH 8.0, 20mM EDTA, pH 8.0, 1.4M Sodium Chloride (NaCl). Microcentrifuge tubes were
159 then incubated at 60 for 30 minutes and gently mixed by inversion every 10 minutes while
160 incubating. Tubes were then spun in a centrifuge at 14,000 rpm for 5 minutes and then the
161 supernatant was transferred to clean 2 ml tubes. The supernatant was vortexed for
162 approximately 20 seconds with 500 ml of chloroform:isoamyl alcohol (24:1 v:v) (Sigma-
163 Aldrich, Inc., Atlanta, GA: Catalogue number C0549), then centrifuged at 14,000 rpm for 5-
164 10 minutes at 4. The clean supernatant was transferred to a new tube, then refrigerated
165 isopropanol (Sigma-Aldrich, Inc., Atlanta, GA: Catalogue number I9516) was added at a rate
166 of 2/3 of the volume of the supernatant. The mixture was then refrigerated at -20 for 20-30
167 minutes. DNA was precipitated by centrifuging the mixture for 20 minutes at 14,000 rpm at
168 4, gently pouring off the supernatant and keeping the precipitated DNA pellet. The pellet
169 was washed in 300 of 70% ethanol and centrifuged for 5 mins at 10,000 rpm. The pellet
170 was then dried overnight in a fume hood. Once dry, 30 of nuclease-free water was added.
171 DNA was stored at -20.

172 Extracted DNA samples were then processed using a Sybgreen method. SsoAdvanced
173 Universal SYBR Green Supermix (Biorad, Hercules, CA) was mixed in a CFX Connect Real-
174 Time PCR Detection System (Biorad, Hercules, CA). HLBr (5'-GCG TTA TCC CGT AGA AAA
175 AGG TAG-3') and LsoF (5'-GTC GAG CGC TTA TTT TTA ATA GGA-3') were used as primers
176 (Li et al. 2006, 2009) and 10 of Sybgreen supermix was added to 150 nM of each primer
177 with 1 of DNA template. The program cycle was as follows: one cycle at 98 for 2 mins
178 followed by 40 cycles of 95 for 10 sec and 62 for 20 sec. The melt curve was 65 to 95, with
179 increments of 0.5 sec⁻¹. DNA of a healthy tuber was used as a negative control. DNA of a
180 Lso-infected tuber was used as a positive control and water was used as a no-template
181 control in all tests. pIDTSmart Kan (Synthetic Genomics, SGI-DNA, CA) with a 250 bp region

182 was amplified with the primer HLBr. The plasmid was diluted 10-fold and used with the
183 following dilutions: 1×10^{-2} , 1×10^{-3} , 1×10^{-4} , 1×10^{-6} , 1×10^{-7} , and 1×10^{-8} ng. Pathogen
184 quantity was reported as copy number of Lso, copy numbers were determined using the
185 methods of (Levy et al. 2011).

186 Each psyllid tested positive for Lso, suggesting a 100% rate of infection for the colony.

187 **No-Choice Behavior Assays**

188 No-choice assays were conducted in a climate-controlled room maintained at 26. Assays
189 were conducted on a wire shelving unit which allowed the testing arena to be lit both from
190 above and below. Three Smith-Victor Digilight fixtures (Smith-Victor Corporation, Bartlett,
191 IL) were used with three Azlo (Akces Media LLC dba ALZO Digital, Bethel, CT) full-spectrum
192 CFL bulbs per light fixture (100-240 volts, 60 Hz, color temp 5500K CRI 91, 750 lumens, 15
193 watts). Two lights were placed with their light sources 35 cm above the testing arena and
194 the light was softened with a diffusion material. The remaining light fixture was placed so
195 that its light source was 45 cm below the testing arena and was softened with diffusion
196 material as well. Illuminance was 3600 lx at the surface of the arena (Sekonic L-308DC-U
197 Light Meter, Sekonic Corporation, Tokyo, Japan).

198 The observation arena (*Fig. 1*) was modeled after the design described by Liu et al. (2004),
199 but modified to use leaflets of intact, potted plants like Butler et al. (2011). This permitted
200 us to observe the psyllids with minimal interference to plant physiology and avoided
201 altering plant volatiles or chemical defenses activated by damaging plant tissues (Klingler
202 et al. 2005). A recording arena was formed by sandwiching a panel of glass, a wetted filter
203 paper, a leaf, and a piece of Plastazote polyethylene foam (Zotefoams Inc., Croydon, UK),
204 with a circular opening in the center (28 mm diameter). The arena was held together with
205 two clips. This arena was then suspended by a suction cup held by an adjustable burette
206 clamp. We used leaves from the upper canopy of the plants. The filter paper was discarded
207 between observations. The glass pane and foam were replaced with each new plant and
208 washed and dried at 90 before reuse to remove potential volatile accumulation. Recordings
209 were done with a L3CMOS C-mount USB camera and ToupView recording software
210 (L3CMOS14000KPA, Hangzhou ToupTek Photonics Co., Ltd, Hangzhou, Zhejiang, China).

211 We collected psyllids from the colony by aspiration and transferred them to 8×35 mm
212 glass shell vials. All psyllids were used within 90 minutes from the time of collection.
213 Psyllids were introduced to the arena and recorded for five minutes. Psyllid sex was
214 identified and psyllids were preserved in 95% ethanol for later testing for Lso by PCR. We
215 recorded similar categories as Butler et al. (2011): probing, walking, cleaning, and whether
216 the psyllid was on or off the leaf. Probing behaviors have putative significance with disease
217 transmission and host selection (Prager, Esquivel, et al. 2014 a, 2014 b). Behavior was
218 scored using CowLog3 (Hänninen and Pastell 2009), which recorded incidence and
219 timestamps for the behaviors observed.

220 **Oviposition Assays**

221 Oviposition assays were conducted with greenhouse conditions, plants, and insects as
222 previously described. A female/male pair of teneral psyllids (identified by their green body

Formatted: Font: 12 pt, Font color: Auto

Formatted: Font: 12 pt, Font color: Auto

223 color) was introduced to a plant covered with an insect rearing sleeve (MegaView Science
224 Co., Ltd., Taiwan). Rearing sleeves were supported over the plant using two lengths of
225 galvanized steel wire with a diameter of 1.63 mm. Each wire was curved into a parabolic
226 shape and each end of the wire was inserted into the soil on opposite corners of the plant
227 pot (*Fig. 2*). Plants were blocked by germplasm accession in rows of four and placed inside
228 60 cm length × 60 cm width × 60 cm height mesh-covered PVC-framed cages. Plants were
229 watered on alternating days by soaking pots in 56 cm length × 28 cm width × 6 cm height
230 plastic trays until the soil became saturated (approximately 45 mins). After a period of six
231 to eight days the males were removed from the plants and the female transferred to a new
232 plant of the same accession. The female psyllid was then transferred to a new plant every
233 four days at three intervals. Eggs were counted on each plant after the female was
234 removed. Nymphs were counted four days, eight days and twelve days later to allow time
235 for hatching (Knowlton and Janes 1931). Each nymph was removed as it was counted. The
236 number of nymphs that hatched was considered an indicator of egg fertility. Fertility
237 percentages were calculated as the ratio of nymphs divided by egg counts for each sample.

238 **Statistical Analysis**

239 Statistical analysis was performed using R Version 3.5.1 (R Core Team 2013) Assumptions
240 of normality were investigated with qqplots and Cullen and Frey graphs from the R
241 package fitdistrplus (Delignette-Muller and Dutang 2015). No-choice experiments and egg
242 count data were analyzed using generalized linear mixed modeling techniques (GLMM)
243 (Stroup 2015) from the glmer function (Bates et al. 2015). A Poisson distribution and log
244 link were used to model count data. Egg fertility was modeled with a binomial distribution
245 and log link to account for ratios. Behavioral models had fixed factors of germplasm
246 accession, sex and the interaction of accession × sex. Psyllid replicate was treated as a
247 random factor. The interaction of accession × sex was excluded from the off-leaf model to
248 low occurrences (n = 20 out of 181 observations), which did not allow an interaction to be
249 estimated by the model. Oviposition models had fixed factors of accession, period and
250 accession × period. Psyllid replicate was considered the random factor. Egg fertility was
251 modeled with accession and period as fixed factors and individual psyllids as the random
252 factor. All data were tested with Wald's χ^2 tests, followed by least-squares means with
253 Tukey's adjustments to test for multiple comparisons. Statistical significance was
254 considered at $\alpha = 0.05$.

Formatted: Font: 12 pt, Font color: Auto



255

256 **Fig. 1.** No-choice arena used for behavioral recordings



257

258 **Fig. 2.** Sleeve cage with potato used in oviposition assays

Results

No-Choice Assays

The number of probing events observed was significantly different among genotypes (*Table 1*). Psyllids probed more frequently on Russet Burbank than on A07781-10LB and A07781-3LB, which did not differ between each other (*Table 2*). Probing frequency on A07781-4LB did not differ among the other genotypes. This effect appeared to reflect the trend of more probing by females on Russet Burbank (*Table 2*). However, the genotype \times sex interaction was not significant (*Table 1*). Probing frequency did not was not affected by sex (*Table 2*). Overall, psyllids spent more time engaged in probing behavior than in the other activities recorded (*Tables 2-5*). Probing duration did not differ among genotypes, between sexes or by their interaction (*Table 1*).

The number of walking events differed significantly among genotypes as well as by the interaction of genotype \times sex (*Table 1*). Psyllids walked more on Russet Burbank than 10LB (*Table 3*). Female psyllids on Russet Burbank walked significantly more often than males and females on 10LB and females on 3LB (*Table 3*). The other means did not differ among each other. Walking duration did not differ among genotypes or between sexes, but the interaction term was significant (*Table 1*). Female psyllids walked significantly longer on Russet Burbank than for all other genotype \times sex combinations (*Table 3*).

Cleaning behaviors generally were uncommon and of short duration. The frequencies and durations of cleaning behaviors were not significantly different among genotypes, between sexes, or by their interaction (*Table 1, Table 4*).

Off-leaf behaviors also tended to occur rarely. Frequency of off-leaf behaviors did not differ among genotypes, between sexes or by their interaction (*Table 1*). However, the duration of off-leaf behaviors differed significantly among genotypes (*Table 1*). Psyllids spent more time off-leaf in the 3LB treatment relative to the 4LB and Russet Burbank treatments. Time spent off-leaf in the 10LB treatment did not differ among the other genotypes (*Table 5*). Off-leaf duration did not differ by sex (*Table 1*). The interaction between genotype and sex could not be analyzed due to the low number psyllids observed leaving the leaf ($n = 20$ out of 181).

Oviposition Assays

Neither the number of eggs nor percent viable eggs differed significantly among genotypes (*Table 6*). However, both the number of eggs and egg fertility were significantly different by period and the interaction of genotype \times period (*Table 6*). For oviposition, this interaction effect was an artifact of calculating multiple comparisons of different genotypes across observation periods. There were no significant differences among genotypes within a given period (*Table 7*). For egg fertility during the last period, there were significantly more fertile eggs on Russet Burbank than 10LB or 3LB and there were significantly more eggs on 4LB than 10LB (*Table 7*). There were no significant differences among genotypes within periods 1-3 (*Table 7*). Overall oviposition (with genotype pooled) was significantly lower during period 4 than for the first (*Table 7*). Similarly, egg fertility (with genotype pooled)

Formatted: Font: 12 pt, Font color: Auto

Formatted: Font: 12 pt, Font color: Auto

Formatted: Font: 12 pt, Font color: Auto

299 tended to decline during the last observation period for all genotypes except for Russet
300 Burbank (Table 7).

301 Discussion

302 It is difficult to separate the mechanisms of host plant resistance or tolerance and how
303 these correlate with psyllid host acceptance (Diaz-Montano et al. 2006, Butler et al. 2011).
304 Our visual observations of settling behavior lack the precision of electrical penetration
305 recordings used in similar studies (Butler et al. 2012, Sandanayaka et al. 2014, Mustafa et
306 al. 2015), but requires less expensive equipment. Our results are like other investigations
307 of putatively resistant potato genotypes. Our study found more probing and walking on
308 Russet Burbank than on the putatively resistant genotypes, which is consistent with results
309 reported by Butler et al. (2011) and Prager et al. [(2014); b]. However, in contrast to Butler
310 et al. (2011), we found cleaning and leaf-leaving behaviors to be rare. Russet Burbank
311 received more probes than two other genotypes, but the psyllids still probed the other
312 genotypes, often for long periods. Sandanayaka et al. (2014) and Mustafa et al [(2015); b]
313 both suggest that it takes *B. cockerelli* approximately two hours to access the phloem and
314 acquire Lso. This suggests that very long recordings may be necessary to determine when
315 probing becomes true feeding. Minimal overnight recordings revealed little activity besides
316 apparent feeding on the genotype where they were placed (ANF, unpublished data). In
317 addition, psyllids rarely abandoned the plants where they began to probe. A single psyllid
318 is enough to transmit Lso and the disease progresses independently of bacterial titer
319 (Buchman et al. 2011, a; Rashed et al. 2012). Therefore, it is unlikely that we were
320 observing phloem feeding which would result in pathogen transmission within the span of
321 our short observation periods. These factors underscore that psyllid probing behavior
322 would have to be nearly eliminated to truly reduce the risk of Lso transmission. We found
323 no evidence for such reductions in probing behavior on these genotypes.

324 Studies on the Asian citrus psyllid, *Diaphorina citri* Kuwayama (Hemiptera: Liviidae), a
325 vector of a similar liberibacter pathogen (Teixeira et al. 2005) have examined how host
326 plant volatiles can alter psyllid behaviors (Wenninger et al. 2009, Davidson et al. 2014).
327 Plant volatiles can induce probing in combination to visual and chemical cues from host
328 plants (Patt et al. 2011).

329 It is possible that Lso infection has altered *B. cockerelli*'s attraction to leaf volatiles (Mayer
330 et al. 2008), and their settling behavior as well (Mas et al. 2014). Lso infection can increase
331 psyllid preferences for undamaged, uninfected hosts for oviposition and settling (Davis et
332 al. 2012) – a behavior which has been seen in other insect-plant-vector relationships (Cao
333 et al. 2016, Eigenbrode et al. 2018). It may be that this encouraged greater acceptance of
334 genotypes which would be rejected by an uninfected psyllid. A high percentage of the
335 psyllids in our colony were infected and our plants were all uninfected, so psyllid infection
336 does not entirely explain the patterns we have seen. Infection status also fails to explain the
337 minor trend we saw between male and female probing on Russet Burbank.

338 Another explanation for differences between genotypes is that the female psyllids are more
339 influenced by familiar cues while selecting host plants for oviposition or feeding (Prager et

Formatted: Font: 12 pt, Font color: Auto

340 al. 2014). Russet Burbank was one of the plants used to rear our colonies, so it is possible
341 that the volatiles from this variety were more stimulating for female psyllids. Further
342 studies into potato psyllid's attraction to plant volatiles while infected can help clarify if
343 these possible explanations correlate with host plant acceptance.

344 Although leaf-leaving duration differed significantly among genotypes, the incidence and
345 duration of leaf-leaving behaviors was very small and probably not biologically significant.
346 It is also important to note that leaf-leaving was defined in the context of leaving the leaf in
347 our small observation arena. On a plant in the field there is a much larger surface area for a
348 psyllid to explore, so the leaf-leaving events might represent questing behavior rather than
349 host rejection. It also is possible that the duration between a psyllid's initial encounter and
350 settling behaviors or eventual plant rejection is longer than the time we allotted for
351 recording.

352 Contrary to previously published studies (Butler et al. 2011, Diaz-Montano et al. 2013,
353 Cooper and Bamberg 2014, Rubio-Covarrubias et al. 2017) our study showed similar
354 oviposition rates among genotypes, consistent with results reported by (Prager et al.
355 2017). Other studies have found psyllids will oviposit on a variety of hosts (Diaz-Montano
356 et al. 2013, Thinakaran et al. 2015), even when it is not beneficial for their survival (Prager,
357 Lewis, et al. (2014) b). Psyllids oviposited on every type of potato offered, like observations
358 by Prager et al. (2017), giving little evidence of antixenosis.

359 We selected the number of days for our observations to correlate with the periods of
360 maximum oviposition reported in the life history tables of Abdullah (Knowlton and Janes
361 1931, 2008) and Yang et al. (2010, 2013). Therefore, it was surprising to see the large
362 reduction of egg fertility for some psyllids in period four (18-24 days). Fertility declined on
363 the resistant genotypes as opposed to the Russet Burbank variety, which suggests that
364 these genotypes may have antibiotic effects over time. Over the course of a growing season,
365 these reductions may have a cumulative effect on psyllid populations. Longer observation
366 periods could help to better quantify these effects.

367 It is possible that Lso infection status played a role in the fertility observed: Lso has been
368 reported to negatively impact female fertility (2012 a, Nachappa, Shapiro, et al. 2012, 2014,
369 Yao et al. 2016, Frias et al. 2018). The antibiotic effects we observed may have different
370 effects on uninfected psyllids.

371 We saw a large degree of variability in fertility for psyllids on all genotypes. We only
372 permitted male access to the female psyllids during the initial period to increase female
373 longevity by preventing possible harassment (Abdullah 2008, Wenninger and Hall 2008,
374 Arnqvist 2013). Abdullah (2008, and Yang and Liu 2009, and Yang et al. 2013) all kept
375 female and male psyllids together to freely mate for the duration their observations, which
376 may explain why they observed greater fertility than we did. It is possible that potato
377 psyllids may require multiple mates and/or multiple matings over time to maintain egg
378 fertility (Wenninger and Hall 2008, Arnqvist 2013). Knowlton and Janes (1931) reported
379 (with a limited number of observations) reductions in egg fertility over time after a single
380 mating. There also may be some variability in female reproductive output created by the
381 physiological interactions of male spermatophores, female spermathecae and/or

382 spermatodose (Marchini et al. 2011), which all influence how long females are able to
383 remain fertile (Qazi and Hogdal 2010, Schnakenberg et al. 2011, Wolfner 2011, Abe and
384 Kamimura 2015).

385 In conclusion, we found little evidence of antixenosis or antibiosis with respect to settling
386 behavior, but we saw a reduction in egg fertility on the resistant genotypes 18-24 days
387 after mating. Taken together, these results suggest that the modality of resistance to Lso for
388 the A07781 genotypes (Rashidi et al. 2017) is not likely related to psyllid settling
389 behaviors, but that reduced Lso symptoms may be due to resistance to the pathogen itself,
390 and not the psyllid vector. Further work will be required to clarify the modality of
391 resistance to Lso in the A07781 genotypes.

392
|

Table 1. Wald's χ^2 tests comparing psyllid behaviors between sexes and among four genotypes: A07781-10LB, A07781-3LB, A07781-4LB and Russet Burbank

Behavior	Factors	Incidence		Duration	
		df	χ^2	Pr > χ^2	Pr > χ^2
Probing	Genotype	3	27.46	0.000*	0.473
	Sex	1	3.24	0.072	0.959
	Genotype \times Sex	3	6.49	0.090	0.192
Walking	Genotype	3	16.17	0.001*	0.199
	Sex	1	1.65	0.200	0.850
	Genotype \times Sex	3	11.13	0.011*	0.013*
Cleaning	Genotype	3	5.98	0.113	0.525
	Sex	1	0.45	0.503	0.490
	Genotype \times Sex	3	0.33	0.955	0.993
Off-Leaf [†]	Genotype	3	1.15	0.765	0.023*
	Sex	1	0.71	0.401	0.832
	Genotype \times Sex	3	—	—	—

* indicates values which are significant at $P > 0.05$

[†] The interaction genotype \times sex was unable to be analyzed due to the low number of psyllids which left the leaf ($n = 20$ out of 181)

Commented [WE(12)]: The tables should be moved after the References

Deleted: :

Deleted: χ^2

Formatted: Superscript

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Centered

Formatted Table

Commented [WE(13)]: Please fix all the other weird and wonky equation formatting that is going on throughout these tables

Deleted: $\text{Pr} > \chi^2$

Formatted: Centered

Deleted: χ^2

Deleted: $\text{Pr} > \chi^2$

Deleted: χ^2

Formatted: Centered

Formatted: Centered

Formatted: Centered

Deleted: ¶

Formatted: Centered

Formatted: Centered

Formatted: Centered

Deleted: ¶

Formatted: Centered

Formatted: Centered

Formatted: Centered

Deleted: ¶

Formatted: Centered

Formatted: Centered

Formatted: Centered

Commented [WE(14)]: This is really not necessary since you report the exact P-values in the table. Please delete.

Commented [WE(15)]: Is this the symbol they want you to use for table notes? Usually for this journal you would use a, b, c, etc. I don't know if they have a hard-and-fast ...

408
409
410

$$\begin{array}{l} 411 \\ 412 \\ 413 \end{array}$$

414

446
447
448

449
450
451
452

449
450
451
452

Table 4. Least-square mean \pm SEM incidence and duration of potato psyllid cleaning
behaviors recorded during 300-s no-choice tests on four different genotypes: A07781-
10LB, A07781-3LB, A07781-4LB and Russet Burbank

Genotype	Sex	N	Incidence	Duration (s)
10LB	Female	21	0.34 ± 0.15	0.008 ± 0.017
	Male	25	0.33 ± 0.13	0.023 ± 0.048
3LB	Female	27	0.13 ± 0.07	0.002 ± 0.003
	Male	21	0.20 ± 0.10	0.003 ± 0.005
4LB	Female	25	0.20 ± 0.10	0.002 ± 0.003
	Male	18	0.26 ± 0.13	0.008 ± 0.018
Russet Burbank	Female	26	0.09 ± 0.05	0.001 ± 0.001
	Male	18	0.13 ± 0.08	0.001 ± 0.002

480

494
495
496

497
498
499

500
501

\$02

Deleted: :	
Deleted: P	
Formatted	...
Formatted	...
Formatted	...
Formatted Table	...
Formatted	...
Formatted	...
Formatted	...
Formatted	...
Formatted	...
Formatted	...
Formatted	...
Formatted	...
Formatted	...
Formatted	...
Deleted: ¶	...
Formatted Table	...
Formatted	...
Formatted	...
Formatted	...
Formatted	...
Formatted	...
Deleted: ¶	...
Formatted Table	...
Formatted	...
Formatted	...
Formatted	...
Formatted	...
Formatted	...
Formatted	...
Deleted: ¶	...
Formatted Table	...
Formatted	...
Formatted	...
Formatted	...
Formatted	...
Formatted	...
Formatted	...
Formatted	...
Commented [WE16]: See comment above about this	...
Deleted: Least-squares means. Differences among	...
Formatted	...

518 Table 7. Mean \pm SEM, (A) total eggs laid and (B) egg fertility of psyllids on four different genotypes.

519 A. Total Eggs

Genotype	N	Period 1 ^a	Period 2	Period 3	Period 4
10LB	20	6.3 \pm 1.5	7.0 \pm 1.7	9.4 \pm 2.3	3.8 \pm 1.0
3LB	13	4.8 \pm 1.4	9.5 \pm 2.8	9.1 \pm 2.7	4.3 \pm 1.3
4LB	19	8.4 \pm 2.0	10.5 \pm 2.6	8.0 \pm 2.0	6.9 \pm 1.8
Russet Burbank	14	5.8 \pm 1.7	7.6 \pm 2.2	7.0 \pm 2.0	6.6 \pm 1.9
Overall	66	9.5 \pm 1.6	12.5 \pm 1.8	12.5 \pm 2.0	8.2 \pm 1.5

520 B. Percent Fertility

Genotype	N	Period 1	Period 2	Period 3	Period 4
10LB	20	68.8 \pm 9.2	59.5 \pm 10.9	61.8 \pm 10.7	3.2 \pm 2.0 ^a
3LB	13	65.9 \pm 12.8	61.0 \pm 12.6	55.7 \pm 13.3	11.9 \pm 6.8 ^{ab}
4LB	19	62.3 \pm 10.5	64.1 \pm 10.1	49.6 \pm 12.2	29.2 \pm 10.4 ^{bc}
Russet Burbank	14	47.0 \pm 13.0	50.9 \pm 12.7	63.9 \pm 11.9	70.1 \pm 10.9 ^c
Overall	66	66.8 \pm 4.2 ^A	68.2 \pm 4.0 ^{AB}	66.0 \pm 5.5 ^{AB}	43.8 \pm 6.2 ^B

519 Means for individual genotypes within a time period that share a letter or overall means within a row that share a letter are
520 not significantly different ($P > 0.05$).

521 ^aPeriod 1 comprised six to eight days, during which a mating pair of psyllids was held on a caged plant. At the end of the first
522 time period, the male was removed and the remaining female was held on a new plant of the same genotype over three
523 successive four-day time periods (Periods 2-4).

Deleted: (

Deleted:)...(A) total eggs laid totals

Moved down [2]: A mating pair of psyllids was placed

Deleted: N

Formatted

Deleted: Period 1

Formatted

Deleted: Period 2

Formatted

Deleted: Period 3

Formatted

Deleted: Period 4

Formatted

Formatted

Formatted Table

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Deleted: ¶

Formatted

Formatted

Formatted

Formatted

Formatted

Deleted: Percent Fertile

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

References Cited

- Abdullah, N. M. H. 2008.** Life history of the potato psyllid *Bactericera cockerelli* (Homoptera: Psyllidae) in controlled environments agriculture in Arizona. *Afr. J. Agric. Res.* 3: 60–67.
- Abe, J., and Y. Kamimura. 2015.** Sperm economy between female mating frequency and male ejaculate allocation. *Am. Nat.* 185: 406–416.
- Aguilar, E., V. G. Sengoda, B. Bextine, K. F. McCue, and J. E. Munyaneza. 2013.** First report of "*Candidatus Liberibacter solanacearum*" on tobacco in Honduras. *Plant Dis.* 97: 1376–1376.
- Alvarado, V. Y., D. Odokonyero, O. Duncan, T. E. Mirkov, and H. B. Scholthof. 2012.** Molecular and physiological properties associated with zebra complex disease in potatoes and its relation with *Candidatus Liberibacter* contents in psyllid vectors. *PLoS ONE.* 7: e37345.
- Anderson, J. A. D., G. P. Walker, P. A. Alspach, M. Jeram, and P. J. Wright. 2012.** Assessment of susceptibility to zebra chip and *Bactericera cockerelli* of selected potato cultivars under different insecticide regimes in New Zealand. *Am. J. Potato Res.* 90: 58–65.
- Arnqvist, G. 2013.** Sexual conflict. Princeton University Press.
- Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015.** Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* 67.
- Buchman, J. L., T. W. Fisher, V. G. Sengoda, and J. E. Munyaneza. 2012.** Zebra chip progression: From inoculation of potato plants with *Liberibacter* to development of disease symptoms in tubers. *Am. J. Potato Res.* 89: 159–168.
- Buchman, J. L., V. G. Sengoda, and J. E. Munyaneza. 2011.** Vector transmission efficiency of *Liberibacter* by *Bactericera cockerelli* (Hemiptera: Triozidae) in zebra chip potato disease: Effects of psyllid life stage and inoculation access period. *J. Econ. Entomol.* 104: 1486–1495.
- Butler, C. D., B. Gonzalez, K. L. Manjunath, R. F. Lee, R. G. Novy, J. C. Miller, and J. T. Trumble. 2011.** Behavioral responses of adult potato psyllid, *Bactericera cockerelli* (Hemiptera: Triozidae), to potato germplasm and transmission of *Candidatus Liberibacter psyllaeus*. *Crop Prot.* 30: 1233–1238.
- Butler, C. D., and J. T. Trumble. 2012.** The potato psyllid, *Bactericera cockerelli* (Sulc) (Hemiptera: Triozidae): Life history, relationship to plant diseases, and management strategies. *Terrestrial arthropod reviews.* 5: 87–111.
- Butler, C. D., G. P. Walker, and J. T. Trumble. 2012.** Feeding disruption of potato psyllid, *Bactericera cockerelli*, by imidacloprid as measured by electrical penetration graphs. *Entomol. Exp. Appl.* 142: 247–257.

Commented [WE(19)]: No acknowledgments section????

Formatted: Indent: Left: 0", Hanging: 0.5"

- 631 **Cao, H., H. Liu, Z. Zhang, and T. Liu. 2016.** The green peach aphid *Myzus persicae* perform
632 better on pre-infested chinese cabbage *Brassica pekinensis* by enhancing host plant
633 nutritional quality. *Sci. Rep.* 6.
- 634 **Casteel, C. L., L. L. Walling, and T. D. Paine. 2006.** Behavior and biology of the tomato
635 psyllid, *Bactericera cockerelli*, in response to the mi-1.2 gene. *Entomol. Exp. Appl.*
636 121: 67–72.
- 637 **Casteel, C. L., L. L. Walling, and T. D. Paine. 2007.** Effect of mi-1.2 gene in natal host
638 plants on behavior and biology of the tomato psyllid *Bactericera cockerelli* (Sulc)
639 (Hemiptera: Psyllidae). *J. Entomol. Sci.* 42: 155–162.
- 640 **Chávez, E. C., O. H. Bautista, J. L. Flores, L. A. Uribe, and Y. M. O. Fuentes. 2015.**
641 Insecticide-resistance ratios of three populations of *Bactericera cockerelli*
642 (Hemiptera: Psylloidea: Triozidae) in regions of northern Mexico. *Fla. Entomol.* 98:
643 950–953.
- 644 **Cicero, J. M., T. W. Fisher, and J. K. Brown. 2016.** Localization of “*Candidatus Liberibacter*
645 *solanacearum*” and evidence for surface appendages in the potato psyllid vector.
646 *Phytopathology.* 106: 142–154.
- 647 **Cooper, W. R., and J. B. Bamberg. 2014.** Variation in *Bactericera cockerelli* (Hemiptera:
648 Triozidae) oviposition, survival, and development on *Solanum bulbocastanum*
649 germplasm. *Am. J. Potato Res.* 91: 532–537.
- 650 **Crosslin, J. M., H. Lin, and J. E. Munyaneza. 2011.** Detection of “*Candidatus Liberibacter*
651 *solanacearum*” in the potato psyllid, *Bactericera cockerelli* (Sulc), by conventional
652 and real-time PCR. *Southwest. Entomol.* 36: 125–135.
- 653 **Crosslin, J. M., N. Olsen, and P. Nolte. 2012.** First report of zebra chip disease and
654 *Candidatus Liberibacter solanacearum* on potatoes in Idaho. *Plant Dis.* 96: 453–453.
- 655 **Dahan, J., E. J. Wenninger, B. Thompson, S. Eid, N. Olsen, and A. V. Karasev. 2017.**
656 Relative abundance of potato psyllid haplotypes in southern Idaho potato fields
657 during 2012 to 2015, and incidence of “*Candidatus Liberibacter solanacearum*”
658 causing zebra chip disease. *Plant Dis.* 101: 822–829.
- 659 **Davidson, M. M., R. C. Butler, N. M. Taylor, M. C. Nielsen, C. E. Sansom, and N. B. Perry.**
660 **2014.** A volatile compound, 2-undecanone, increases walking, but not flying, tomato
661 potato psyllid movement toward an odour source. *New Zealand plant protection.*
662 67: 184–190.
- 663 **Davis, T. S., D. R. Horton, J. E. Munyaneza, and P. J. Landolt. 2012.** Experimental
664 infection of plants with an herbivore-associated bacterial endosymbiont influences
665 herbivore host selection behavior. *PLoS ONE.* 7: e49330.
- 666 **Delignette-Muller, M. L., and C. Dutang. 2015.** fitdistrplus: An R package for fitting
667 distributions. *J. Stat. Softw.* 64.

- 668 **Diaz-Montano, J., J. C. Reese, W. T. Schapaugh, and L. R. Campbell. 2006.**
 669 Characterization of antibiosis and antixenosis to the soybean aphid (Hemiptera:
 670 Aphididae) in several soybean genotypes. *J. Econ. Entomol.* 99: 1884–1889.
- 671 **Diaz-Montano, J., B. G. Vindiola, N. Drew, R. G. Novy, J. C. Miller, and J. T. Trumble.**
 672 **2013.** Resistance of selected potato genotypes to the potato psyllid (Hemiptera:
 673 Trioziidae). *Am. J. Potato Res.* 91: 363–367.
- 674 **Dwelle, R. B., J. M. Alvarez, P. Bain, C. R. Baird, E. J. Bechinski, W. H. Bohl, D. L. Corsini,**
 675 **C. V. Eberlein, L. L. Ewing, B. F. Finnigan, B. D. Geary, J. F. Guenther, S. L. Hafez,**
 676 **P. J. S. Hutchinson, W. B. Jones, B. A. King, G. E. Kleinkopf, J. S. Miller, P. Nolte, R.**
 677 **Novy, N. Olsen, S. Palanisamy, P. E. Patterson, L. E. Sandvol, R. L. Stoltz, D. T.**
 678 **Westermann, and J. C. Whitmore. 2003.** Potato production systems, pp. 12–14. *In*.
 679 The University of Idaho agricultural communications.
- 680 **Echegaray, E. R., and S. I. Rondon. 2017.** Incidence of *Bactericera cockerelli* (Hemiptera:
 681 Trioziidae) under different pesticide regimes in the lower Columbia Basin. *J. Econ.*
 682 *Entomol.* 110: 1639–1647.
- 683 **Eigenbrode, S. D., N. A. Bosque-Pérez, and T. S. Davis. 2018.** Insect-borne plant
 684 pathogens and their vectors: Ecology, evolution, and complex interactions. *Annu.*
 685 *Rev. Entomol.* 63: 169–191.
- 686 **Eyer, J. R., and R. F. Crawford. 1933.** Observations on the feeding habits of the potato
 687 psyllid (*Paratrioza cockerelli* Sulc.) and the pathological history of the "psyllid
 688 yellows" which it produces. *J. Econ. Entomol.* 26: 846–850.
- 689 **Frias, A. A. T., F. Ibanez, A. Mendoza, W. M. de Carvalho Nunes, and C. Tamborindeguy.**
 690 **2018.** Effects of "*Candidatus Liberibacter solanacearum*" (haplotype b) on
 691 *Bactericera cockerelli* fitness and vitellogenesis. *Insect Sci.*
- 692 **Gharalari, A. H., C. Nansen, D. S. Lawson, J. Gilley, J. E. Munyaneza, and K. Vaughn.**
 693 **2009.** Knockdown mortality, repellency, and residual effects of insecticides for
 694 control of adult *Bactericera cockerelli* (Hemiptera: Psyllidae). *J. Econ. Entomol.* 102:
 695 1032–1038.
- 696 **Goolsby, J. A., J. Adamczyk, B. Bextine, D. Lin, J. E. Munyaneza, and G. Bester. 2007.**
 697 Development of an IPM program for management of the potato psyllid to reduce
 698 incidence of zebra chip disorder in potatoes. *Subtropical Plant Science.* 59: 85–94.
- 699 **Goolsby, J. A., B. Bextine, J. E. Munyaneza, M. Setamou, J. Adamczyk, and G. Bester.**
 700 **2007.** Seasonal abundance of sharpshooters, leafhoppers, and psyllids associated
 701 with potatoes affected by zebra chip disorder. *Subtropical Plant Science.* 59: 15–23.
- 702 **Greenway, G. 2014.** Economic impact of zebra chip control costs on grower returns in
 703 seven US states. *Am. J. Potato Res.* 91: 714–719.
- 704 **Greenway, G. A., and S. Rondon. 2018.** Economic impacts of zebra chip in Idaho, Oregon,
 705 and Washington. *Am. J. Potato Res.*

- 706 **Guenthner, J., J. Goolsby, and G. Greenway. 2012.** Use and cost of insecticides to control
707 potato psyllids and zebra chip on potatoes. *Southwest. Entomol.* 37: 263–270.
- 708 **Hansen, A. K., J. T. Trumble, R. Stouthamer, and T. D. Paine. 2008.** A new
709 huanglongbing species, "*Candidatus Liberibacter psyllae*," found to infect
710 tomato and potato, is vectored by the psyllid (*Bactericera cockerelli*) (Sulc). *Appl.*
711 *Environ. Microbiol.* 74: 5862–5865.
- 712 **Hänninen, L., and M. Pastell. 2009.** CowLog: Open-source software for coding behaviors
713 from digital video. *Behav. Res. Methods.* 41: 472–476.
- 714 **Hernández-Bautista, O., E. Cerna-Chávez, J. Landeros-Flores, Y. Ochoa-Fuentes, J.**
715 **Chacón-Hernández, and S. Castillo-Arriaga. 2013.** Resistance proportion of
716 *Bactericera cockerelli* (Sulc) in regions from Villa de Arista, San Luis Potosí and
717 Saltillo, Coahuila. *Entomología mexicana.*
- 718 **Hodkinson, I. D., C. V. Achterberg, M. Ahola, M. Barták, V. Behan-Pelletier, J. M. Bird, K.**
719 **Bøg, F. Brodo, P. N. Buhl, C. Dahl, R. H. L. Disney, K. Dittmar, A. Fjellberg, Ø.**
720 **Gammelmo, M. Forshage, R. Gerecke, C. A. Gertsson, M. M. L. Haastriter, J. P.**
721 **Haenni, O. E. Heie, J. M. Heraty, C. Hodgson, D. Horsfield, J. T. Huber, M. Jaschoff,**
722 **F. Jensen, K. A. Johanson, R. Jussila, O. Karsholt, E. Krzeminska, V. I. Lantsov, P.**
723 **Láska, C. Lindegaard, L. Lyneborg, O. Makarova, Y. M. Marusik, W. N. Mathis, L.**
724 **Mazánek, V. Michelsen, T. Munk, W. L. Murphy, S. A. Nielsen, T. R. Nielsen, J. S.**
725 **Noyes, P. Oosterbroek, A. L. Ozerov, T. Pape, J. D. Pinto, M. Pollet, E. Rindal, J.**
726 **Roháček, T. J. Simonsen, V. S. Smith, G. Söli, J. Stary, R. Z. Strassen, B. W.**
727 **Svensson, L. Vilhelmsen, P. Vilkamaa, M. Wilson, and T. Zatwarnicki. 2015.**
728 Psyllidae (jumping plant-lice, psyllids), p. 113. *In* The Greenland Entomofauna: An
729 Identification Manual of Insects, Spiders and Their Allies (Fauna Entomologica
730 Scandinavica). Brill Academic Pub.
- 731 **Kaloshian, I. 2004.** Gene-for-gene disease resistance: Bridging insect pest and pathogen
732 defense. *J. Chem. Ecol.* 30: 2419–2438.
- 733 **Kennedy, G. G., F. Gould, O. M. B. Deponti, and R. E. Stinner. 1987.** Ecological,
734 agricultural, genetic, and commercial considerations in the deployment of insect-
735 resistant germplasm. *Environ. Entomol.* 16: 327–338.
- 736 **Klingler, J., R. Creasy, L. Gao, R. M. Nair, A. S. Calix, H. S. Jacob, O. R. Edwards, and K. B.**
737 **Singh. 2005.** Aphid resistance in *Medicago truncatula* involves antixenosis and
738 phloem-specific, inducible antibiosis, and maps to a single locus flanked by NBS-LRR
739 resistance gene analogs. *Plant Physiol.* 137: 1445–1455.
- 740 **Knowlton, G. F., and M. J. Janes. 1931.** Studies on the biology of *Paratrioza cockerelli*
741 (Sulc). *Ann. Entomol. Soc. Am.* 24: 283–292.
- 742 **Knowlton, G. F., and W. L. Thomas. 1934.** Host plants of the potato psyllid. *J. Econ.*
743 *Entomol.* 27: 547–549.

- 744 **Kogan, M. 1988.** Integrated pest management theory and practice. Entomol. Exp. Appl. 49:
745 59–70.
- 746 **Levy, J., A. Ravindran, D. Gross, C. Tamborindeguy, and E. Pierson. 2011.** Translocation
747 of "*Candidatus Liberibacter solanacearum*", the zebra chip pathogen, in potato and
748 tomato. Phytopathology. 101: 1285–1291.
- 749 **Li, W., J. A. Abad, R. D. French-Monar, R. J., A. Wen, N. C. Gudmestad, G. A. Secor, I. M.**
750 **Lee, Y. Duan, and L. Levy. 2009.** Multiplex real-time PCR for detection,
751 identification and quantification of "*Candidatus Liberibacter solanacearum*" in
752 potato plants with zebra chip. J. Microbiol. Methods. 78: 59–65.
- 753 **Li, W., J. S. Hartung, and L. Levy. 2006.** Quantitative real-time PCR for detection and
754 identification of *Candidatus Liberibacter* species associated with citrus
755 huanglongbing. J. Microbiol. Methods. 66: 104–115.
- 756 **Liefting, L. W., L. I. Ward, J. B. Shiller, and G. R. G. Clover. 2008.** A new "*Candidatus*
757 *Liberibacter*" species in *Solanum betaceum* (tamarillo) and *Physalis peruviana* (cape
758 gooseberry) in New Zealand. Plant Dis. 92: 1588–1588.
- 759 **Liefting, L. W., B. S. Weir, S. R. Pennycook, and G. R. G. Clover. 2009.** '*Candidatus*
760 *Liberibacter solanacearum*', associated with plants in the family Solanaceae. Int. J.
761 Syst. Evol. Microbiol. 59: 2274–2276.
- 762 **Lin, H., H. Doddapaneni, J. E. Munyaneza, E. L. Civerolo, V. G. Sengoda, J. L. Buchman,**
763 **and D. C. Stenger. 2009.** Molecular characterization and phylogenetic analysis of
764 16S rRNA from a new "*Candidatus Liberibacter*" strain associated with zebra chip
765 disease of potato (*Solanum tuberosum* L.) and the potato psyllid (*Bactericera*
766 *cockerelli* Sulc). J. Plant Pathol. 91: 215–219.
- 767 **Liu, D., and J. T. Trumble. 2004.** Tomato psyllid behavioral responses to tomato plant
768 lines and interactions of plant lines with insecticides. J. Econ. Entomol. 97: 1078–
769 1085.
- 770 **Marchini, D., G. D. Bene, R. Viscuso, and R. Dallai. 2011.** Sperm storage by
771 spermatodotes in the spermatheca of *Trioza alacris* (Flor, 1861) Hemiptera,
772 Psylloidea, Triozidae: A structural and ultrastructural study. J. Morphol. 273: 195–
773 210.
- 774 **Martin, N. A. 2008.** Host plants of the potato/tomato psyllid: A cautionary tale. The Weta.
775 35: 12–16.
- 776 **Marzachi, C., F. Beratti, and D. Bosco. 1998.** Direct PCR detection of phytoplasmas in
777 experimentally infected insects. Ann. Appl. Biol. 133: 45–54.
- 778 **Mas, F., J. Vereijssen, and D. M. Suckling. 2014.** Influence of the pathogen *Candidatus*
779 *Liberibacter solanacearum* on tomato host plant volatiles and psyllid vector
780 settlement. J. Chem. Ecol. 40: 1197–1202.

- 781 **Mayer, C. J., A. Vilcinskis, and J. Gross. 2008.** Phytopathogen lures its insect vector by
782 altering host plant odor. *J. Chem. Ecol.* 34: 1045–1049.
- 783 **Munyanze, J. E. 2012.** Zebra chip disease of potato: Biology, epidemiology, and
784 management. *Am. J. Potato Res.* 89: 329–350.
- 785 **Munyanze, J. E., J. L. Buchman, V. G. Sengoda, T. W. Fisher, and C. C. Pearson. 2011.**
786 Susceptibility of selected potato varieties to zebra chip potato disease. *Am. J. Potato*
787 *Res.* 88: 435–440.
- 788 **Munyanze, J. E., J. L. Buchman, J. E. Upton, J. A. Goolsby, J. M. Crosslin, G. Bester, G. P.**
789 **Miles, and V. G. Sengoda. 2008.** Main content area impact of different potato
790 psyllid populations on zebra chip disease incidence, severity, and potato yield.
791 *Subtropical plant science.* 60: 27–37.
- 792 **Munyanze, J. E., J. M. Crosslin, and J. E. Upton. 2007.** Association of *Bactericera*
793 *cockerelli* (Homoptera: Psyllidae) with 'zebra chip,' a new potato disease in
794 southwestern United States and Mexico. *J. Econ. Entomol.* 100: 656–663.
- 795 **Munyanze, J. E., V. G. Sengoda, J. M. Crosslin, G. D. la Rosa-Lozano, and A. Sanchez.**
796 **2009.** First report of "*Candidatus Liberibacter psyllaeus*" in potato tubers with
797 zebra chip disease in Mexico. *Plant Dis.* 93: 552–552.
- 798 **Murphy, A. F., S. I. Rondon, and A. S. Jensen. 2012.** First report of potato psyllids,
799 *Bactericera cockerelli*, overwintering in the Pacific Northwest. *Am. J. Potato Res.* 90:
800 294–296.
- 801 **Mustafa, T., D. R. Horton, W. R. Cooper, K. D. Swisher, R. S. Zack, H. R. Pappu, and J. E.**
802 **Munyanze. 2015.** Use of electrical penetration graph technology to examine
803 transmission of "*Candidatus Liberibacter solanacearum*" to potato by three
804 haplotypes of potato psyllid (*Bactericera cockerelli*; (Hemiptera: Triozidae). *PLoS*
805 *ONE.* 10: e0138946.
- 806 **Nachappa, P., J. Levy, and C. Tamborindeguy. 2012.** Transcriptome analyses of
807 *Bactericera cockerelli* adults in response to "*Candidatus Liberibacter solanacearum*"
808 infection. *Mol. Genet. Genomics.* 287: 803–817.
- 809 **Nachappa, P., A. A. Shapiro, and C. Tamborindeguy. 2012.** Effect of "*Candidatus*
810 *Liberibacter solanacearum*" on fitness of its insect vector, *Bactericera cockerelli*
811 (Hemiptera: Triozidae), on tomato. *Phytopathology.* 102: 41–46.
- 812 **Nachappa, P. J. Levy, E. Pierson, and C. Tamborindeguy. 2014.** Correlation between
813 "*Candidatus Liberibacter solanacearum*" infection levels and fecundity in its psyllid
814 vector. *J. Invertebr. Pathol.* 115: 55–61.
- 815 **NASS Northwest Regional Field Office, U. S. D. A. 2017.** Potato size and grade summary –
816 2017 crop. United States Department of Agriculture - National Agricultural Statistics
817 Service.

§18 **Navarre, D. A., R. Shakya, J. Holden, and J. M. Crosslin. 2009.** LC-MS analysis of phenolic
 819 compounds in tubers showing zebra chip symptoms. *Am. J. Potato Res.* 86: 88–95.

§20 **Patt, J. M., W. G. Meikle, A. Mafra-Neto, M. Sétamou, R. Mangan, C. Yang, N. Malik, and J.**
 821 **J. Adamczyk. 2011.** Multimodal cues drive host-plant assessment in asian citrus
 822 psyllid (*Diaphorina citri*). *Environ. Entomol.* 40: 1494–1502.

§23 **Pfeiffer, D. G., and E. C. Burts. 1983.** Effect of tree fertilization on numbers and
 824 development of pear psylla (Homoptera: Psyllidae) and on fruit damage. *Environ.*
 825 *Entomol.* 12: 895–901.

§26 **Pfeiffer, D. G., and E. C. Burts. 1984.** Effect of tree fertilization on protein and free amino
 827 acid content and feeding rate of pear psylla (Homoptera: Psyllidae). *Environ.*
 828 *Entomol.* 13: 1487–1490.

§29 **Prager, S. M., I. Esquivel, and J. T. Trumble. 2014.** Factors influencing host plant choice
 830 and larval performance in *Bactericera cockerelli*. *PLoS ONE.* 9: e94047.

§31 **Prager, S. M., O. M. Lewis, J. Michels, and C. Nansen. 2014.** The influence of maturity and
 832 variety of potato plants on oviposition and probing of *Bactericera cockerelli*
 833 (Hemiptera: Triozidae). *Environ. Entomol.* 43: 402–409.

§34 **Prager, S. M., B. Vindiola, G. S. Kund, F. J. Byrne, and J. T. Trumble. 2013.**
 835 Considerations for the use of neonicotinoid pesticides in management of *Bactericera*
 836 *cockerelli* (Sulky) (Hemiptera: Triozidae). *Crop Prot.* 54: 84–91.

§37 **Prager, S. M., C. M. Wallis, M. Jones, R. Novy, and J. T. Trumble. 2017.** Examining the
 838 potential role of foliar chemistry in imparting potato germplasm tolerance to potato
 839 psyllid, green peach aphid, and zebra chip disease. *J. Econ. Entomol.* 111: 327–336.

§40 **Putten, W. H. V. der, L. E. M. Vet, J. A. Harvey, and F. L. Wäckers. 2001.** Linking above -
 841 and belowground multitrophic interactions of plants, herbivores, pathogens, and
 842 their antagonists. *Trends in Ecology & Evolution.* 16: 547–554.

§43 **Qazi, M. C. B., and L. Hogdal. 2010.** Hold on: Females modulate sperm depletion from
 844 storage sites in the fly *drosophila melanogaster*. *J. Insect Physiol.* 56: 1332–1340.

§45 **Rashed, A., T. D. Nash, L. Paetzold, F. Workneh, and C. M. Rush. 2012.** Transmission
 846 efficiency of “*Candidatus Liberibacter solanacearum*” and potato zebra chip disease
 847 progress in relation to pathogen titer, vector numbers, and feeding sites.
 848 *Phytopathology.* 102: 1079–1085.

§49 **Rashidi, M., R. G. Novy, C. M. Wallis, and A. Rashed. 2017.** Characterization of host plant
 850 resistance to zebra chip disease from species-derived potato genotypes and the
 851 identification of new sources of zebra chip resistance. *PLoS ONE.* 12: e0183283.

§52 **R Core Team. 2013.** R: A language and environment for statistical computing. R foundation
 853 for statistical computing, Vienna, Austria.

§54 **Rehman, M., J. C. Melgar, C. J. M. Rivera, A. M. Idris, and J. K. Brown. 2010.** First report
 855 of "*Candidatus* Liberibacter psyllae" or "*ca.* Liberibacter solanacearum"
 856 associated with severe foliar chlorosis, curling, and necrosis and tuber discoloration
 857 of potato plants in Honduras. *Plant Dis.* 94: 376–376.

§58 **Richards, B. L. 1928.** A new and destructive disease of the potato in Utah and its relation
 859 to the potato psylla. *Phytopathology.* 18.

§60 **Richards, H. L., and H. L. Blood. 1973.** Psyllid yellows of the potato. Readings in insect-
 861 plant disease relationships. 46: 139.

§62 **Rosson, P., M. Niemeyer, M. Palma, and L. Ribera. 2006.** Economic impacts of zebra
 863 chips on the Texas potato industry. Center for North American studies, department
 864 of agricultural economics, Texas A&M university, College Station, TX.

§65 **Rubio-Covarrubias, O. A., M. A. Cadena-Hinojosa, S. M. Prager, C. M. Wallis, and J. T.**
 866 **Trumble. 2017.** Characterization of the tolerance against zebra chip disease in
 867 tubers of advanced potato lines from Mexico. *Am. J. Potato Res.* 94: 342–356.

§68 **Sandanayaka, W. R. M., A. Moreno, L. K. Tooman, N. E. M. Page-Weir, and A. Fereres.**
 869 **2014.** Stylet penetration activities linked to the acquisition and inoculation of
 870 *Candidatus* Liberibacter solanacearum by its vector tomato potato psyllid. *Entomol.*
 871 *Exp. Appl.* 151: 170–181.

§72 **Schnakenberg, S. L., W. R. Matias, and M. L. Siegal. 2011.** Sperm-storage defects and live
 873 birth in *drosophila* females lacking spermathecal secretory cells. *PLoS Biology.* 9.

§74 **Secor, G. A., and V. V. Rivera-Varas. 2004.** Emerging diseases of cultivated potato and
 875 their impact on latin america. *Revista Latinoamericana de la Papa (Suplemento).* 1:
 876 1–8.

§77 **Stroup, W. W. 2015.** Rethinking the analysis of non-normal data in plant and soil science.
 878 *Agron. J.* 107: 811.

§79 **Swisher, K. D., and J. M. Crosslin. 2014.** Restriction digestion method for haplotyping the
 880 potato psyllid, *Bactericera cockerelli*. *Southwest. Entomol.* 39: 49–56.

§81 **Šulc, K. 1909.** *Trioza cockerelli* n. Sp., a novelty from North America, being also of economic
 882 importance. *Acta Societatis Entomologicae Bohemiae.* 6: 102–108.

§83 **Teixeira, D. do C., C. Saillard, S. Eveillard, J. L. Danet, P. I. da Costa, A. J. Ayres, and J.**
 884 **Brové. 2005.** '*Candidatus* Liberibacter americanus', associated with citrus
 885 huanglongbing (greening disease) in Sao Paulo State, Brazil. *Int. J. Syst. Evol.*
 886 *Microbiol.* 55: 1857–1862.

§87 **Teulon, D. A. J., P. J. Workman, K. L. Thomas, and M. C. Nielsen. 2009.** *Bactericera*
 888 *cockerelli*: Incurion, dispersal and current distribution on vegetable crops in New
 889 Zealand. *New Zealand Plant Protection.* 62: 136–144.

- §90 **Thinakaran, J., E. A. Pierson, M. Longnecker, C. Tamborindeguy, J. E. Munyaneza, C. M.**
 891 **Rush, and D. C. Henne. 2015.** Settling and ovipositional behavior of *Bactericera*
 892 *cockerelli* (Hemiptera: Triozidae) on solanaceous hosts under field and laboratory
 893 conditions. J. Econ. Entomol. 108: 904–916.
- §94 **Vega-Gutiérrez, M. T., J. C. Rodríguez-Maciel, O. Díaz-Gómez, R. Bujanos-Muñiz, D.**
 895 **Mota-Sánchez, J. L. Martínez-Carrillo, A. Lagunes-Tejeda, and J. A. Garzón-**
 896 **Tiznado. 2008.** Susceptibility to insecticides in two Mexican population of tomato-
 897 potato psyllid, *Bactericera cockerelli* (Sulc.) (Hemiptera: Triozidae). Agrociencia. 42:
 898 463–471.
- §99 **Wallis, R. L. 1955.** Ecological studies on the potato psyllid as a pest of potatoes. U.S.
 900 Department of Agriculture; US Department of Agriculture.
- §01 **Wenninger, E. J., A. Carroll, J. Dahan, A. V. Karasev, M. Thornton, J. Miller, P. Nolte, N.**
 902 **Olsen, and W. Price. 2017.** Phenology of the potato psyllid, *Bactericera cockerelli*
 903 (Hemiptera: Triozidae), and "*Candidatus Liberibacter solanacearum*" in commercial
 904 potato fields in Idaho. Environ. Entomol. 46: 1179–1188.
- §05 **Wenninger, E. J., and D. G. Hall. 2008.** Importance of multiple mating to female
 906 reproductive output in *Diaphorina citri*. Physiol. Entomol. 33: 316–321.
- §07 **Wenninger, E. J., L. L. Stelinski, and D. G. Hall. 2009.** Roles of olfactory cues, visual cues,
 908 and mating status in orientation of *Diaphorina citri* Kuwayama (Hemiptera:
 909 Psyllidae) to four different host plants. Environ. Entomol. 38: 225–234.
- §10 **Wolfner, M. F. 2011.** Precious essences: Female secretions promote sperm storage in
 911 *Drosophila*. PLoS Biology. 9.
- §12 **Yang, X. B., and T. X. Liu. 2009.** Life history and life tables of *Bactericera cockerelli*
 913 (Homoptera: Psyllidae) on eggplant and bell pepper. Environ. Entomol. 38: 1661–
 914 1667.
- §15 **Yang, X. B., Y. M. Zhang, D. C. Henne, and T. X. Liu. 2013.** Life tables of *Bactericera*
 916 *cockerelli* (Hemiptera: Triozidae) on tomato under laboratory and field conditions in
 917 southern Texas. Fla. Entomol. 96: 904–913.
- §18 **Yang, X. B., Y. M. Zhang, L. Hua, and T. X. Liu. 2010.** Life history and life tables of
 919 *Bactericera cockerelli* (Hemiptera: Psyllidae) on potato under laboratory and field
 920 conditions in the Lower Rio Grande Valley of Texas. J. Econ. Entomol. 103: 1729–
 921 1734.
- §22 **Yao, J., P. Saenkham, J. Levy, F. Ibanez, C. Noroy, A. Mendoza, O. Huot, D. F. Meyer, and**
 923 **C. Tamborindeguy. 2016.** Interactions of "*Candidatus Liberibacter*" solanacearum -
 924 *Bactericera cockerelli*: Haplotype effect on vector fitness and gene expression
 925 analyses. Front. Cell. Infect. Microbiol. 6.