

**1. KSU Medicine Ortho Final 2022**

1. In correction of clubfoot BY manipulation. Which deformity should be corrected first
 - A) Forefoot adduction
 - B) Varus
 - C) Upper end tibia
 - D) Calcaneus
2. Non-healing sinus is a common clinical feature in chronic osteomyelitis. The most common frequent cause for this presentation is
 - A) Resistant organism
 - B) Retained foreign body
 - C) Presence of sequestrum
 - D) Intraosseous cavities
3. Grade 3 spondylolisthesis is best treated by
 - A) Medical treatment
 - B) Surgical transpedicular fixation
 - C) Reduction and inter body fusion
 - D) Brace
4. Axillary nerve injury signs include
 - A) Deltoid atrophy
 - B) Numbness of medial side of the upper arm
 - C) Both of the above
 - D) None of the above
5. All of the following are complications of supracondylar fracture in pediatrics except
 - A) Posterior interosseous nerve injury
 - B) Myositis ossificans
 - C) Volkmann ischemic contracture
 - D) Cubitus varus
6. All of the following are signs of compartment syndrome except
 - A) Paresthesia
 - B) Exaggerated pain
 - C) Pain with passive toe stretch
 - D) Hotness of the affected leg
7. The most common organism causing osteomyelitis in sickle cell anemia is
 - A) Streptococcus
 - B) Staph epidermidis
 - C) Pseudomonas
 - D) Salmonella
8. Sure diagnostic sign of septic arthritis
 - A) Raised ESR
 - B) Pus aspiration
 - C) Leukocytosis
 - D) Elevated CRP
9. All are true about Brodies abscess except
 - A) Staph aureus is the most common organism
 - B) Common in long bones of lower extremity
 - C) Represent subacute osteomyelitis
 - D) Sclerotic lesion with irregular rim
10. Cortical destruction in chronic osteomyelitis is
 - A) Cloacae
 - B) Involucrum



- C) Sequestrum
 - D) None of the above
11. All is true about osteochondroma except
- A) Most common benign tumor
 - B) Cartilaginous cap
 - C) Bony outgrowth from epiphysis of long bone
 - D) May cause pressure on adjacent structures
12. Popcorn calcifications usually seen in
- A) Osteosarcoma
 - B) Aneurysmal bone cyst
 - C) Enchondroma
 - D) Fibrous dysplasia
13. All of the following are true about plasmacytoma except
- A) Most common primary bone tumor in adults
 - B) Usually between 40 to 60 years
 - C) Nuclear spokes or wheel fashion
 - D) Associated with hypocalcemia
14. First radiological sign of osteoarthritis is
- A) Joint space narrowing
 - B) Osteophyte formation
 - C) Subchondral sclerosis
 - D) Subchondral cyst
15. The main pathology of chondromalacia patellae
- A) Patellar tendon degeneration
 - B) Tender joint line
 - C) Fibrillation of cartilage
 - D) none of the above
16. Unhappy triad of the knee includes injury to all except
- A) Medial collateral ligament
 - B) Anterior cruciate ligament
 - C) Medial meniscus
 - D) Medial patellar ligament
17. All is true about pathological angular deformity except
- A) Symmetrical
 - B) Severe
 - C) Progressive
 - D) Not expected for age
18. Fragmentation & collapse >50% of head represents Catterall classification
- A) I
 - B) II
 - C) III
 - D) IV
19. Signs of slipped capital femoral epiphysis include all except
- A) Moderate shortening
 - B) Leg held in internal rotation
 - C) Sudden pain
 - D) None of the above
20. Most common osteoporotic fractures include all except
- A) Tibial fracture
 - B) Neck femur fracture





- C) Proximal Humeral fracture
- D) Colles fracture
21. All risk factors for post-menopausal osteoporosis include all except
 - A) Lack of exercises
 - B) Obesity
 - C) Oophorectomy
 - D) Family History
22. Endochondral ossification is a type of
 - A) Epiphyseal Growth
 - B) Diaphyseal Growth
 - C) None of the above
 - D) Surface Growth
23. All are good for bone except
 - A) Vit D
 - B) Corticosteroid
 - C) Calcitonin
 - D) Estrogen
24. Looser's zones commonly seen in all of the following except
 - A) Neck femur
 - B) Neck scapula
 - C) Upper humerus
 - D) Pubis
25. Subperiosteal resorption of middle phalanges is a sign of
 - A) Hypoparathyroidism
 - B) Hyperparathyroidism
 - C) Hypothyroidism
 - D) Hyperthyroidism
26. Ring epiphysis is pathognomonic of
 - A) Rickets
 - B) Scurvy
 - C) Osteomalacia
 - D) Hyperparathyroidism
27. Site for most osteosarcoma
 - A) Upper tibia
 - B) Middle tibia
 - C) Lower tibia
 - D) Lower humerus
28. All are surface signs of osteosarcoma except
 - A) Periosteal new bone formation
 - B) Metaphyseal bone destructive lesion
 - C) Fluffy cotton wool calcifications
 - D) Sunburst appearance



Answers

1-A	2-C	3-C	4-C	5-A	6-D	7-D	8-B	9-D	10-C	11-C	12-C	13-D	14-A
15-C	16-D	17-A	18-D	19-B	20-A	21-B	22-A	23-B	24-B	25-B	26-B	27-A	28-C



2. KSU Medicine Ortho Final 2023

1. Transverse fracture olecranon most commonly treated by plate and screws

- A. False
- B. True

2. The treatment of advanced osteoarthritis is joint replacement

- A. True
- B. False

3. Middle phalangeal subperiosteal resorption are sign for:

- A. Osteomalacia
- B. Hyperparathyroidism
- C. Rickets
- D. Osteoporosis

4. Cobb method is used to:

- A. Method used to treat scoliosis
- B. Measure the angle of scoliosis curve
- C. Measure the prognosis of scoliosis
- D. Measure the rotational deformity

5. The epiphyseal growth is endochondral ossification

- A. False
- B. True

6. Which of the following is a type of spinal fracture:

- A. All answers are correct
- B. Burst fracture
- C. Fracture dislocation
- D. Compression wedge fracture

7. Grade 2 spondylolisthesis, there will be slippage of:

- A. 70-80% of the upper vertebral body
- B. 25% of the upper vertebral body
- C. 25-50% of the upper vertebral body
- D. 100% of the upper vertebral body

8. The most dangerous complication of fracture femur is:

- A. Pulmonary embolism
- B. Non-union
- C. Malunion
- D. Arthritis

9. Pathology of chondromalacia patella is fibrillation of cartilage

- A. True
- B. False

10. Ewing sarcoma arises from:

- A. Periosteum
- B. Cortical bone
- C. Bone marrow
- D. Epiphysis

11. The most common complication of ankle fractures is arthritis

- A. False
- B. True





12. In osteoarthritis the pathology starts in:

- A. Articular cartilage
- B. Menisci
- C. Synovial membrane
- D. Anterior cruciate ligament

13. Osteomyelitis in children older than 2 years, the epiphysis can be involved

- A. True
- B. False

14. Unhappy triad of the knee all except:

- A. ACL injury
- B. Capsular injury
- C. Medial collateral ligament injury
- D. Medial meniscus injury

15. Menisci function are all except:

- A. Absorb shock
- B. Increase stability
- C. Increase friction
- D. Increase contact area

16. Paget disease of bone is characterized by all except:

- A. Cotton wool exudate skull
- B. Long bone bowing
- C. Coarse trabeculae
- D. Young adult

17. Looser's zones are sign for:

- A. Osteomalacia
- B. Scurvy
- C. Rickets
- D. Osteoporosis

18. Scoliosis incidence is more common in:

- A. Male
- B. Both
- C. Female
- D. Elderly

19. The most diagnostic sign of septic arthritis is:

- A. Pain
- B. Fever
- C. No range of motion
- D. Swelling

20. Osteoporotic fractures common except:

- A. Colles fracture
- B. Shaft femur
- C. Vertebral fracture
- D. Neck of femur

21. Characteristic of malignant tumors all true except:

- A. Cortical destruction
- B. Soft tissue masses
- C. Short zone of transition
- D. Periosteal response





22. The bone metastases are common with:

- A. Colon cancer
- B. Breast cancer
- C. Lung cancer
- D. Stomach cancer

23. The most common bone tumor in adult is lymphoma

- A. True
- B. False

24. The most common benign bone tumor is:

- A. Osteochondroma
- B. Chondroblastoma
- C. Osteoid osteoma
- D. Fibrous dysplasia

25. Radiological sign of chronic osteomyelitis all except:

- A. Sequestrum
- B. Cloaca
- C. Involucrum
- D. Codman triangle

26. Colles fracture associated with radial nerve injury

- A. False
- B. True

27. Most common organisms in osteomyelitis in IV drug abusers are:

- A. E. coli
- B. Staph aureus
- C. H. influenzae
- D. Pseudomonas

28. The best radiological method for diagnosis of sequestrum

- A. MRI
- B. CT 3D
- C. X-ray
- D. Ultrasound

29. The most common joint affected in osteoarthritis is:

- A. Ankle
- B. Hip
- C. Shoulder
- D. Knee

30. Kocher criteria of septic arthritis of hip all except:

- A. Leukocytosis
- B. Fever 38.5
- C. ESR 40
- D. Ability to ambulate

31. Signs of myelopathy include the following except:

- A. Spastic quadriparesis
- B. +ve Hoffman sign
- C. Hypotonia
- D. Ataxic gait



32. Wrist drop occurs when which nerve is injured:

- A. Ulnar nerve
- B. Radial nerve
- C. Axillary nerve
- D. Median nerve

33. In lumbar disc prolapse all the following true except:

- A. Radiculopathy
- B. Hyper-reflexia
- C. Low back pain
- D. Weak foot dorsiflexion

34. Cervical disc C5-6 affects which nerve root:

- A. C3
- B. C6
- C. C5
- D. C7



Answers (2023)

1-B 2-A 3-B 4-B 5-B 6-A 7-C 8-A 9- A 10-C 11-A 12-A
13-A 14-B 15-C 16-D 17-A 18-C 19-C 20-B 21-C 22-B 23-B 24-A
25-D 26-A 27-D 28-B 29-D 30-D 31-C 32-B 33-B 34-B

**3. KSU Medicine Ortho End Module 2023**

1. The most common organisms associated with shunt infections include coagulase-negative Staphylococci (particularly Staphylococcus epidermidis), Staphylococcus aureus.
 - A. True
 - B. False
 2. ACL tears commonly associate:
 - A. MCL tears
 - B. LCL tears
 - C. PCL tears
 - D. Meniscal tears
 3. Ring epiphysis is a sign for:
 - A. Hyperparathyroidism
 - B. Scurvy
 - C. Rickets
 - D. Osteomalacia
 4. Brodie's abscess is a type of:
 - A. Subacute osteomyelitis
 - B. Acute on top of chronic osteomyelitis
 - C. Acute osteomyelitis
 - D. Chronic osteomyelitis
 5. The most common site of enchondromas is:
 - A. Fingers
 - B. Radius
 - C. Vertebra
 - D. Skull
 6. Unicameral bone cyst, all are true except:
 - A. Well demarcated
 - B. Metaphyseal
 - C. Asymptomatic
 - D. Sclerosis
 7. The second most common malignant bone tumor:
 - A. Chondrosarcoma
 - B. Osteblastoma
 - C. Ewing sarcoma
 - D. Osteosarcoma
 8. Nonstructural scoliosis involves the spine and is:
 - A. Idiopathic
 - B. Posttraumatic
 - C. With rotation
 - D. Reversible
 9. The most common bone affected by osteosarcoma:
 - A. Spine
 - B. Fingers
 - C. Skull
 - D. Femur
 10. Brown tumors are signs for:
 - A. Osteomalacia
 - B. Osteoporosis
 - C. Hyperparathyroidism
 - D. Rickets
- 



11. The most common complaint of ACL injury is:
 - A. Lateral knee pain
 - B. Limited ROM
 - C. Giving way
 - D. Medial knee pain
12. Mild compression spine fracture, the patient usually is neurologically intact.
 - A. True
 - B. False
13. Spondyloptosis, according to Meyerding grading system:
 - A. Grade IV
 - B. Grade V
 - C. Grade I
 - D. Grade III
14. The most common complication of posterior hip dislocation:
 - A. Sciatic nerve injury
 - B. Hemorrhage
 - C. Femoral nerve injury
 - D. Lateral cutaneous nerve
15. Most common cause of scoliosis is:
 - A. Congenital
 - B. Idiopathic
 - C. Paralytic
 - D. Traumatic
16. Most common cause of death on fracture pelvis:
 - A. Nonunion
 - B. Neurological injury
 - C. Hemorrhage
 - D. Infection
17. The first radiological sign of osteoarthritis is:
 - A. Subchondral cyst
 - B. Subchondral sclerosis
 - C. Osteophyte formation
 - D. Joint space narrowing
18. Burst spinal fracture associated with retropulsed bone fragment inside the spinal canal
 - A. True
 - B. False
19. Most common site of osteomyelitis is:
 - A. Epiphysis
 - B. Physis
 - C. Metaphysis
 - D. Diaphysis
20. Congenital scoliosis characterized by:
 - A. Vertebral bar
 - B. All of the above
 - C. Rib fusion
 - D. Hemivertebrae
21. Traumatic spondylolisthesis of the axis (C2 vertebra) is also called Hangman's fracture
 - A. True
 - B. False



22. The most common complication of humeral surgical neck fracture is:
- A. Malunion
 - B. Axillary nerve injury
 - C. Stiffness
 - D. Nonunion
23. A direct anterior blow to the knee causes:
- A. Meniscal injury
 - B. Collateral injuries
 - C. PCL injury
 - D. ACL injury
24. All are radiological signs of rickets except:
- A. Cupping of metaphysis
 - B. Irregular metaphysis
 - C. Salt and pepper
 - D. Widened physis

Answers (End Module 2023)

- 1-A
- 2-A
- 3-B
- 4-A
- 5-A
- 6-D
- 7-C
- 8-D
- 9-D
- 10-C
- 11-C
- 12-A
- 13-B
- 14-A
- 15-B
- 16-C
- 17-D
- 18-A
- 19-C
- 20-B
- 21-A
- 22-B
- 23-C
- 24-C



4. Final 3/9/2025 لائحہ قدیمہ جزء الاورثو

1. Give short account on knee ligamentous injury (4 marks)
2. Give short account on Shoulder dislocation (4 Marks)
3. Give short account on radiological criteria of Perthes disease (3.5 marks)
4. Discuss Fracture Neck Femur (3.5 Marks)





5. Final Ortho 2025 جزء الازرثو لانه قديمه

1. **Lachman test is usually done at**
 - A. Full extension
 - B. 20-30 flexion
 - C. 60 flexion
 - D. 90 flexion
2. **Positive apprehension test signifies**
 - A. Subacromial impingement
 - B. Supraspinatus tear
 - C. Infrapinatus tear
 - D. Shoulder instability
3. **Apparent limb length is measured from**
 - A. Umbilicus to medial malleolus
 - B. Anterior superior iliac spine to medial malleolus
 - C. Chin to medial malleolus
 - D. All of the above
4. **Straight leg raising test is used to assess**
 - A. Cervical radiculopathy
 - B. Lumbar radiculopathy
 - C. Myelopathy
 - D. Sacroiliitis
5. **C8 nerve root motor function is tested by**
 - A. Finger flexion
 - B. Wrist flexion
 - C. Wrist extension
 - D. Finger abduction
6. **What are uses of gamma nail**
 - A. Trochanteric fractures
 - B. Subtrochanteric fractures
 - C. Shaft fractures
 - D. All of the above
7. **Most common deformity after pediatric supracondylar humeral fracture is**
 - A. Cubitus valgus
 - B. Genu valgus
 - C. Cubitus varus
 - D. Coxa varus
8. **Dashboard injury can lead to**
 - A. Posterior hip dislocation
 - B. Posterior cruciate ligament injury
 - C. Patellar fracture
 - D. All of the above
9. **External fixator is indicated in**
 - A. Pelvic fractures
 - B. Open fractures
 - C. Comminuted fractures
 - D. All of the above
10. **Tension band wiring can be used in**
 - A. Olecranon fractures
 - B. Patellar fractures
 - C. Medial malleolar fracture
 - D. All of the above

**6. Ortho Final 2024 (PT Ortho Diseases)**

1. Osteomyelitis that is less than 2 weeks duration is called
 - A. Acute om
 - B. Subacute om
 - C. Chronic om
 - D. None of above
2. Commonest site of acute osteomyelitis in children is
 - A. Metaphyseal
 - B. Diaphyseal
 - C. Epiphyseal
 - D. Physeal
3. Lab tests of acute osteomyelitis all except
 - A. Increased ESR
 - B. Increased CRP
 - C. Leucopenia
 - D. Aspiration reveals pus cells
4. Treatment of acute om
 - A. No role for surgical ttt
 - B. Bed rest - antibiotic - good general condition for mild cases
 - C. Physical therapy for pain relief
 - D. Fixation by plate and screw
5. Indolent course of subacute om is not due to
 - A. Increased host resistance
 - B. Decreased bacterial virulence
 - C. Antibiotic administration before onset of om
 - D. Repeated multiple debridement
6. Regarding subacute osteomyelitis all are true except
 - A. Leucocytosis in 100% of cases
 - B. Increased ESR in 50% of cases
 - C. Blood culture usually negative
 - D. Plain x-ray generally positive
7. Gold standard for diagnosis of chronic OM
 - A. Infected dead bone with compromised patient
 - B. Presence of sequestrum
 - C. New bone formation by periosteum
 - D. All of the above
8. Surgical ttt of Chronic osteomyelitis is not
 - A. Sequestrectomy
 - B. Radical debridement
 - C. Antibiotic loaded cement
 - D. Avoid bone resection
9. Most common joint affected in septic arthritis is
 - A. Knee
 - B. Hip
 - C. Shoulder
 - D. Elbow
10. Most diagnostic sign of septic arthritis is
 - A. Marked tender limitation of ROM
 - B. Increase ESR CRP



- C. Positive Blood culture
- D. Fever
11. Treatment of acute septic arthritis
 - A. Surgical emergency for debridement and drainage
 - B. Antibiotic therapy in mild cases with follow up
 - C. Splinting of affected joint
 - D. All of above
12. Knee OA show in x-ray all except
 - A. Deformity
 - B. Subchondral lucency
 - C. Joint space narrowing
 - D. Osteophytes
13. Behavioral ttt of knee OA include except
 - A. Weight loss
 - B. Intra Articular injection
 - C. Mechanical aid
 - D. Exercises
14. Regarding surgical ttt of knee OA all are true except
 - A. Total knee replacement
 - B. Deformity correction
 - C. Arthroscopic debridement
 - D. Ligament reconstruction
15. Osteoporosis can lead to
 - A. Colles fracture
 - B. Femoral neck fracture
 - C. Vertebral fracture
 - D. All of the above
16. Osteoporosis is mainly due to
 - A. Decreased bone mass
 - B. Decreased ca & vit d
 - C. Increased phosphorus in bone matrix
 - D. Increased alkaline phosphatase enzyme
17. Medications not used in ttt of osteoporosis
 - A. Ca supplements
 - B. Vit d
 - C. Bisphosphonates
 - D. Steroid
18. Rickets is mainly due to
 - A. Decreased mineralization in growing bone
 - B. Decreased mineralization in adult bone
 - C. Reduction of bone mass
 - D. Decreased alkaline phosphatase enzyme
19. Manifestations of rickets is not
 - A. Enlarged metaphysis
 - B. Rosary chest
 - C. Cod fish vertebrae
 - D. Deformity
20. X-ray of rickets include
 - A. Narrow metaphysis
 - B. Normal shaft





- C. Wide physis
- D. Abnormal epiphysis
21. Lab results of rickets most diagnostic
 - A. Low ca
 - B. Low phosphorus
 - C. Low vit D
 - D. Marked Increased ALK Phosphatase
22. X ray of osteomalacia not include
 - A. Trefoil pelvis
 - B. Looser zone
 - C. Biconcave vertebra
 - D. Codman triangle
23. Brown tumor is seen in
 - A. Hyperparathyroidism
 - B. Osteomalacia
 - C. Osteoporosis
 - D. Rickets
24. Causes of out toeing gait is not due to
 - A. Abnormal Femoral version
 - B. Tibial external torsion
 - C. Forefoot abduction
 - D. Genu varus deformity
25. Genu valgus is not
 - A. Known as bow legs
 - B. Physiological at 2.5 age
 - C. Increase stress on lateral epiphysis of knee
 - D. Corrected by proximal tibial osteotomy
26. Flat foot is known as
 - A. Pes planus
 - B. Talipes equino varus
 - C. Pes cavus
 - D. Cavovarus foot
27. Increased neck shaft angle of proximal femur
 - A. Coxa valga
 - B. Coxa vara
 - C. Coxa magna
 - D. Coxa previa
28. The following are etiologies of scoliosis except
 - A. Idiopathic
 - B. Congenital
 - C. Neuromuscular
 - D. Adolescent scoliosis
29. Hemivertebra causes
 - A. Congenital scoliosis
 - B. Neuromuscular scoliosis
 - C. Idiopathic scoliosis
 - D. Adolescent scoliosis
30. Positive signs used for diagnosis of scoliosis
 - A. Cobb angle >20
 - B. Adam bending test positive





- C. Scoliometer positive
- D. All of the above
- 31. Rib fusion is a component of
 - A. Congenital scoliosis
 - B. Idiopathic scoliosis
 - C. Neuromuscular scoliosis
 - D. Adult type scoliosis
- 32. Forward slipping of all or part of vertebra is called
 - A. Spondylosis
 - B. Spondylolisthesis
 - C. Spondylolysis
 - D. Spondyloptosis
- 33. The most common symptom of spondylolisthesis is
 - A. Low back pain
 - B. Referred lower back pain
 - C. Intermittent claudication
 - D. Foot drop
- 34. Slippage of >25% of vertebra on another is spondylolisthesis
 - A. Type I
 - B. Type II
 - C. Type III
 - D. Type IV
- 35. Most common bone malignancy
 - A. Ewing sarcoma
 - B. Bone metastasis
 - C. Osteosarcoma
 - D. Giant cell tumor
- 36. Most common benign bone tumor
 - A. Osteoid osteoma
 - B. Osteochondroma
 - C. Osteoblastoma
 - D. Aneurysmal bone cyst
- 37. Enchondroma is not
 - A. Most common in fingers
 - B. Bone cell inside cartilage
 - C. Popcorn calcification in x-ray
 - D. Require no ttt in non symptomatic cases
- 38. Central nidus is found in
 - A. Osteoid osteoma
 - B. Enchondroma
 - C. Chondroblastoma
 - D. Simple bone cyst
- 39. Giant cell tumor commonly found in
 - A. Epiphysis
 - B. Metaphysis
 - C. Diaphysis
 - D. Flat bones
- 40. Giant cell tumor differs from benign tumors in
 - A. Send distal metastasis
 - B. Can recur after excision






- C. Severe periosteal reaction
 - D. Chemotherapy is preferred ttt
41. Osteosarcoma can be
- A. Telangiectatic
 - B. Parosteal
 - C. Periosteal
 - D. Mesenchymal
42. Codman triangle is not
- A. A periosteal reaction in a tumor
 - B. Known as sun ray appearance
 - C. Found on all bone tumor
 - D. Represent new bone formation
43. Onion skin layers is found in
- A. Osteosarcoma
 - B. Ewing sarcoma
 - C. Chondrosarcoma
 - D. Multiple myeloma
44. Most common bone tumor in adult is
- A. Ewing sarcoma
 - B. Multiple myeloma
 - C. Osteosarcoma
 - D. Chondrosarcoma
45. Lab finding in multiple myeloma is not
- A. Bence Jones protein in urine
 - B. Increase alkaline phosphatase enzyme
 - C. Increase Ca
 - D. Increase serum globulin
46. Most common site that metastasizes to bone is
- A. Kidney
 - B. Liver
 - C. Breast
 - D. Uterus
47. Most sensitive method for detection of metastasis
- A. X-ray
 - B. Bone scan
 - C. MRI
 - D. CT scan
48. CTS is entrapment of which nerve
- A. Median
 - B. Ulnar
 - C. Radial
 - D. Posterior interosseous
49. CTS treatment is all except
- A. Splints
 - B. Neurotonics
 - C. Surgical neurectomy
 - D. Local injection
50. Shaking wrist relief symptoms of CTS is called
- A. Tinel sign
 - B. Flick sign






- C. Phalen sign
- D. Erb's sign
51. Pronator syndrome is entrapment of which nerve
 - A. Radial nerve
 - B. Nerve to pronator quadratus
 - C. Anterior interosseous nerve
 - D. Ulnar nerve
52. Failure to do OK sign is pathognomonic to
 - A. Median nerve affection
 - B. Ulnar nerve affection
 - C. Radial nerve affection
 - D. Posterior interosseous nerve affection
53. Ulnar nerve can be entrapped in
 - A. Cubital tunnel
 - B. Guyon's canal
 - C. All of the above
 - D. None of the above
54. Clawing seen due to ulnar nerve affection is at
 - A. Ring & little
 - B. Index & middle
 - C. Thumb
 - D. All fingers
55. Radial nerve entrapment at different sites cause except
 - A. Posterior interosseous nerve syndrome
 - B. Radial tunnel syndrome
 - C. Guyon canal syndrome
 - D. Wartenberg syndrome
56. Piriformis syndrome is due to
 - A. Disc bulge causing sciatic nerve entrapment
 - B. Compression by tendon or muscle causing sciatica
 - C. Compression of lateral femoral cutaneous nerve affection
 - D. Compression of obturator nerve
57. Entrapment of LFCN is called
 - A. Meralgia paresthetica
 - B. Piriformis syndrome
 - C. Sciatica
 - D. None of the above
58. Tarsal tunnel syndrome is compression of
 - A. Tibial nerve branches under flexor retinaculum
 - B. Superficial peroneal nerve under extensor retinaculum
 - C. Deep peroneal nerve under extensor retinaculum
 - D. Common peroneal nerve at fibular head
59. Rotator cuff muscles include except
 - A. Deltoid muscle
 - B. Supraspinatus muscle
 - C. Teres minor muscle
 - D. Subscapularis muscle
60. Job's test best evaluates
 - A. Abduction of shoulder by deltoid muscle
 - B. Abduction of shoulder by supraspinatus muscle





- C. Internal rotation of shoulder by subscapularis muscle
- D. Internal rotation of shoulder by supraspinatus muscle
61. Drop arm sign is pathognomonic for
 - A. Rotator cuff tear
 - B. Shoulder dislocation
 - C. Shoulder impingement
 - D. Frozen shoulder
62. Limited active but preserved passive movement of the shoulder is seen in
 - A. Rotator cuff tear
 - B. Shoulder dislocation
 - C. Frozen shoulder
 - D. None of the above
63. Neer test is best to evaluate
 - A. Rotator cuff tear
 - B. Shoulder dislocation
 - C. Frozen shoulder
 - D. Shoulder impingement
64. Failure to press the abdomen by palm of the hand in adducted shoulder is due to
 - A. Supraspinatus tear
 - B. Subscapularis tear
 - C. Shoulder impingement
 - D. Teres minor tear
65. The following muscle is responsible for external rotation of the shoulder
 - A. Teres minor
 - B. Infraspinatus
 - C. All of the above
 - D. None of the above
66. Most common affected nerve in shoulder dislocation
 - A. Axillary
 - B. Radial
 - C. Median
 - D. Ulnar
67. Most common variant of shoulder dislocation
 - A. Anterior
 - B. Posterior
 - C. Inferior
 - D. None of the above
68. All of the following are methods of shoulder reduction except
 - A. Hippocrates method
 - B. Kocher method
 - C. Stimpson method
 - D. Flynn method
69. Best test for evaluation of shoulder stability
 - A. Neer test
 - B. Hawkins test
 - C. Apprehension test
 - D. Belly press test
70. Shoulder joint is composed of
 - A. 3 bones 3 joints
 - B. 3 bones 2 joints





- C. 2 bones 2 joints
 - D. 2 bones 3 joints
71. Shoulder dislocation is not
- A. Emergency reduction is mandatory
 - B. Reduction under anesthesia may be needed
 - C. Can be associated with fractures
 - D. Posterior dislocation is more common
72. Exaggeration of lumbar spine curve is
- A. Lordosis
 - B. Kyphosis
 - C. Scoliosis
 - D. Ankylosis
73. All of the following are malignant tumors of bone and cartilage except
- A. Osteosarcoma
 - B. Ewing sarcoma
 - C. Rhabdomyosarcoma
 - D. Chondrosarcoma
74. Regarding chondromalacia patellae all except
- A. May be traumatic in origin
 - B. Fissures in articular surface of the patella
 - C. Also known as Runner's knee
 - D. Treated by artificial joint replacement
75. Kocher criteria for septic arthritis
- A. Temperature ≥ 38
 - B. White cell count > 12000
 - C. Mild limitation of ROM
 - D. ESR, CRP not always high
76. Cerebral palsy is associated with
- A. Neuromuscular scoliosis
 - B. Congenital scoliosis
 - C. Idiopathic scoliosis
 - D. None of the above
77. Moderate scoliosis is Cobb's angle
- A. <20
 - B. 20-40
 - C. 40
 - D. None of the above
78. The following brace is used in TTT of scoliosis
- A. Milwaukee brace
 - B. Boston brace
 - C. Charleston night bending brace
 - D. All of the above
79. Spondylolisthesis can result from except
- A. Congenital
 - B. Degenerative
 - C. Traumatic
 - D. Rheumatic
80. Regarding spondylolisthesis all are true except
- A. Low back pain is major symptoms
 - B. Lumbar fusion is not mandatory





- C. Grade III is displacement >50%
- D. Can cause lumbar canal stenosis

Answers – Ortho Final 2024

1-A 2-A 3-C 4-B 5-D 6-A 7-D 8-D 9-A
 10-A 11-D 12-B 13-B 14-D 15-D 16-A 17-D 18-A
 19-C 20-C 21-D 22-D 23-A 24-D 25-A 26-A 27-A
 28-D 29-A 30-D 31-A 32-B 33-A 34-B 35-B 36-B
 37-A 38-A 39-A 40-B 41-D 42-C 43-B 44-B 45-B
 46-C 47-B 48-A 49-C 50-B 51-C 52-A 53-C 54-A
 55-C 56-B 57-A 58-A 59-A 60-B 61-A 62-A 63-D
 64-B 65-C 66-A 67-A 68-D 69-C 70-B 71-D 72-A
 73-C 74-D 75-B 76-A 77-B 78-D 79-D 80-B



**7. Ortho Final 2025 (PT Ortho Diseases)**

1. To test Shoulder Stability, which test is included?
 - a) Lift off
 - b) Neer
 - c) Hawkins
 - d) Apprehension
2. Restricted active and preserved passive shoulder range of motion may be caused by:
 - a) Rotator Cuff tear
 - b) Frozen Shoulder
 - c) Shoulder Instability
 - d) Bursitis
3. Most common cause of Rotator Cuff tear is degenerative:
 - a) True
 - b) False
4. Mild cases of osteomyelitis can have conservative treatment:
 - a) True
 - b) False
5. Mild acute osteomyelitis treated by antibiotics and follow up:
 - a) True
 - b) False
6. The gold standard for diagnosis of chronic osteomyelitis is laparoscopic investigations:
 - a) True
 - b) False
7. Most common site of septic arthritis is hip:
 - a) True
 - b) False
8. Most diagnostic sign of septic arthritis is positive culture or aspiration:
 - a) True
 - b) False
9. First radiological sign of knee osteoarthritis is subchondral cyst:
 - a) True
 - b) False
10. Osteoporosis can cause fracture of the neck of femur:
 - a) True
 - b) False
11. X-ray radiological sign in rickets is wide metaphysis:
 - a) True
 - b) False
12. Genu varum except:
 - a) Bow legs
 - b) Medial thrust
13. Most common cause of genu varum is Blount disease:
 - a) True
 - b) False
14. Out-toeing can't be because of external tibial torsion:
 - a) True
 - b) False
15. One of the criteria of pathological deformity is regressive:
 - a) True
 - b) False



1. Rheumatology Final 2022

Q/ True or False

1. HLA-B27 is present in 100% of patients with Ankylosing Spondylitis.
2. Subclinical synovitis in RA patients can be detected by X-ray imaging.
3. Anti-Sm antibodies are the most specific and sensitive antibodies in SLE.
4. Avascular necrosis in SLE is a common musculoskeletal manifestation that usually affects the hip.
5. Sacroilitis is diagnosed early in the course of SpA using MRI.
6. The most typical enthesitis is heel pain, related to inflammation of the Achilles tendon or the plantar fascia insertion.
7. Upper limb involvement is typically associated with psoriatic arthritis.
8. Secondary osteoporosis is usually related to disease, medications, or post-menopausal state.
9. The aim of osteoporosis treatment is the prevention of fragility fracture.
10. In soft tissue rheumatism, pain is elicited by passive movement of the joint.
11. Most patients with hyperuricemia do not suffer from clinical gout.
12. The classical presentation of acute gout is characterized by a typical rapid development of severe pain and swelling of a joint, without overlying erythema or tenderness, that reaches its maximum within 6–12 hours.
13. Chronic urate arthropathy is a late feature of neglected gout that is destructive with frequent erosions.
14. JIA is a definite arthritis of unknown etiology that begins before the 16th birthday and persists for at least 6 weeks.
15. Skin lesions are not required for the diagnosis of psoriatic JIA.
16. Osteoarthritis has been relabeled as a whole organ disease.
17. The highest risk for the development of chronic asymptomatic anterior uveitis in JIA is in the oligoarticular category.
18. Obesity, hyperglycemia, hypercholesterolemia, and low levels of vitamin D are predisposing factors for osteoarthritis.
19. Rheumatoid arthritis is a chronic systemic disease that usually affects females after menopause.
20. Anti-CCP antibodies are found in 60–70% of RA patients, but hardly ever in other diseases or healthy subjects.

True / False

1 F 2 F 3 F 4 T 5 T 6 T 7 F 8 T 9 T 10 F
11 T 12 F 13 T 14 T 15 T 16 T 17 T 18 T 19 F 20 T



Q/ Multiple Choice Questions

1. Triad of rash, fever, and arthritis in a female in childbearing period, may raise the diagnosis of:
 - a) RA
 - b) SLE
 - c) Systemic sclerosis
 - d) SpA
2. What is the main diagnostic modality for osteoarthritis?
 - a) Dual energy X-ray absorptiometry (DEXA) scan
 - b) Plain radiography
 - c) Ultrasonography
 - d) MRI
3. Dactylitis occurs most frequently in which of the following among all categories of Spondarthropathy (SpA)?
 - a) Ankylosing spondylitis
 - b) Enteropathic arthritis
 - c) Undifferentiated Spondarthropathy
 - d) Psoriatic arthritis
4. Which one of the following domains is NOT involved in ACR/EULAR criteria for RA diagnosis?
 - a) Joint distribution
 - b) Acute phase reactant
 - c) Serology
 - d) Radiological assessment
5. A common cause of chronic polyarthritis with predominant articular manifestation includes:
 - a) RA
 - b) Systemic sclerosis
 - c) SLE
 - d) Systemic vasculitis
6. The main cytokine involved in the pathogenesis of Rheumatoid arthritis is
 - a) Tumour necrosis factor alpha
 - b) Interferon
 - c) Interleukin 10
 - d) Interleukin 4
7. Which statement is true about Enthesitis related arthritis category of Juvenile idiopathic arthritis?
 - a) It occurs more frequently in females.
 - b) It involves joints of the hand and feet.
 - c) It involves joints of the upper limbs.
 - d) Enthesitis occurs mostly in plantar fascia and tendoachilles.
8. Which statement of the following is true about articular features of Spondarthropathy (SpA)?
 - a) Involvement of large joints of upper limbs.
 - b) Low back pain is uncommon.
 - c) Symmetrical small joint arthritis is common.
 - d) Sacroiliitis is common.
9. Which statement is true about stiffness in osteoarthritis?
 - a) It lasts for 30 minutes and more.
 - b) It lasts for one hour and more.
 - c) It lasts for 15 minutes or less.
 - d) It occurs after a long period of rest but not after a long period of sleep.
10. Which drug is used during the acute attack of gouty arthritis?
 - a) Uricosuric drugs
 - b) Allopurinol
 - c) Colchicine
 - d) Methotrexate

1. B

2. B

3. D

4 D

5 A

6 A

7 D

8 D

9 C

10 C

**2. End Round (1) 2022****1. What is Polyarthritis?**

- a. Arthritis affecting two to four joints during the first six months of disease
- b. Arthritis affecting more than four joints during the first six months of disease
- c. Arthritis affecting one joint during the first six months of disease
- d. Arthritis affecting more than one joint during the first six months of disease

2. A Common cause of acute monoarthritis

- a. gouty arthritis
- b. Rheumatoid arthritis
- c. psoriatic arthritis
- d. systemic sclerosis

3. What is the commonest pattern of arthritis in Rheumatoid arthritis?

- a. Symmetrical small joint arthritis.
- b. Symmetrical large joint arthritis.
- c. Asymmetrical small joint arthritis.
- d. Asymmetrical large joint arthritis

4. Which statement is true about rheumatoid factor (RF)?

- a. It is specific for Rheumatoid arthritis.
- b. It is not specific for Rheumatoid arthritis and occurs in many other conditions.
- c. It occurs in 100% of Rheumatoid arthritis patients.
- d. not included in the diagnostic criteria of Rheumatoid arthritis

5. Which disease of the following is Not one of Spondyloarthropathy (SpA) subgroups?

- a. Ankylosing spondylitis
- b. Reactive arthritis
- c. Sarcoidosis
- d. Psoriatic arthritis

6. The main pathogenic finding in osteoarthritis is

- a. cartilage degeneration
- b. subchondral bone cyst
- c. synovitis
- d. Capsulitis

7. Gouty arthritis is disease caused by the deposition of In joints

- a. Monosodium urate crystals
- b. calcium oxalate crystals
- c. calcium pyrophosphate dihydrate crystals
- d. Calcium phosphate crystals

8. Which immunosuppressant drugs are commonly used to induce remission in lupus nephritis

- a. MTX
- b. Cyclosporine
- c. Leflunomide
- d. Mycophenolate Mofetil

9. Which of the following autoantibodies are the most specific for SLE?

- a. anti-Sm.
- b. ANA
- c. Anti CCP
- d. RF

10. What is the best diagnostic modality for osteoporosis?

- a. Dual energy X-ray absorptiometry (DEXA) scan.
- b. Plain radiography
- c. Ultrasonography
- d. MRI

**11. Which statement is true about inflammatory arthritis?**

- a. Associated morning stiffness of less than 30 minutes
- b. Improvement of symptoms on gentle use of joints
- c. Range of motion is usually normal
- d. not associated with joint tenderness or swelling

12. Which one of the following domains is Not involved in ACR/EULAR diagnosis criteria for RA?

- a. Joint distribution
- b. Acute phase reactant
- c. Serology
- d. Radiological assessment

13. Which statement is true about Enthesitis related arthritis category of Juvenile idiopathic arthritis?

- a. It occurs more frequently in females.
- b. It involves joints of the hand and feet
- c. It involves joints of upper limb
- d. Enthesitis occurs mostly in plantar fascia and tendoachilles

14. The commonest extraarticular manifestations occurring in spondyloarthropathy is

- a. Acute anterior uveitis
- b. Heart block
- c. Aortic insufficiency
- d. Restrictive lung disease

15. Dactylitis occurs most frequently in among all categories of spondyloarthropathy

- a. Ankylosing spondyloarthropathy
- b. Enteropathic arthritis
- c. Undifferentiated spondyloarthropathy
- d. Psoriatic arthritis

16. Which statement is true about stiffness in osteoarthritis

- a. It last for 30 minutes or more
- b. It lasts for one hour or more
- c. It lasts for 15 minutes or less
- d. It occurs after long period of rest but not after long period of sleep

17. The WHO defined osteoporosis as

- a. T score -2.5 or less
- b. T score of -1
- c. T score between 1 and -2.5
- d. T score -3.5 or less

18. Hyperuricemia occurs when serum uric acid levels are

- a. More 6 mg/dl
- b. Less 6 mg/dl
- c. More 10 mg/dl
- d. More 5 mg/dl

19. What is the first line drug therapy for osteoporosis patients

- a. Steroid
- b. Bisphosphonate
- c. Vitamin D
- d. Calcium

20. A higher rate of mortality and morbidity in lupus patients associated with

- a. Renal and neurophycartic changes
- b. Neuropsychiatric changes
- c. Renal involvement
- d. Chronic cutaneous involvement

**3. End Round (2) 2022****1. What is Oligoarthritis?**

- a. Arthritis affecting two to four joints during the first six month of disease
- b. Arthritis affecting more than five joints during the first six months of disease
- c. Arthritis affecting one joint during the first six months of disease
- d. Arthritis affecting more than one joint during the first six months of disease

2. Which statement is true about inflammatory arthritis

- a. Associated morning stillness of less than 30 minutes
- b. Improvement of symptoms on gentle use of joints
- c. Range of motion is usually normal
- d. not associated with tenderness or swelling

3. A Common cause of acute monoarthritis

- a. gouty arthritis
- b. Rheumatoid arthritis
- c. psoriatic arthritis
- d. systemic sclerosis

4. What is the most common Extra-articular Manifestations of Rheumatoid arthritis

- a. Sicca syndrome
- b. Rheumatoid nodules
- c. Pericarditis and pleuritis
- d. Interstitial lung fibrosis

5. What is the commonest pattern of arthritis in Rheumatoid arthritis?

- a. Symmetrical small joint arthritis
- b. Symmetrical large joint arthritis
- c. Asymmetrical small joint arthritis
- d. Asymmetrical large joint arthritis

6. Which one of the following domains is Not involved in ACR/EULAR criteria for RA diagnosis?

- a. Joint distribution
- b. Acute phase reactant
- c. Serology
- d. Radiological assessment

7. The main cytokine involved in the pathogenesis of Rheumatoid arthritis is

- a. Tumor necrosis factor alpha
- b. Interferon
- c. Interleukin 10
- d. Interleukin 4

8. The most frequent category of Juvenile idiopathic arthritis with chronic asymptomatic anterior uveitis

- a. systemic Juvenile idiopathic arthritis
- b. undifferentiated Juvenile idiopathic arthritis
- c. polyarticular Juvenile idiopathic arthritis
- d. oligoarticular Juvenile idiopathic arthritis



9. What is the commonest site of rash in systemic Juvenile idiopathic arthritis?

- a. The face
- b. Upper chest and underarm
- c. Chin of tibia
- d. Hands and feet

10. Which category of Juvenile idiopathic arthritis presents with fever, rash and lymphadenopathy?

- a. Psoriatic Juvenile idiopathic arthritis
- b. Polyarticular Juvenile idiopathic arthritis
- c. Oligoarticular Juvenile idiopathic arthritis
- d. Systemic Juvenile idiopathic arthritis

11. Which statement of the following is true about articular features of Spondylarthropathy (SpA)?

- a. Sacroiliitis is common.
- b. Involvement of large joints of upper limbs.
- c. Low back pain is uncommon.
- d. Symmetrical small joint arthritis is common.

12. The axial affection of Spondylarthropathy is best treated with

- a. NSAIDs.
- b. Steroids
- c. Conventional DMARDs
- d. Biologics

13. The main pathogenic finding in osteoarthritis is

- a. cartilage degeneration
- b. subchondral bone cyst
- c. Synovitis
- d. Capsulitis

14. is mandatory for the diagnosis of knee osteoarthritis.

- a. Weight bearing plain X ray
- b. ultrasonography
- c. MRI
- d. serology

15. The WHO defined osteoporosis as :

- a. T score -2.5 or less
- b. T score of -1
- c. T score between 1 and -2.5
- d. T score -3.5 or less

16. Which one of the following condition is a cause of primary osteoporosis

- a. Hyperparathyroidism
- b. Chronic steroid intake
- c. Ankylosing spondylitis
- d. Post menopausal state



17. Hyperuricemia occurs when serum uric acid levels are

- a. More 6 mg/dl
- b. Less 6 mg/dl
- c. More 10 mg/dl
- d. More 5 mg/dl

18. Which statement is true about acute gouty arthritis?

- a. It presents mostly as acute polyarthritis
- b. It involves a single joint in the upper extremity
- c. In particular, the first metatarsophalangeal joint is involved
- d. Serum uric acid is usually elevated during the attack

19. Which of the following autoantibodies are the most specific for SLE?

- a. anti-dsDNA and anti-Sm.
- b. ANA
- c. Anti CCP
- d. RF

20. Which statement is true about SLE?

- a. It typically affects women between 16 years and 55 years
- b. It typically affects men between 16 years and 55 years
- c. It typically affects pediatric age group
- d. It equally affects women and men between 16 years and 55 years

1 a 2 b 3 a 4 b 5 a 6 d 7 a 8 d 9 b 10 d
11 a 12 a 13 a 14 a 15 a 16 d 17 a 18 c 19 a 20 a

**4. End Round (3) 2022****1. What is Oligoarthritis?**

- a. Arthritis affecting two to four joints during the first six month of disease
- b. Arthritis affecting more than five joints during the first six months of disease
- c. Arthritis affecting one joint during the first six months of disease
- d. Arthritis affecting more than one joint during the first six months of disease

2. A Common cause of acute monoarthritis

- a. gouty arthritis
- b. Rheumatoid arthritis
- c. psoriatic arthritis
- d. systemic sclerosis

3. What is the commonest pattern of arthritis in Rheumatoid arthritis?

- a. Symmetrical small joint arthritis.
- b. Symmetrical large joint arthritis.
- c. Asymmetrical small joint arthritis.
- d. Asymmetrical large joint arthritis

4. Which statement is true about ACPA?

- a. It is specific for Rheumatoid arthritis.
- b. It is not specific for Rheumatoid arthritis and occurs in many other conditions.
- c. It occurs in 100% of Rheumatoid arthritis patients.
- d. not included in the diagnostic criteria of Rheumatoid arthritis

5. Which disease of the following is one of Spondyloarthropathy (SpA) subgroups?

- a. Rheumatoid arthritis
- b. Systemic sclerosis
- c. Sarcoidosis
- d. Psoriatic arthritis

6. The main pathogenic finding in RA is

- a. cartilage degeneration
- b. subchondral bone cyst
- c. synovitis
- d. Capsulitis

7. Gout is a disease caused by the deposition of in joints

- a. monosodium urate crystals
- b. calcium oxalate crystals
- c. calcium pyrophosphate dihydrate crystals
- d. Calcium phosphate crystals

8. Which immunosuppressant drugs are commonly used to induce remission in lupus nephritis

- a. MTX
- b. Cyclosporine
- c. Leflunomide
- d. Mycophenolate Mofetil



9. Which of the following autoantibodies are the most specific for SLE?

- a. anti-Sm.
- b. ANA
- c. Anti CCP
- d. RF

10. What is the best diagnostic modality for osteoarthritis?

- a. Dual energy X-ray absorptiometry (DEXA) scan.
- b. Plain radiography
- c. Ultrasonography
- d. MRI

11. Which statement is true about inflammatory arthritis?

- a. Associated morning stiffness of less than 30 minutes
- b. Improvement of symptoms on gentle use of joints
- c. Range of motion is usually normal
- d. not associated with joint tenderness or swelling

12. Which one of the following domains is Not involved in ACR/EULAR diagnosis criteria for RA?

- a. Joint distribution
- b. Acute phase reactant
- c. Serology
- d. Radiological assessment

13. Which statement is true about Enthesitis related arthritis category of Juvenile idiopathic arthritis?

- a. It occurs more frequently in females.
- b. It involves joints of the hand and feet
- c. It involves joints of upper limb
- d. Enthesitis occurs mostly in plantar fascia and tendoachilles

14. The commonest extraarticular manifestations occurring in RA is

- a. Acute anterior uveitis
- b. Subcutaneous nodules
- c. Aortic insufficiency
- d. ILF

15. Dactylitis occurs most frequently in among all categories of spondyloarthropathy

- a. Ankylosing spondyloarthropathy
- b. Enteropathic arthritis
- c. Undifferentiated spondyloarthropathy
- d. Psoriatic arthritis

16. Which statement is true about symptoms of osteoarthritis

- a. Crepitus is not common
- b. Morning stiffness last for one hour or more
- c. Pain is very common
- d. Deformity is not a finding



17. The WHO defined osteoporosis as

- a. T score -2.5 or less
- b. T score of -1
- c. T score between 1 and -2.5
- d. T score -3.5 or less

18. Hyperuricemia occurs when serum uric acid levels are

- a. More 6 mg/dl
- b. Less 6 mg/dl
- c. More 10 mg/dl
- d. More 5 mg/dl

19. What is the first line drug therapy for osteoporosis patients

- a. Steroid
- b. Bisphosphonate
- c. Vitamin D
- d. Calcium

20. A triad of rash, fever and arthritis in a child bearing period female must raise diagnose of

- a. RA
- b. SLE
- c. Systemic sclerosis
- d. SpA



1 a 2 a 3 a 4 a 5 d 6 c 7 a 8 d 9 a 10 b
11 b 12 d 13 d 14 b 15 d 16 c 17 a 18 a 19 b 20 b

**5. (2023 Final Rheumatology – Not Complete)**

1) Which of the following autoantibodies are the most specific for SLE?

- a. anti CCP
- b. Anti Ro and anti La
- c. anti-dsDNA and anti-Sm
- d. ANA

2) The main aim of the treatment of osteoporosis is prevention of fragility fracture:

- a. true
- b. false

3) A 26-year-old man presents with inflammatory back pain... diagnosis:

- a. Psoriatic arthritis
- b. Diffuse idiopathic skeletal hyperostosis
- c. Ankylosing spondylitis
- d. Spinal canal stenosis

4) Avascular necrosis in SLE is restricted to the hip

- a. true
- b. false

5) The WHO defined osteopenia as:

- a. T score -3.5 or less
- b. T score between -1 and -2.5
- c. T score -2.5 or less
- d. T score of -1



1 c 2 true 3 c 4 false 5 b

**6. Old Book Questions****1. What is Polyarthritis?**

- a. Arthritis affecting two to four joints during the first six months of disease.
- b. Arthritis affecting more than five joints during the first six months of disease.
- c. Arthritis affecting one joint during the first six months of disease.
- d. Arthritis affecting more than one joint during the first six months of disease.

2. Heberden nodes, in hand osteoarthritis, are present on

- a. Metatarsophalangeal joints.
- b. Proximal interphalangeal joints.
- c. Distal interphalangeal joints.
- d. Thumb base joint

3. A Common cause of acute monoarthritis

- a. gouty arthritis
- b. Rheumatoid arthritis
- c. psoriatic arthritis
- d. systemic sclerosis

4. What is the most common Extra-articular Manifestations of Rheumatoid arthritis?

- a. Sicca syndrome
- b. Rheumatoid nodules
- c. Pericarditis and pleuritis
- d. Interstitial lung fibrosis

5. What is the earliest pathogenic finding in Rheumatoid arthritis affected joint?

- a. cartilage degeneration
- b. subchondral bone cyst
- c. synovitis
- d. capsulitis

6. What is the commonest pattern of arthritis in Rheumatoid arthritis

- a. Symmetrical small joint arthritis.
- b. Symmetrical large joint arthritis.
- c. Asymmetrical small joint arthritis.
- d. Asymmetrical large joint arthritis

7. Which one of the following drugs is one of the DMARDs?

- a. Allopurinol.
- b. NSAIDs.
- c. Methotrexate.
- d. Colchicine.

8. Which one of the following domains Is Not Involved in ACR/EULAR criteria for RA diagnosis?

- a. Joint distribution
- b. Acute phase reactant
- c. Serology
- d. Radiological assessment

**9. Which statement is true about rheumatoid factor (RF)?**

- a. It is specific for Rheumatoid arthritis.
- b. It is not specific for Rheumatoid arthritis and occurs in many other conditions
- c. It occurs in 100% of Rheumatoid arthritis patients.
- d. not included in the diagnostic criteria of Rheumatoid arthritis

10. Which statement is true about the Anti CCP antibody?

- a. It is Specific for Rheumatoid arthritis.
- b. It is not specific for Rheumatoid arthritis and occurs in other diseases.
- c. It occurs in 100% of Rheumatoid arthritis patient
- d. It is not included in the diagnostic criteria of Rheumatoid arthritis

11. Which statement is true about Enthesitis related arthritis category of Juvenile idiopathic arthritis?

- a. It occurs more frequently in females.
- b. It involves joints of the hand and feet.
- c. It involves joints of the upper limbs.
- d. Enthesitis occurs mostly in planter fascia and tendoachilis

12. What is the most frequent form of glucocorticoids used in Juvenile idiopathic arthritis patients?

- a. Bridge therapy
- b. Pulse therapy
- c. Intra articular injection
- d. Oral forms

13. The commonest extra articular manifestation occurring in Spondyloarthropathy (SpA) is

- a. Acute anterior uveitis
- b. Heart block
- c. Aortic insufficiency
- d. Restrictive lung disease

14. Dactylitis occurs most frequently in among all categories of Spondarthropathy

- a. ankylosing spondylitis
- b. enteropathic arthritis
- c. undifferentiated Spondarthropathy
- d. psoriatic arthritis

15. The axial affection of Spondarthropathy (SpA) is best treated with

- a. NSAIDs
- b. Steroids
- c. Conventional DMARDS
- d. Biologics

16. The main pathogenic finding in osteoarthritis is:

- a. cartilage degeneration
- b. subchondral bone cyst
- c. synovitis
- d. capsulitis

**17. The WHO defined osteoporosis as**

- a. T score -2.5 or less
- b. T score of 1
- c. T score between 1 and -2.5
- d. T score -3.5 or less

18. What is the most accurate site for bone mineral density measurement?

- a. lumbar spine
- b. distal end radius
- c. distal end tibia
- d. femoral neck

19. What is the best diagnostic modality for osteoporosis?

- a. Dual energy X-ray absorptiometry (DEXA) scan
- b. Plain radiography
- c. Ultrasonography
- d. MRI

20. Which drug is used during the acute attack of gouty arthritis?

- a. Uricosuric drugs
- b. Allopurinol
- c. Colchicine
- d. Methotrexate

21. Which statement is true about SLE?

- a. It typically affects women between 16 years and 55 years
- b. It typically affects men between 16 years and 55 years
- c. It typically affects pediatric age group
- d. It equally affects women and men between 16 years and 55 years

22. Which of the following autoantibodies are the most specific for SLE

- a. anti-dsDNA and anti-Sm
- b. ANA
- c. Anti CCP
- d. RF

23. Which immunosuppressant drugs are most commonly used to induce remission in lupus nephritis?

- a. MTX
- b. Cyclosporine
- c. Leflunomide
- d. Mycophenolate Mofetil and Cyclophosphamide

1 b 2 c 3 a 4 b 5 c 6 a 7 c 8 d 9 b 10 a
11 d 12 c 13 a 14 d 15 a 16 a 17 a 18 d 19 a 20 c
21 a 22 a 23 d

**7. New Book Questions (MCQs)****1) What is oligo arthritis?**

- a. Arthritis affecting two to four joints during the first six months of disease.
- b. Arthritis affecting more than four joints during the first six months of disease.
- c. Arthritis affecting one joint during the first six months of disease.
- d. Arthritis affecting more than one joint during the first six months of disease.

2) A Common cause of acute monoarthritis is

- a. gouty arthritis
- b. Rheumatoid arthritis
- c. psoriatic arthritis
- d. systemic sclerosis

3) What is the commonest pattern of arthritis in Rheumatoid arthritis?

- a. Asymmetrical small joint arthritis.
- b. Symmetrical large joint arthritis.
- c. symmetrical small joint arthritis.
- d. Asymmetrical large joint arthritis

4) Which statement is true about ACPA?

- a. It is specific for Rheumatoid arthritis.
- b. It is not specific for Rheumatoid arthritis and occurs in many other conditions.
- c. It occurs in 100% of Rheumatoid arthritis patients.
- d. not included in the diagnostic criteria of Rheumatoid arthritis

5) Which disease of the following is one of Spondarthropathy (SpA) subgroups?

- a. Rheumatoid arthritis
- b. Systemic sclerosis
- c. Sarcoidosis
- d. Psoriatic arthritis

6) The main pathogenic finding in RA is;

- a. cartilage degeneration
- b. subchondral bone cyst
- c. synovitis
- d. Capsulitis

7) Gout is disease caused by the deposition ofin joints

- a. monosodium urate crystals
- b. calcium oxalate crystals
- c. calcium pyrophosphate dihydrate crystals
- d. Calcium phosphate crystals

8) Which immune suppressant drugs are most commonly used to induce remission in lupus nephritis?

- a. MTX
- b. Cyclosporine
- c. Leflunomide
- d. Mycophenolate Mofetil

9) Which of the following autoantibodies are the most specific for SLE?

- a. anti-Sm
- b. ANA
- c. Anti CCP
- d. RF

10) What is the main diagnostic modality for osteoarthritis?

- a. Dual energy X-ray absorptiometry (DEXA) scan
- b. Plain radiography
- c. Ultrasonography
- d. MRI