

OBJECTIVE: Plasmid DNA Isolation from *E. coli* culture (Manual Method).

Chemical required: Glucose, Tris-Cl, EDTA, potassium acetate, glacial acetic acid, NaOH, Sodium dodecyl sulfate, Iso-propanol, RNase (10mg/ml, Qiagen) ddH₂O/MilliQ. LE Agarose (EEO: 0.09-0.13)

Plastic ware/ glassware: Measuring cylinder, autoclaved micro centrifuge tubes, 1.0 ml tips, 0.02ml-0.2 ml tips, enzyme tips, 100ml/250 ml flasks, reagent bottle, 0.2 µ syringe filter.

Equipment required: Horizontal electrophoresis unit, power supply, casting tray, combs, minispin (centrifuge), pH meter.

Recipe for solutions:

1. Solution I (100 ml)

50 mM glucose	1.8 ml of 50 % glucose
25 mM Tris HCl (pH 8.0)	2.5 ml of 1M Tris-HCl (pH 8.0)
10 mM EDTA (pH 8.0)	2.0 ml of 0.5 M EDTA (pH 8.0)
	93.7 ml of H ₂ O (autoclaved)

2. Solution II (100 ml)

(This should be freshly prepared)

0.2 N NaOH	10 ml 2N NaOH
1.0 % SDS	10 ml 10% SDS
	80 ml of H ₂ O (autoclaved)

3. Solution III (100 ml)

Potassium acetate (5M, pH 4.8)	60 ml 5M Potassium Acetate
Glacial acetic acid	11.5 ml Glacial acetic acid
Sterile distilled water	28.5 ml of H ₂ O (autoclaved)

4. TE (100 ml)

10 mM Tris- HCL (pH 8.0)	1 ml 1 M Tris-HCl (pH 8.0)
1 mM EDTA-sodium salt	0.5 ml 0.5 M EDTA (pH 8.0)

98.5 ml of H₂O (autoclaved)

PROTOCOL.

Inoculate a single colony of *E. coli* DH5 α in 5ml LB and incubate at 37°C, 180 rpm overnight.

Harvest cells from overnight grown culture (\sim O.D_{600nm} 1.0- 3.0) at 12000 rpm for 2min.

Wash the pellet with MilliQ.

Resuspend pellet in 100 μ l of Solution I and keep on ice for 5 min.

Add 200 μ l solution II and thoroughly mix by inverting tube 4-5 times.

Add 150 μ l ice-cold solution III and mix by inverting tube 4-5 times, incubate for 5- 10 min on ice.

Centrifuge mixture at 12000 rpm for 15 min. Collect supernatant in separate tube and add 0.8 volume of isopropanol followed by incubation on ice for 10-15 min.

To get plasmid DNA centrifuge mixture at 12000 rpm for 5 min.

Wash the pellet with 1 ml 70% ethanol followed by centrifugation at 12000 rpm for 2 min.

Air dry the pellet and dissolve in 40 μ l of TE buffer.

Add RNase 100 μ g/ml to final concentration and incubate for 1 hr at 37°C.

Check the plasmid DNA on 1 % agarose gel.