


Manipulation of DNA- 3

Artificial Restriction Enzymes


ZINC FINGER NUCLEASES



Basic structure and design of a zinc finger nuclease (ZFN). ZFNs are created by joining a DNA-binding region to the catalytic domain of the nonspecific FokI endonuclease. Zinc fingers are a protein motif capable of DNA binding, whose sequence specificity can be predetermined. Each zinc finger, illustrated by an individual circle, recognizes 3-4 nucleotides, and, by assembling three or four suitable zinc finger motifs, a sequence-specific DNA-binding domain can be created. FokI nuclease activity requires dimerization, and so the customized ZFNs function in pairs. As shown, the zinc finger-binding domain brings two FokI units together in the right orientation over the target sequence; this induces FokI dimerization and target sequence cleavage.

Transcription Activator-like Effector Nucleases (TALENs)

Transcription activator-like effectors (TALEs) are DNA-binding domains that can be linked together modularly and fused with nuclease domains to create TALE nucleases, or TALENs



Pioneers of revolutionary CRISPR gene editing win chemistry Nobel

Emmanuelle Charpentier and Jennifer Doudna share the award for developing the precise genome-editing technology.

Heidi Ledford & Ewen Callaway



Jennifer Doudna and Emmanuelle Charpentier share the 2020 Nobel chemistry prize for their discovery of a game-changing gene-editing technique. Credit: Alexander Heinel/Picture Alliance/DPA

RELATED ARTICLES

The quiet revolutionary: How the co-discovery of CRISPR explosively changed Emmanuelle Charpentier's life


Genome-editing revolution: My whirlwind year with CRISPR

CRISPR, the disruptor

The unsung heroes of CRISPR

<https://www.nature.com/articles/d41586-020-02765-9>

CRISPR (clustered regularly interspaced short palindromic repeats) is a family of DNA sequences found in the genomes of prokaryotic organisms such as bacteria and archaea. These sequences are derived from DNA fragments of bacteriophages that had previously infected the prokaryote



The CRISPR loci include Cas genes, a leader sequence, and several spacer sequences (in green) derived from engineered or foreign DNA that are separated by short repeat sequences (in black). Individual mature CRISPR RNA (crRNAs) are created by processing of the CRISPR transcript. Cas proteins and the crRNA make an effector complex that recognizes a target nucleic acid sequence via complementary base pairing to the crRNA. Cleavage of the target sequence occurs and is followed by DNA repair by the endogenous cellular repair machinery.

For Zinc Finger Nuclease

<https://www.youtube.com/watch?v=iGmq1O6mly0>

For transcriptional activator-like effector nucleases (TALENs)

<https://www.thermofisher.com/in/en/home/life-science/genome-editing/geneart-tals.html>


For CRISPR

<https://www.youtube.com/watch?v=MnYppmstxls>

<https://www.livescience.com/58790-crispr-explained.html>


For ethical concerns read this news

<https://www.sciencemag.org/news/2019/12/chinese-scientist-who-produced-genetically-altered-babies-sentenced-3-years-jail>




Meganuclease

20-40 bp/Enzyme




ZFN

3 bp/Finger



TALEN

1 bp/Module



CRISPR-Cas

1 bp/Base

