Teleinformatics Engineering Department, Federal University of Ceará

Introduction to Gaussian Processes

Filipe P. de Farias, IC filipepfarias@fisica.ufc.br

October 23, 2018





The Gaussian Processes are the widely used stochastic processes for modeling dependent data observed over time, space or even time and space. Here, we'll iniciate our study with a Probability and Random Process Theory Review taking some point to base our journey, going through Linear Regression and finally the Gaussian Processes.

The material here presented isn't sufficient to guide you over basic probability, so it's recommended to have some knowledge, once we'll just take a simple review.

Outline

- Probability and Random Process Theory Review
 - 1.1 Basic Concepts of Probability Theory
 - 1.2 The Gaussian distribution
 - 1.3 Independency of two random variables
- 2 Linear Regression
 - 2.1 Basic Concepts of Curve Fitting
- 2.2 Bayesian Curve Fitting

Theory Review

Probability and Random Process



Specifying Random Experiments

Let's start with the sentence

Sentence

A random experiment is specified by stating an experimental procedure and a set of one or more measurements or observations.



The Gaussian distribution

The **Gaussian distribution** is defined as

$$\mathcal{N}(x|\mu,\sigma^2) = \frac{1}{(2\pi\sigma^2)^{1/2}} \exp\left\{-\frac{1}{2\sigma^2}(x-\mu)^2\right\}$$
 (1)





Sentence

X and **Y** are independent random variables if *any* event A_1 defined in terms of X is independent of *any* event A_2 defined in terms of Y



Sentence

X and **Y** are independent random variables if *any* event A_1 defined in terms of X is independent of *any* event A_2 defined in terms of Y

The sentence above is equivalent to say mathematically that



Sentence

X and **Y** are independent random variables if *any* event A_1 defined in terms of X is independent of *any* event A_2 defined in terms of Y

The sentence above is equivalent to say mathematically that

$$P[X \text{ in } A_1, Y \text{ in } A_2] = P[X \text{ in } A_1]P[Y \text{ in } A_2]$$
(2)



Sentence

X and **Y** are independent random variables if *any* event A_1 defined in terms of X is independent of *any* event A_2 defined in terms of Y

The sentence above is equivalent to say mathematically that

$$P[X \text{ in } A_1, Y \text{ in } A_2] = P[X \text{ in } A_1]P[Y \text{ in } A_2]$$
(2)

that means in other words that if X and Y are independent discrete random variables, then the **joint probability mass function (pmf)** is equal to the product of the marginal pmf's.

Linear Regression



So, we'll start to look the regression with a statistical approach. To encourage you, let's take the sentence.



So, we'll start to look the regression with a statistical approach. To encourage you, let's take the sentence.

Sentence

If we could update the **regression weights** as we acquire some new values of the experiment?



Let's take a look again at the Bayes Theorem



Let's take a look again at the Bayes Theorem

Bayes Theorem

the weights probability
$$p(\mathbf{w}|\mathcal{D}) = \underbrace{\frac{p(\mathcal{D}|\mathbf{w})}{p(\mathbf{w})}}_{\text{the data probability}}$$
(3)



Let's take a look again at the Bayes Theorem

Bayes Theorem

the weights probability
$$p(\mathbf{w}|\mathcal{D}) = \underbrace{\frac{p(\mathcal{D}|\mathbf{w})}{p(\mathbf{w})}}_{\text{the data probability}}$$
(3)

So, if we have the probability of the data, we'll could estimate the future weights.



Let's take a look again at the Bayes Theorem

Bayes Theorem

the weights probability
$$p(\mathbf{w}|\mathcal{D}) = \underbrace{\frac{p(\mathcal{D}|\mathbf{w})}{p(\mathbf{w})}}_{\text{the data probability}}$$
(3)

So, if we have the probability of the data, we'll could estimate the future weights.

But, how?



Taking some steps back, let's re-visit the **Curve Fitting**. There, the strategy was minimize the error function.



Taking some steps back, let's re-visit the **Curve Fitting**. There, the strategy was minimize the error function.

Now we'll try to view the same problem with a *probabilistic perspective*. We're trying to make predictions for the target value \mathbf{t} given some new values of x.



Taking some steps back, let's re-visit the **Curve Fitting**. There, the strategy was minimize the error function.

Now we'll try to view the same problem with a *probabilistic perspective*. We're trying to make predictions for the target value **t** given some new values of *x*.

A good ideia is to express our target values ${\bf t}$ in terms of **gaussians distributions** with the mean equals to $y(x,{\bf w})$.






Figure: Schematic of the polynomial function $y(x, \mathbf{w})$ and the gaussian distribution p.



By the 12, we assume the relation

(5)

(6)



By the 12, we assume the relation

$$p(t|x, \mathbf{w}, \beta) = \mathcal{N}(t|y(x, \mathbf{w}), \beta^{-1})$$
(4)



By the 12, we assume the relation

$$p(t|x, \mathbf{w}, \beta) = \mathcal{N}(t|y(x, \mathbf{w}), \beta^{-1})$$
(4)

and then, assume that the training data $\{x, t\}$ is independent and identically distributed (i.i.d.) and put on **product form**, i.e. the joint probability is

$$p(\mathbf{t}|\mathbf{x},\mathbf{w},\beta) = \mathcal{N}(t_0|y(x_1,\mathbf{w}),\beta^{-1}) \cap \mathcal{N}(t_n|y(x_0,\mathbf{w}),\beta^{-1}) \dots \cap \mathcal{N}(t_n|y(x_0,\mathbf{w}),\beta^{-1})$$
(5)

(6)



By the 12, we assume the relation

$$p(t|x, \mathbf{w}, \beta) = \mathcal{N}(t|y(x, \mathbf{w}), \beta^{-1})$$
(4)

and then, assume that the training data $\{x, t\}$ is independent and identically distributed (i.i.d.) and put on **product form**, i.e. the joint probability is

$$p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \beta) = \mathcal{N}(t_0|y(x_1, \mathbf{w}), \beta^{-1}) \cap \mathcal{N}(t_n|y(x_0, \mathbf{w}), \beta^{-1}) \dots \cap \mathcal{N}(t_n|y(x_0, \mathbf{w}), \beta^{-1})$$
(5)
$$= \prod_{n=1}^{N} \mathcal{N}(t_n|y(x_n, \mathbf{w}), \beta^{-1})$$
(6)

regarding that $\beta^{-1} = \sigma^2$.



And we'll have a function to maximize if we apply the logarithm function to p, so



And we'll have a function to maximize if we apply the logarithm function to p, so

$$\ln \left(p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \beta) \right) = \sum_{n=1}^{N} \ln \left(\mathcal{N}(t_n|y(x_n, \mathbf{w}), \beta^{-1}) \right)$$
 (7)



And we'll have a function to maximize if we apply the logarithm function to p, so

$$\ln\left(p(\mathbf{t}|\mathbf{x},\mathbf{w},\beta)\right) = \sum_{n=1}^{N} \ln\left(\mathcal{N}(t_n|y(x_n,\mathbf{w}),\beta^{-1})\right)$$
(7)

Applying the Gaussian distribution (see 1) will result

$$\ln\left(p(\mathbf{t}|\mathbf{x},\mathbf{w},\beta)\right) = -\frac{\beta}{2} \sum_{n=1}^{N} \left\{y(x_n,\mathbf{w}-t_n)\right\}^2 + \frac{N}{2} \ln(\beta) - \frac{N}{2} \ln(2\pi)$$
 (8)



And taking the derivatives with respect to β to minimize the error

(9)

(11)



And taking the derivatives with respect to β to minimize the error

$$\frac{\partial}{\partial \beta} \ln \left(p(\mathbf{t} | \mathbf{x}, \mathbf{w}, \beta) \right) = 0 \tag{9}$$

(11)



And taking the derivatives with respect to β to minimize the error

$$\frac{\partial}{\partial \beta} \ln \left(p(\mathbf{t} | \mathbf{x}, \mathbf{w}, \beta) \right) = 0 \tag{9}$$

$$\frac{\partial}{\partial \beta} \ln \left(p(\mathbf{t} | \mathbf{x}, \mathbf{w}, \beta) \right) = 0$$

$$-\frac{1}{2} \sum_{n=1}^{N} \left\{ y(x_n, \mathbf{w} - t_n) \right\}^2 + \frac{N}{2} \frac{1}{\beta} = 0$$
(10)

(11)



And taking the derivatives with respect to β to minimize the error

$$\frac{\partial}{\partial \beta} \ln \left(p(\mathbf{t} | \mathbf{x}, \mathbf{w}, \beta) \right) = 0 \tag{9}$$

$$-\frac{1}{2}\sum_{n=1}^{N}\left\{y(x_n,\mathbf{w}-t_n)\right\}^2 + \frac{N}{2}\frac{1}{\beta} = 0$$
 (10)

$$\frac{1}{N} \sum_{n=1}^{N} \{ y(x_n, \mathbf{w} - t_n) \}^2 = \frac{1}{\beta_{ML}}$$
 (11)

Where β_{ML} is the maximum likelihood.